Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

SKU: 9780134439020-SOLUTIONS

CHAPTER 2 LIMITS AND CONTINUITY

2.1 RATES OF CHANGE AND TANGENTS TO CURVES

1. (a) $\frac{\Delta f}{\Delta x} = \frac{f(3)-f(2)}{3-2} = \frac{28-9}{1} = 19$ (b) $\frac{\Delta f}{\Delta x} = \frac{f(1)-f(-1)}{1-(-1)} = \frac{2-0}{2} = 1$

2. (a) $\frac{\Delta g}{\Delta x} = \frac{g(3)-g(1)}{3-1} = \frac{3-(-1)}{2} = 2$ (b) $\frac{\Delta g}{\Delta x} = \frac{g(4)-g(-2)}{4-(-2)} = \frac{8-8}{6} = 0$

3. (a) $\frac{\Delta h}{\Delta t} = \frac{h\left(\frac{3\pi}{4}\right)-h\left(\frac{\pi}{4}\right)}{\frac{3\pi}{4}-\frac{\pi}{4}} = \frac{-1-1}{\frac{\pi}{2}} = -\frac{4}{\pi}$ (b) $\frac{\Delta h}{\Delta t} = \frac{h\left(\frac{\pi}{2}\right)-h\left(\frac{\pi}{6}\right)}{\frac{\pi}{2}-\frac{\pi}{6}} = \frac{0-\sqrt{3}}{\frac{\pi}{3}} = \frac{-3\sqrt{3}}{\pi}$

4. (a) $\frac{\Delta g}{\Delta t} = \frac{g(\pi)-g(0)}{\pi-0} = \frac{(2-1)-(2+1)}{\pi-0} = -\frac{2}{\pi}$ (b) $\frac{\Delta g}{\Delta t} = \frac{g(\pi)-g(-\pi)}{\pi-(-\pi)} = \frac{(2-1)-(2-1)}{2\pi} = 0$

5. $\frac{\Delta R}{\Delta \theta} = \frac{R(2)-R(0)}{2-0} = \frac{\sqrt{8+1}-\sqrt{1}}{2} = \frac{3-1}{2} = 1$

6. $\frac{\Delta P}{\Delta \theta} = \frac{P(2)-P(1)}{2-1} = \frac{(8-16+10)-(1-4+5)}{1} = 2-2 = 0$

7. (a) $\frac{\Delta y}{\Delta x} = \frac{((2+h)^2-5)-(2^2-5)}{h} = \frac{4+4h+h^2-5+1}{h} = \frac{4h+h^2}{h} = 4+h$. As $h \to 0, 4+h \to 4 \Rightarrow$ at $P(2,-1)$ the slope is 4.
(b) $y-(-1) = 4(x-2) \Rightarrow y+1 = 4x-8 \Rightarrow y = 4x-9$

8. (a) $\frac{\Delta y}{\Delta x} = \frac{(7-(2+h)^2)-(7-2^2)}{h} = \frac{7-4-4h-h^2-3}{h} = \frac{-4h-h^2}{h} = -4-h$. As $h \to 0, -4-h \to -4 \Rightarrow$ at $P(2,3)$ the slope is -4.
(b) $y-3 = (-4)(x-2) \Rightarrow y-3 = -4x+8 \Rightarrow y = -4x+11$

9. (a) $\frac{\Delta y}{\Delta x} = \frac{((2+h)^2-2(2+h)-3)-(2^2-2(2)-3)}{h} = \frac{4+4h+h^2-4-2h-3-(-3)}{h} = \frac{2h+h^2}{h} = 2+h$. As $h \to 0, 2+h \to 2 \Rightarrow$ at $P(2,-3)$ the slope is 2.
(b) $y-(-3) = 2(x-2) \Rightarrow y+3 = 2x-4 \Rightarrow y = 2x-7$.

10. (a) $\frac{\Delta y}{\Delta x} = \frac{((1+h)^2-4(1+h))-(1^2-4(1))}{h} = \frac{1+2h+h^2-4-4h-(-3)}{h} = \frac{h^2-2h}{h} = h-2$. As $h \to 0, h-2 \to -2 \Rightarrow$ at $P(1,-3)$ the slope is -2.
(b) $y-(-3) = (-2)(x-1) \Rightarrow y+3 = -2x+2 \Rightarrow y = -2x-1$.

11. (a) $\frac{\Delta y}{\Delta x} = \frac{(2+h)^3-2^3}{h} = \frac{8+12h+4h^2+h^3-8}{h} = \frac{12h+4h^2+h^3}{h} = 12+4h+h^2$. As $h \to 0, 12+4h+h^2 \to 12, \Rightarrow$ at $P(2,8)$ the slope is 12.
(b) $y-8 = 12(x-2) \Rightarrow y-8 = 12x-24 \Rightarrow y = 12x-16$.

12. (a) $\frac{\Delta y}{\Delta x} = \frac{2-(1+h)^3-(2-1^3)}{h} = \frac{2-1-3h-3h^2-h^3-1}{h} = \frac{-3h-3h^2-h^3}{h} = -3-3h-h^2$. As $h \to 0, -3-3h-h^2 \to -3, \Rightarrow$ at $P(1,1)$ the slope is -3.
(b) $y-1 = (-3)(x-1) \Rightarrow y-1 = -3x+3 \Rightarrow y = -3x+4$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

13. (a) $\frac{\Delta y}{\Delta x} = \frac{(1+h)^3 - 12(1+h) - (1^3 - 12(1))}{h} = \frac{1+3h+3h^2+h^3-12-12h-(-11)}{h} = \frac{-9h+3h^2+h^3}{h} = -9+3h+h^2.$
As $h \to 0$, $-9+3h+h^2 \to -9 \Rightarrow$ at $P(1,-11)$ the slope is -9.
(b) $y-(-11) = (-9)(x-1) \Rightarrow y+11 = -9x+9 \Rightarrow y = -9x-2.$

14. (a) $\frac{\Delta y}{\Delta x} = \frac{(2+h)^3 - 3(2+h)^2 + 4 - (2^3 - 3(2)^2 + 4)}{h} = \frac{8+12h+6h^2+h^3-12-12h-3h^2+4-0}{h} = \frac{3h^2+h^3}{h} = 3h+h^2.$
As $h \to 0$, $3h+h^2 \to 0 \Rightarrow$ at $P(2,0)$ the slope is 0.
(b) $y-0 = 0(x-2) \Rightarrow y = 0.$

15. (a) $\frac{\Delta y}{\Delta x} = \frac{\frac{1}{-2+h} - \frac{1}{-2}}{h} = \frac{2+(-2+h)}{2(-2+h)} \cdot \frac{1}{h} = \frac{1}{2(-2+h)}.$
As $h \to 0$, $\frac{1}{2(-2+h)} \to \frac{-1}{4}$, $\Rightarrow$ at $P\left(-2, \frac{-1}{2}\right)$ the slope is $\frac{-1}{4}$.
(b) $y-\left(\frac{-1}{2}\right) = \frac{-1}{4}(x-(-2)) \Rightarrow y+\frac{1}{2} = \frac{-1}{4}x - \frac{1}{2} \Rightarrow y = \frac{-1}{4}x - 1$

16. (a) $\frac{\Delta y}{\Delta x} = \frac{\frac{(4+h)}{2-(4+h)} - \frac{4}{2-4}}{h} = \left(\frac{4+h}{-2-h} + \frac{2}{1}\right) \cdot \frac{1}{h} = \frac{4+h+2(-2-h)}{-2-h} \cdot \frac{1}{h} = \frac{-1}{-2-h} = \frac{1}{2+h}.$
As $h \to 0$, $\frac{1}{2+h} \to \frac{1}{2}$, $\Rightarrow$ at $P(4,-2)$ the slope is $\frac{1}{2}$.
(b) $y-(-2) = \frac{1}{2}(x-4) \Rightarrow y+2 = \frac{1}{2}x - 2 \Rightarrow y = \frac{1}{2}x - 4$

17. (a) $\frac{\Delta y}{\Delta x} = \frac{\sqrt{4+h}-\sqrt{4}}{h} = \frac{\sqrt{4+h}-2}{h} \cdot \frac{\sqrt{4+h}+2}{\sqrt{4+h}+2} = \frac{(4+h)-4}{h(\sqrt{4+h}+2)} = \frac{1}{\sqrt{4+h}+2}.$
As $h \to 0$, $\frac{1}{\sqrt{4+h}+2} \to \frac{1}{\sqrt{4}+2} = \frac{1}{4}$, $\Rightarrow$ at $P(4,2)$ the slope is $\frac{1}{4}$.
(b) $y-2 = \frac{1}{4}(x-4) \Rightarrow y-2 = \frac{1}{4}x - 1 \Rightarrow y = \frac{1}{4}x + 1$

18. (a) $\frac{\Delta y}{\Delta x} = \frac{\sqrt{7-(-2+h)}-\sqrt{7-(-2)}}{h} = \frac{\sqrt{9-h}-3}{h} = \frac{\sqrt{9-h}-3}{h} \cdot \frac{\sqrt{9-h}+3}{\sqrt{9-h}+3} = \frac{(9-h)-9}{h(\sqrt{9-h}+3)} = \frac{-1}{\sqrt{9-h}+3}.$
As $h \to 0$, $\frac{-1}{\sqrt{9-h}+3} \to \frac{-1}{\sqrt{9}+3} = \frac{-1}{6}$, $\Rightarrow$ at $P(-2,3)$ the slope is $\frac{-1}{6}$.
(b) $y-3 = \frac{-1}{6}(x-(-2)) \Rightarrow y-3 = \frac{-1}{6}x - \frac{1}{3} \Rightarrow y = \frac{-1}{6}x + \frac{8}{3}$

19. (a)

Q	Slope of $PQ = \frac{\Delta p}{\Delta t}$
$Q_1(10,225)$	$\frac{650-225}{20-10} = 42.5$ m/sec
$Q_2(14,375)$	$\frac{650-375}{20-14} = 45.83$ m/sec
$Q_3(16.5,475)$	$\frac{650-475}{20-16.5} = 50.00$ m/sec
$Q_4(18,550)$	$\frac{650-550}{20-18} = 50.00$ m/sec

(b) At $t = 20$, the sportscar was traveling approximately 50 m/sec or 180 km/h.

20. (a)

Q	Slope of $PQ = \frac{\Delta p}{\Delta t}$
$Q_1(5,20)$	$\frac{80-20}{10-5} = 12$ m/sec
$Q_2(7,39)$	$\frac{80-39}{10-7} = 13.7$ m/sec
$Q_3(8.5,58)$	$\frac{80-58}{10-8.5} = 14.7$ m/sec
$Q_4(9.5,72)$	$\frac{80-72}{10-9.5} = 16$ m/sec

(b) Approximately 16 m/sec

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

21. (a)



(b) $\frac{\Delta p}{\Delta t} = \frac{174-62}{2014-2012} = \frac{112}{2} = 56$ thousand dollars per year

(c) The average rate of change from 2011 to 2012 is $\frac{\Delta p}{\Delta t} = \frac{62-27}{2012-2011} = 35$ thousand dollars per year.

The average rate of change from 2012 to 2013 is $\frac{\Delta p}{\Delta t} = \frac{111-62}{2013-2012} = 49$ thousand dollars per year.

So, the rate at which profits were changing in 2012 is approximately $\frac{1}{2}(35+49) = 42$ thousand dollars per year.

22. (a) $F(x) = (x+2)/(x-2)$

x	1.2	1.1	1.01	1.001	1.0001	1
$F(x)$	-4.0	$-3.\overline{4}$	$-3.0\overline{4}$	$-3.00\overline{4}$	$-3.000\overline{4}$	-3

$\frac{\Delta F}{\Delta x} = \frac{-4.0-(-3)}{1.2-1} = -5.0;$ $\frac{\Delta F}{\Delta x} = \frac{-3.\overline{4}-(-3)}{1.1-1} = -4.\overline{4};$

$\frac{\Delta F}{\Delta x} = \frac{-3.04-(-3)}{1.01-1} = -4.0\overline{4};$ $\frac{\Delta F}{\Delta x} = \frac{-3.004-(-3)}{1.001-1} = -4.00\overline{4};$

$\frac{\Delta F}{\Delta x} = \frac{-3.0004-(-3)}{1.0001-1} = -4.000\overline{4};$

(b) The rate of change of $F(x)$ at $x = 1$ is -4.

23. (a) $\frac{\Delta g}{\Delta x} = \frac{g(2)-g(1)}{2-1} = \frac{\sqrt{2}-1}{2-1} \approx 0.414213$ $\frac{\Delta g}{\Delta x} = \frac{g(1.5)-g(1)}{1.5-1} = \frac{\sqrt{1.5}-1}{0.5} \approx 0.449489$

$\frac{\Delta g}{\Delta x} = \frac{g(1+h)-g(1)}{(1+h)-1} = \frac{\sqrt{1+h}-1}{h}$

(b) $g(x) = \sqrt{x}$

$1+h$	1.1	1.01	1.001	1.0001	1.00001	1.000001
$\sqrt{1+h}$	1.04880	1.004987	1.0004998	1.0000499	1.000005	1.0000005
$\left(\sqrt{1+h}-1\right)/h$	0.4880	0.4987	0.4998	0.499	0.5	0.5

(c) The rate of change of $g(x)$ at $x = 1$ is 0.5.

(d) The calculator gives $\lim_{h\to 0} \frac{\sqrt{1+h}-1}{h} = \frac{1}{2}$.

24. (a) i) $\frac{f(3)-f(2)}{3-2} = \frac{\frac{1}{3}-\frac{1}{2}}{1} = \frac{\frac{-1}{6}}{1} = -\frac{1}{6}$

ii) $\frac{f(T)-f(2)}{T-2} = \frac{\frac{1}{T}-\frac{1}{2}}{T-2} = \frac{\frac{2}{2T}-\frac{T}{2T}}{T-2} = \frac{2-T}{2T(T-2)} = \frac{2-T}{-2T(2-T)} = -\frac{1}{2T}, T \neq 2$

(b)

T	2.1	2.01	2.001	2.0001	2.00001	2.000001
$f(T)$	0.476190	0.497512	0.499750	0.4999750	0.499997	0.499999
$(f(T)-f(2))/(T-2)$	-0.2381	-0.2488	-0.2500	-0.2500	-0.2500	-0.2500

(c) The table indicates the rate of change is -0.25 at $t = 2$.

(d) $\lim_{T\to 2}\left(\frac{1}{-2T}\right) = -\frac{1}{4}$

NOTE: Answers will vary in Exercises 25 and 26.

25. (a) $[0, 1]$: $\frac{\Delta s}{\Delta t} = \frac{15-0}{1-0} = 15$ mph; $[1, 2.5]$: $\frac{\Delta s}{\Delta t} = \frac{20-15}{2.5-1} = \frac{10}{3}$ mph; $[2.5, 3.5]$: $\frac{\Delta s}{\Delta t} = \frac{30-20}{3.5-2.5} = 10$ mph

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b) At $P\left(\frac{1}{2}, 7.5\right)$: Since the portion of the graph from $t = 0$ to $t = 1$ is nearly linear, the instantaneous rate of change will be almost the same as the average rate of change, thus the instantaneous speed at $t = \frac{1}{2}$ is $\frac{15-7.5}{1-0.5} = 15$ mi/hr. At $P(2, 20)$: Since the portion of the graph from $t = 2$ to $t = 2.5$ is nearly linear, the instantaneous rate of change will be nearly the same as the average rate of change, thus $v = \frac{20-20}{2.5-2} = 0$ mi/hr. For values of t less than 2, we have

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(1, 15)$	$\frac{15-20}{1-2} = 5$ mi/hr
$Q_2(1.5, 19)$	$\frac{19-20}{1.5-2} = 2$ mi/hr
$Q_3(1.9, 19.9)$	$\frac{19.9-20}{1.9-2} = 1$ mi/hr

Thus, it appears that the instantaneous speed at $t = 2$ is 0 mi/hr.
At $P(3, 22)$:

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(4, 35)$	$\frac{35-22}{4-3} = 13$ mi/hr
$Q_2(3.5, 30)$	$\frac{30-22}{3.5-3} = 16$ mi/hr
$Q_3(3.1, 23)$	$\frac{23-22}{3.1-3} = 10$ mi/hr

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(2, 20)$	$\frac{20-22}{2-3} = 2$ mi/hr
$Q_2(2.5, 20)$	$\frac{20-22}{2.5-3} = 4$ mi/hr
$Q_3(2.9, 21.6)$	$\frac{21.6-22}{2.9-3} = 4$ mi/hr

Thus, it appears that the instantaneous speed at $t = 3$ is about 7 mi/hr.

(c) It appears that the curve is increasing the fastest at $t = 3.5$. Thus for $P(3.5, 30)$

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(4, 35)$	$\frac{35-30}{4-3.5} = 10$ mi/hr
$Q_2(3.75, 34)$	$\frac{34-30}{3.75-3.5} = 16$ mi/hr
$Q_3(3.6, 32)$	$\frac{32-30}{3.6-3.5} = 20$ mi/hr

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(3, 22)$	$\frac{22-30}{3-3.5} = 16$ mi/hr
$Q_2(3.25, 25)$	$\frac{25-30}{3.25-3.5} = 20$ mi/hr
$Q_3(3.4, 28)$	$\frac{28-30}{3.4-3.5} = 20$ mi/hr

Thus, it appears that the instantaneous speed at $t = 3.5$ is about 20 mi/hr.

26. (a) $[0, 3]$: $\frac{\Delta A}{\Delta t} = \frac{10-15}{3-0} \approx -1.67 \frac{\text{gal}}{\text{day}}$; $[0, 5]$: $\frac{\Delta A}{\Delta t} = \frac{3.9-15}{5-0} \approx -2.2 \frac{\text{gal}}{\text{day}}$; $[7, 10]$: $\frac{\Delta A}{\Delta t} = \frac{0-1.4}{10-7} \approx -0.5 \frac{\text{gal}}{\text{day}}$

(b) At $P(1, 14)$:

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(2, 12.2)$	$\frac{12.2-14}{2-1} = -1.8$ gal/day
$Q_2(1.5, 13.2)$	$\frac{13.2-14}{1.5-1} = -1.6$ gal/day
$Q_3(1.1, 13.85)$	$\frac{13.85-14}{1.1-1} = -1.5$ gal/day

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(0, 15)$	$\frac{15-14}{0-1} = -1$ gal/day
$Q_2(0.5, 14.6)$	$\frac{14.6-14}{0.5-1} = -1.2$ gal/day
$Q_3(0.9, 14.86)$	$\frac{14.86-14}{0.9-1} = -1.4$ gal/day

Thus, it appears that the instantaneous rate of consumption at $t = 1$ is about -1.45 gal/day.
At $P(4, 6)$:

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(5, 3.9)$	$\frac{3.9-6}{5-4} = -2.1$ gal/day
$Q_2(4.5, 4.8)$	$\frac{4.8-6}{4.5-4} = -2.4$ gal/day
$Q_3(4.1, 5.7)$	$\frac{5.7-6}{4.1-4} = -3$ gal/day

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(3, 10)$	$\frac{10-6}{3-4} = -4$ gal/day
$Q_2(3.5, 7.8)$	$\frac{7.8-6}{3.5-4} = -3.6$ gal/day
$Q_3(3.9, 6.3)$	$\frac{6.3-6}{3.9-4} = -3$ gal/day

Thus, it appears that the instantaneous rate of consumption at $t = 1$ is -3 gal/day.

(*solution continues on next page*)

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

At $P(8, 1)$:

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(9, 0.5)$	$\frac{0.5-1}{9-8} = -0.5$ gal/day
$Q_2(8.5, 0.7)$	$\frac{0.7-1}{8.5-8} = -0.6$ gal/day
$Q_3(8.1, 0.95)$	$\frac{0.95-1}{8.1-8} = -0.5$ gal/day

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(7, 1.4)$	$\frac{1.4-1}{7-8} = -0.6$ gal/day
$Q_2(7.5, 1.3)$	$\frac{1.3-1}{7.5-8} = -0.6$ gal/day
$Q_3(7.9, 1.04)$	$\frac{1.04-1}{7.9-8} = -0.6$ gal/day

Thus, it appears that the instantaneous rate of consumption at $t = 1$ is -0.55 gal/day.

(c) It appears that the curve (the consumption) is decreasing the fastest at $t = 3.5$. Thus for $P(3.5, 7.8)$

Q	Slope of $PQ = \frac{\Delta A}{\Delta t}$
$Q_1(4.5, 4.8)$	$\frac{4.8-7.8}{4.5-3.5} = -3$ gal/day
$Q_2(4, 6)$	$\frac{6-7.8}{4-3.5} = -3.6$ gal/day
$Q_3(3.6, 7.4)$	$\frac{7.4-7.8}{3.6-3.5} = -4$ gal/day

Q	Slope of $PQ = \frac{\Delta s}{\Delta t}$
$Q_1(2.5, 11.2)$	$\frac{11.2-7.8}{2.5-3.5} = -3.4$ gal/day
$Q_2(3, 10)$	$\frac{10-7.8}{3-3.5} = -4.4$ gal/day
$Q_3(3.4, 8.2)$	$\frac{8.2-7.8}{3.4-3.5} = -4$ gal/day

Thus, it appears that the rate of consumption at $t = 3.5$ is about -4 gal/day.

2.2 LIMIT OF A FUNCTION AND LIMIT LAWS

1. (a) Does not exist. As x approaches 1 from the right, $g(x)$ approaches 0. As x approaches 1 from the left, $g(x)$ approaches 1. There is no single number L that all the values $g(x)$ get arbitrarily close to as $x \to 1$.
   (b) 1 (c) 0 (d) 0.5

2. (a) 0
   (b) -1
   (c) Does not exist. As t approaches 0 from the left, $f(t)$ approaches -1. As t approaches 0 from the right, $f(t)$ approaches 1. There is no single number L that $f(t)$ gets arbitrarily close to as $t \to 0$.
   (d) -1

3. (a) True (b) True (c) False
   (d) False (e) False (f) True
   (g) True (h) False (i) True
   (j) True (k) False

4. (a) False (b) False (c) True
   (d) True (e) True (f) True
   (g) False (h) True (i) False

5. $\lim_{x \to 0} \frac{x}{|x|}$ does not exist because $\frac{x}{|x|} = \frac{x}{x} = 1$ if $x > 0$ and $\frac{x}{|x|} = \frac{x}{-x} = -1$ if $x < 0$. As x approaches 0 from the left, $\frac{x}{|x|}$ approaches -1. As x approaches 0 from the right, $\frac{x}{|x|}$ approaches 1. There is no single number L that all the function values get arbitrarily close to as $x \to 0$.

6. As x approaches 1 from the left, the values of $\frac{1}{x-1}$ become increasingly large and negative. As x approaches 1 from the right, the values become increasingly large and positive. There is no number L that all the function values get arbitrarily close to as $x \to 1$, so $\lim_{x \to 1} \frac{1}{x-1}$ does not exist.

7. Nothing can be said about $f(x)$ because the existence of a limit as $x \to x_0$ does not depend on how the function is defined at x_0. In order for a limit to exist, $f(x)$ must be arbitrarily close to a single real number L when x is close enough to x_0. That is, the existence of a limit depends on the values of $f(x)$ for x near x_0, not on the definition of $f(x)$ at x_0 itself.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

8. Nothing can be said. In order for $\lim_{x\to 0} f(x)$ to exist, $f(x)$ must close to a single value for x near 0 regardless of the value $f(0)$ itself.

9. No, the definition does not require that f be defined at $x = 1$ in order for a limiting value to exist there. If $f(1)$ is defined, it can be any real number, so we can conclude nothing about $f(1)$ from $\lim_{x\to 1} f(x) = 5$.

10. No, because the existence of a limit depends on the values of $f(x)$ when x is near 1, not on $f(1)$ itself. If $\lim_{x\to 1} f(x)$ exists, its value may be some number other than $f(1) = 5$. We can conclude nothing about $\lim_{x\to 1} f(x)$, whether it exists or what its value is if it does exist, from knowing the value of $f(1)$ alone.

11. $\lim_{x\to -3} (x^2 - 13) = (-3)^2 - 13 = 9 - 13 = -4$

12. $\lim_{x\to 2} (-x^2 + 5x - 2) = -(2)^2 + 5(2) - 2 = -4 + 10 - 2 = 4$

13. $\lim_{t\to 6} 8(t-5)(t-7) = 8(6-5)(6-7) = -8$

14. $\lim_{x\to -2} (x^3 - 2x^2 + 4x + 8) = (-2)^3 - 2(-2)^2 + 4(-2) + 8 = -8 - 8 - 8 + 8 = -16$

15. $\lim_{x\to 2} \frac{2x+5}{11-x^3} = \frac{2(2)+5}{11-(2)^3} = \frac{9}{3} = 3$

16. $\lim_{t\to 2/3} (8-3s)(2s-1) = \left(8 - 5\left(\frac{2}{3}\right)\right)\left(2\left(\frac{2}{3}\right) - 1\right) = (8-2)\left(\left(\frac{4}{3}\right) - 1\right) = (6)\left(\frac{1}{3}\right) = 2$

17. $\lim_{x\to -1/2} 4x(3x+4)^2 = 4\left(-\frac{1}{2}\right)\left(3\left(-\frac{1}{2}\right) + 4\right)^2 = (-2)\left(-\frac{3}{2} + 4\right)^2 = (-2)\left(\frac{5}{2}\right)^2 = -\frac{25}{2}$

18. $\lim_{y\to 2} \frac{y+2}{y^2+5y+6} = \frac{2+2}{(2)^2+5(2)+6} = \frac{4}{4+10+6} = \frac{4}{20} = \frac{1}{5}$

19. $\lim_{y\to -3} (5-y)^{4/3} = [5-(-3)]^{4/3} = (8)^{4/3} = \left((8)^{1/3}\right)^4 = 2^4 = 16$

20. $\lim_{z\to 4} \sqrt{z^2 - 10} = \sqrt{4^2 - 10} = \sqrt{16 - 10} = \sqrt{6}$

21. $\lim_{h\to 0} \frac{3}{\sqrt{3h+1}+1} = \frac{3}{\sqrt{3(0)+1}+1} = \frac{3}{\sqrt{1}+1} = \frac{3}{2}$

22. $\lim_{h\to 0} \frac{\sqrt{5h+4}-2}{h} = \lim_{h\to 0} \frac{\sqrt{5h+4}-2}{h} \cdot \frac{\sqrt{5h+4}+2}{\sqrt{5h+4}+2} = \lim_{h\to 0} \frac{(5h+4)-4}{h\left(\sqrt{5h+4}+2\right)} = \lim_{h\to 0} \frac{5h}{h\left(\sqrt{5h+4}+2\right)} = \lim_{h\to 0} \frac{5}{\sqrt{5h+4}+2} = \frac{5}{\sqrt{4}+2} = \frac{5}{4}$

23. $\lim_{x\to 5} \frac{x-5}{x^2-25} = \lim_{x\to 5} \frac{x-5}{(x+5)(x-5)} = \lim_{x\to 5} \frac{1}{x+5} = \frac{1}{5+5} = \frac{1}{10}$

24. $\lim_{x\to -3} \frac{x+3}{x^2+4x+3} = \lim_{x\to -3} \frac{x+3}{(x+3)(x+1)} = \lim_{x\to -3} \frac{1}{x+1} = \frac{1}{-3+1} = -\frac{1}{2}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

25. $\lim\limits_{x\to -5}\dfrac{x^2+3x-10}{x+5}=\lim\limits_{x\to -5}\dfrac{(x+5)(x-2)}{x+5}=\lim\limits_{x\to -5}(x-2)=-5-2=-7$

26. $\lim\limits_{x\to 2}\dfrac{x^2-7x-10}{x-2}=\lim\limits_{x\to 2}\dfrac{(x-5)(x-2)}{x-2}=\lim\limits_{x\to 2}(x-5)=2-5=-3$

27. $\lim\limits_{t\to 1}\dfrac{t^2+t-2}{t^2-1}=\lim\limits_{t\to 1}\dfrac{(t+2)(t-1)}{(t-1)(t+1)}=\lim\limits_{t\to 1}\dfrac{t+2}{t+1}=\dfrac{1+2}{1+1}=\dfrac{3}{2}$

28. $\lim\limits_{t\to -1}\dfrac{t^2+3t+2}{t^2-t-2}=\lim\limits_{t\to -1}\dfrac{(t+2)(t+1)}{(t-2)(t+1)}=\lim\limits_{t\to -1}\dfrac{t+2}{t-2}=\dfrac{-1+2}{-1-2}=-\dfrac{1}{3}$

29. $\lim\limits_{x\to -2}\dfrac{-2x-4}{x^3+2x^2}=\lim\limits_{x\to -2}\dfrac{-2(x+2)}{x^2(x+2)}=\lim\limits_{x\to -2}\dfrac{-2}{x^2}=\dfrac{-2}{4}=-\dfrac{1}{2}$

30. $\lim\limits_{y\to 0}\dfrac{5y^3+8y^2}{3y^4-16y^2}=\lim\limits_{y\to 0}\dfrac{y^2(5y+8)}{y^2(3y^2-16)}=\lim\limits_{y\to 0}\dfrac{5y+8}{3y^2-16}=\dfrac{8}{-16}=-\dfrac{1}{2}$

31. $\lim\limits_{x\to 1}\dfrac{x^{-1}-1}{x-1}=\lim\limits_{x\to 1}\dfrac{\frac{1-x}{x}}{x-1}=\lim\limits_{x\to 1}\left(\dfrac{1-x}{x}\cdot\dfrac{1}{x-1}\right)=\lim\limits_{x\to 1}-\dfrac{1}{x}=-1$

32. $\lim\limits_{x\to 0}\dfrac{\frac{1}{x-1}+\frac{1}{x+1}}{x}=\lim\limits_{x\to 0}\dfrac{\frac{(x+1)+(x-1)}{(x-1)(x+1)}}{x}=\lim\limits_{x\to 0}\left(\dfrac{2x}{(x-1)(x+1)}\cdot\dfrac{1}{x}\right)=\lim\limits_{x\to 0}\dfrac{2}{(x-1)(x+1)}=\dfrac{2}{-1}=-2$

33. $\lim\limits_{u\to 1}\dfrac{u^4-1}{u^3-1}=\lim\limits_{u\to 1}\dfrac{(u^2+1)(u+1)(u-1)}{(u^2+u+1)(u-1)}=\lim\limits_{u\to 1}\dfrac{(u^2+1)(u+1)}{u^2+u+1}=\dfrac{(1+1)(1+1)}{1+1+1}=\dfrac{4}{3}$

34. $\lim\limits_{v\to 2}\dfrac{v^3-8}{v^4-16}=\lim\limits_{v\to 2}\dfrac{(v-2)(v^2+2v+4)}{(v-2)(v+2)(v^2+4)}=\lim\limits_{v\to 2}\dfrac{v^2+2v+4}{(v+2)(v^2+4)}=\dfrac{4+4+4}{(4)(8)}=\dfrac{12}{32}=\dfrac{3}{8}$

35. $\lim\limits_{x\to 9}\dfrac{\sqrt{x}-3}{x-9}=\lim\limits_{x\to 9}\dfrac{\sqrt{x}-3}{(\sqrt{x}-3)(\sqrt{x}+3)}=\lim\limits_{x\to 9}\dfrac{1}{\sqrt{x}+3}=\dfrac{1}{\sqrt{9}+3}=\dfrac{1}{6}$

36. $\lim\limits_{x\to 4}\dfrac{4x-x^2}{2-\sqrt{x}}=\lim\limits_{x\to 4}\dfrac{x(4-x)}{2-\sqrt{x}}=\lim\limits_{x\to 4}\dfrac{x(2+\sqrt{x})(2-\sqrt{x})}{2-\sqrt{x}}=\lim\limits_{x\to 4}x\left(2+\sqrt{x}\right)=4(2+2)=16$

37. $\lim\limits_{x\to 1}\dfrac{x-1}{\sqrt{x+3}-2}=\lim\limits_{x\to 1}\dfrac{(x-1)\left(\sqrt{x+3}+2\right)}{\left(\sqrt{x+3}-2\right)\left(\sqrt{x+3}+2\right)}=\lim\limits_{x\to 1}\dfrac{(x-1)\left(\sqrt{x+3}+2\right)}{(x+3)-4}=\lim\limits_{x\to 1}\left(\sqrt{x+3}+2\right)=\sqrt{4}+2=4$

38. $\lim\limits_{x\to -1}\dfrac{\sqrt{x^2+8}-3}{x+1}=\lim\limits_{x\to -1}\dfrac{\left(\sqrt{x^2+8}-3\right)\left(\sqrt{x^2+8}+3\right)}{(x+1)\left(\sqrt{x^2+8}+3\right)}=\lim\limits_{x\to -1}\dfrac{(x^2+8)-9}{(x+1)\left(\sqrt{x^2+8}+3\right)}=\lim\limits_{x\to -1}\dfrac{(x+1)(x-1)}{(x+1)\left(\sqrt{x^2+8}+3\right)}$

$=\lim\limits_{x\to -1}\dfrac{x-1}{\sqrt{x^2+8}+3}=\dfrac{-2}{3+3}=-\dfrac{1}{3}$

39. $\lim\limits_{x\to 2}\dfrac{\sqrt{x^2+12}-4}{x-2}=\lim\limits_{x\to 2}\dfrac{\left(\sqrt{x^2+12}-4\right)\left(\sqrt{x^2+12}+4\right)}{(x-2)\left(\sqrt{x^2+12}+4\right)}=\lim\limits_{x\to 2}\dfrac{(x^2+12)-16}{(x-2)\left(\sqrt{x^2+12}+4\right)}=\lim\limits_{x\to 2}\dfrac{(x-2)(x+2)}{(x-2)\left(\sqrt{x^2+12}+4\right)}$

$=\lim\limits_{x\to 2}\dfrac{x+2}{\sqrt{x^2+12}+4}=\dfrac{4}{\sqrt{16}+4}=\dfrac{1}{2}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

40. $\lim\limits_{x\to -2}\dfrac{x+2}{\sqrt{x^2+5}-3}=\lim\limits_{x\to -2}\dfrac{(x+2)\left(\sqrt{x^2+5}+3\right)}{\left(\sqrt{x^2+5}-3\right)\left(\sqrt{x^2+5}+3\right)}=\lim\limits_{x\to -2}\dfrac{(x+2)\left(\sqrt{x^2+5}+3\right)}{(x^2+5)-9}=\lim\limits_{x\to -2}\dfrac{(x+2)\left(\sqrt{x^2+5}+3\right)}{(x+2)(x-2)}$

$=\lim\limits_{x\to -2}\dfrac{\sqrt{x^2+5}+3}{x-2}=\dfrac{\sqrt{9}+3}{-4}=-\dfrac{3}{2}$

41. $\lim\limits_{x\to -3}\dfrac{2-\sqrt{x^2-5}}{x+3}=\lim\limits_{x\to -3}\dfrac{\left(2-\sqrt{x^2-5}\right)\left(2+\sqrt{x^2-5}\right)}{(x+3)\left(2+\sqrt{x^2-5}\right)}=\lim\limits_{x\to -3}\dfrac{4-(x^2-5)}{(x+3)\left(2+\sqrt{x^2-5}\right)}=\lim\limits_{x\to -3}\dfrac{9-x^2}{(x+3)\left(2+\sqrt{x^2-5}\right)}$

$=\lim\limits_{x\to -3}\dfrac{(3-x)(3+x)}{(x+3)\left(2+\sqrt{x^2-5}\right)}=\lim\limits_{x\to -3}\dfrac{3-x}{2+\sqrt{x^2-5}}=\dfrac{6}{2+\sqrt{4}}=\dfrac{3}{2}$

42. $\lim\limits_{x\to 4}\dfrac{4-x}{5-\sqrt{x^2+9}}=\lim\limits_{x\to 4}\dfrac{(4-x)\left(5+\sqrt{x^2+9}\right)}{\left(5-\sqrt{x^2+9}\right)\left(5+\sqrt{x^2+9}\right)}=\lim\limits_{x\to 4}\dfrac{(4-x)\left(5+\sqrt{x^2+9}\right)}{25-(x^2+9)}=\lim\limits_{x\to 4}\dfrac{(4-x)\left(5+\sqrt{x^2+9}\right)}{16-x^2}$

$=\lim\limits_{x\to 4}\dfrac{(4-x)\left(5+\sqrt{x^2+9}\right)}{(4-x)(4+x)}=\lim\limits_{x\to 4}\dfrac{5+\sqrt{x^2+9}}{4+x}=\dfrac{5+\sqrt{25}}{8}=\dfrac{5}{4}$

43. $\lim\limits_{x\to 0}(2\sin x-1)=2\sin 0-1=0-1=-1$

44. $\lim\limits_{x\to 0}\sin^2 x=\left(\lim\limits_{x\to 0}\sin x\right)^2=(\sin 0)^2=0^2=0$

45. $\lim\limits_{x\to 0}\sec x=\lim\limits_{x\to 0}\dfrac{1}{\cos x}=\dfrac{1}{\cos 0}=\dfrac{1}{1}=1$

46. $\lim\limits_{x\to 0}\tan x=\lim\limits_{x\to 0}\dfrac{\sin x}{\cos x}=\dfrac{\sin 0}{\cos 0}=\dfrac{0}{1}=0$

47. $\lim\limits_{x\to 0}\dfrac{1+x+\sin x}{3\cos x}=\dfrac{1+0+\sin 0}{3\cos 0}=\dfrac{1+0+0}{3}=\dfrac{1}{3}$

48. $\lim\limits_{x\to 0}(x^2-1)(2-\cos x)=(0^2-1)(2-\cos 0)=(-1)(2-1)=(-1)(1)=-1$

49. $\lim\limits_{x\to -\pi}\sqrt{x+4}\cos(x+\pi)=\lim\limits_{x\to -\pi}\sqrt{x+4}\cdot\lim\limits_{x\to -\pi}\cos(x+\pi)=\sqrt{-\pi+4}\cdot\cos 0=\sqrt{4-\pi}\cdot 1=\sqrt{4-\pi}$

50. $\lim\limits_{x\to 0}\sqrt{7+\sec^2 x}=\sqrt{\lim\limits_{x\to 0}(7+\sec^2 x)}=\sqrt{7+\lim\limits_{x\to 0}\sec^2 x}=\sqrt{7+\sec^2 0}=\sqrt{7+(1)^2}=2\sqrt{2}$

51. (a) quotient rule
(b) difference and power rules
(c) sum and constant multiple rules

52. (a) quotient rule
(b) power and product rules
(c) difference and constant multiple rules

53. (a) $\lim\limits_{x\to c}f(x)g(x)=\left[\lim\limits_{x\to c}f(x)\right]\left[\lim\limits_{x\to c}g(x)\right]=(5)(-2)=-10$

(b) $\lim\limits_{x\to c}2f(x)g(x)=2\left[\lim\limits_{x\to c}f(x)\right]\left[\lim\limits_{x\to c}g(x)\right]=2(5)(-2)=-20$

(c) $\lim\limits_{x\to c}[f(x)+3g(x)]=\lim\limits_{x\to c}f(x)+3\lim\limits_{x\to c}g(x)=5+3(-2)=-1$

(d) $\lim\limits_{x\to c}\dfrac{f(x)}{f(x)-g(x)}=\dfrac{\lim\limits_{x\to c}f(x)}{\lim\limits_{x\to c}f(x)-\lim\limits_{x\to c}g(x)}=\dfrac{5}{5-(-2)}=\dfrac{5}{7}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

54. (a) $\lim\limits_{x\to 4}[g(x)+3] = \lim\limits_{x\to 4} g(x) + \lim\limits_{x\to 4} 3 = -3+3 = 0$

(b) $\lim\limits_{x\to 4} xf(x) = \lim\limits_{x\to 4} x \cdot \lim\limits_{x\to 4} f(x) = (4)(0) = 0$

(c) $\lim\limits_{x\to 4}[g(x)]^2 = \left[\lim\limits_{x\to 4} g(x)\right]^2 = [-3]^2 = 9$

(d) $\lim\limits_{x\to 4} \dfrac{g(x)}{f(x)-1} = \dfrac{\lim\limits_{x\to 4} g(x)}{\lim\limits_{x\to 4} f(x) - \lim\limits_{x\to 4} 1} = \dfrac{-3}{0-1} = 3$

55. (a) $\lim\limits_{x\to b}[f(x)+g(x)] = \lim\limits_{x\to b} f(x) + \lim\limits_{x\to b} g(x) = 7+(-3) = 4$

(b) $\lim\limits_{x\to b} f(x)\cdot g(x) = \left[\lim\limits_{x\to b} f(x)\right]\left[\lim\limits_{x\to b} g(x)\right] = (7)(-3) = -21$

(c) $\lim\limits_{x\to b} 4g(x) = \left[\lim\limits_{x\to b} 4\right]\left[\lim\limits_{x\to b} g(x)\right] = (4)(-3) = -12$

(d) $\lim\limits_{x\to b} f(x)/g(x) = \lim\limits_{x\to b} f(x) / \lim\limits_{x\to b} g(x) = \dfrac{7}{-3} = -\dfrac{7}{3}$

56. (a) $\lim\limits_{x\to -2}[p(x)+r(x)+s(x)] = \lim\limits_{x\to -2} p(x) + \lim\limits_{x\to -2} r(x) + \lim\limits_{x\to -2} s(x) = 4+0+(-3) = 1$

(b) $\lim\limits_{x\to -2} p(x)\cdot r(x)\cdot s(x) = \left[\lim\limits_{x\to -2} p(x)\right]\left[\lim\limits_{x\to -2} r(x)\right]\left[\lim\limits_{x\to -2} s(x)\right] = (4)(0)(-3) = 0$

(c) $\lim\limits_{x\to -2}[-4p(x)+5r(x)]/s(x) = \left[-4\lim\limits_{x\to -2} p(x) + 5\lim\limits_{x\to -2} r(x)\right] \Big/ \lim\limits_{x\to -2} s(x) = [-4(4)+5(0)]/-3 = \dfrac{16}{3}$

57. $\lim\limits_{h\to 0} \dfrac{(1+h)^2-1^2}{h} = \lim\limits_{h\to 0} \dfrac{1+2h+h^2-1}{h} = \lim\limits_{h\to 0} \dfrac{h(2+h)}{h} = \lim\limits_{h\to 0} (2+h) = 2$

58. $\lim\limits_{h\to 0} \dfrac{(-2+h)^2-(-2)^2}{h} = \lim\limits_{h\to 0} \dfrac{4-4h+h^2-4}{h} = \lim\limits_{h\to 0} \dfrac{h(h-4)}{h} = \lim\limits_{h\to 0} (h-4) = -4$

59. $\lim\limits_{h\to 0} \dfrac{[3(2+h)-4]-[3(2)-4]}{h} = \lim\limits_{h\to 0} \dfrac{3h}{h} = 3$

60. $\lim\limits_{h\to 0} \dfrac{\left(\frac{1}{-2+h}\right)-\left(\frac{1}{-2}\right)}{h} = \lim\limits_{h\to 0} \dfrac{\frac{-2}{-2+h}-1}{-2h} = \lim\limits_{h\to 0} \dfrac{-2-(-2+h)}{-2h(-2+h)} = \lim\limits_{h\to 0} \dfrac{-h}{h(4-2h)} = -\dfrac{1}{4}$

61. $\lim\limits_{h\to 0} \dfrac{\sqrt{7+h}-\sqrt{7}}{h} = \lim\limits_{h\to 0} \dfrac{\left(\sqrt{7+h}-\sqrt{7}\right)\left(\sqrt{7+h}+\sqrt{7}\right)}{h\left(\sqrt{7+h}+\sqrt{7}\right)} = \lim\limits_{h\to 0} \dfrac{(7+h)-7}{h\left(\sqrt{7+h}+\sqrt{7}\right)} = \lim\limits_{h\to 0} \dfrac{h}{h\left(\sqrt{7+h}+\sqrt{7}\right)} = \lim\limits_{h\to 0} \dfrac{1}{\sqrt{7+h}+\sqrt{7}} = \dfrac{1}{2\sqrt{7}}$

62. $\lim\limits_{h\to 0} \dfrac{\sqrt{3(0+h)+1}-\sqrt{3(0)+1}}{h} = \lim\limits_{h\to 0} \dfrac{\left(\sqrt{3h+1}-1\right)\left(\sqrt{3h+1}+1\right)}{h\left(\sqrt{3h+1}+1\right)} = \lim\limits_{h\to 0} \dfrac{(3h+1)-1}{h\left(\sqrt{3h+1}+1\right)} = \lim\limits_{h\to 0} \dfrac{3h}{h\left(\sqrt{3h+1}+1\right)} = \lim\limits_{h\to 0} \dfrac{3}{\sqrt{3h+1}+1} = \dfrac{3}{2}$

63. $\lim\limits_{x\to 0} \sqrt{5-2x^2} = \sqrt{5-2(0)^2} = \sqrt{5}$ and $\lim\limits_{x\to 0} \sqrt{5-x^2} = \sqrt{5-(0)^2} = \sqrt{5}$; by the sandwich theorem, $\lim\limits_{x\to 0} f(x) = \sqrt{5}$

64. $\lim\limits_{x\to 0} (2-x^2) = 2-0 = 2$ and $\lim\limits_{x\to 0} 2\cos x = 2(1) = 2$; by the sandwich theorem, $\lim\limits_{x\to 0} g(x) = 2$

65. (a) $\lim\limits_{x\to 0}\left(1-\dfrac{x^2}{6}\right) = 1-\dfrac{0}{6} = 1$ and $\lim\limits_{x\to 0} 1 = 1$; by the sandwich theorem, $\lim\limits_{x\to 0} \dfrac{x\sin x}{2-2\cos x} = 1$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b) For $x \neq 0$, $y = (x \sin x)/(2 - 2\cos x)$ lies between the other two graphs in the figure, and the graphs converge as $x \to 0$.



66. (a) $\lim_{x\to 0}\left(\frac{1}{2} - \frac{x^2}{24}\right) = \lim_{x\to 0}\frac{1}{2} - \lim_{x\to 0}\frac{x^2}{24} = \frac{1}{2} - 0 = \frac{1}{2}$ and $\lim_{x\to 0}\frac{1}{2} = \frac{1}{2}$; by the sandwich theorem, $\lim_{x\to 0}\frac{1-\cos x}{x^2} = \frac{1}{2}$.

(b) For all $x \neq 0$, the graph of $f(x) = (1-\cos x)/x^2$ lies between the line $y = \frac{1}{2}$ and the parabola $y = \frac{1}{2} - x^2/24$, and the graphs converge as $x \to 0$.



67. (a) $f(x) = (x^2 - 9)/(x + 3)$

x	–3.1	–3.01	–3.001	–3.0001	–3.00001	–3.000001
$f(x)$	–6.1	–6.01	–6.001	–6.0001	–6.00001	–6.000001

x	–2.9	–2.99	–2.999	–2.9999	–2.99999	–2.999999
$f(x)$	–5.9	–5.99	–5.999	–5.9999	–5.99999	–5.999999

The estimate is $\lim_{x\to -3} f(x) = -6$.

(b)



(c) $f(x) = \frac{x^2-9}{x+3} = \frac{(x+3)(x-3)}{x+3} = x - 3$ if $x \neq -3$, and $\lim_{x\to -3}(x-3) = -3-3 = -6$.

68. (a) $g(x) = (x^2 - 2)/\left(x - \sqrt{2}\right)$

x	1.4	1.41	1.414	1.4142	1.41421	1.414213
$g(x)$	2.81421	2.82421	2.82821	2.828413	2.828423	2.828426

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b)



(c) $g(x)=\frac{x^2-2}{x-\sqrt{2}}=\frac{\left(x+\sqrt{2}\right)\left(x-\sqrt{2}\right)}{\left(x-\sqrt{2}\right)}=x+\sqrt{2}$ if $x\neq\sqrt{2}$, and $\lim\limits_{x\to\sqrt{2}}\left(x+\sqrt{2}\right)=\sqrt{2}+\sqrt{2}=2\sqrt{2}$.

69. (a) $G(x)=(x+6)/(x^2+4x-12)$

x	−5.9	−5.99	−5.999	−5.9999	−5.99999	−5.999999
$G(x)$	−.126582	−.1251564	−.1250156	−.1250015	−.1250001	−.1250000

x	−6.1	−6.01	−6.001	−6.0001	−6.00001	−6.000001
$G(x)$	−.123456	−.124843	−.124984	−.124998	−.124999	−.124999

(b)



(c) $G(x)=\frac{x+6}{(x^2+4x-12)}=\frac{x+6}{(x+6)(x-2)}=\frac{1}{x-2}$ if $x\neq-6$, and $\lim\limits_{x\to-6}\frac{1}{x-2}=\frac{1}{-6-2}=-\frac{1}{8}=-0.125$.

70. (a) $h(x)=(x^2-2x-3)/(x^2-4x+3)$

x	2.9	2.99	2.999	2.9999	2.99999	2.999999
$h(x)$	2.052631	2.005025	2.000500	2.000050	2.000005	2.0000005

x	3.1	3.01	3.001	3.0001	3.00001	3.000001
$h(x)$	1.952380	1.995024	1.999500	1.999950	1.999995	1.999999

(b)



(c) $h(x)=\frac{x^2-2x-3}{x^2-4x+3}=\frac{(x-3)(x+1)}{(x-3)(x-1)}=\frac{x+1}{x-1}$ if $x\neq3$, and $\lim\limits_{x\to3}\frac{x+1}{x-1}=\frac{3+1}{3-1}=\frac{4}{2}=2$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

71. (a) $f(x) = (x^2 - 1)/(|x| - 1)$

x	-1.1	-1.01	-1.001	-1.0001	-1.00001	-1.000001
$f(x)$	2.1	2.01	2.001	2.0001	2.00001	2.000001

x	$-.9$	$-.99$	$-.999$	$-.9999$	$-.99999$	$-.999999$
$f(x)$	1.9	1.99	1.999	1.9999	1.99999	1.999999

(b)



(c) $f(x) = \dfrac{x^2-1}{|x|-1} = \begin{cases} \dfrac{(x+1)(x-1)}{x-1} = x+1, & x \geq 0 \text{ and } x \neq 1 \\ \dfrac{(x+1)(x-1)}{-(x+1)} = 1-x, & x < 0 \text{ and } x \neq -1 \end{cases}$, and $\lim\limits_{x \to -1} (1-x) = 1-(-1) = 2.$

72. (a) $F(x) = (x^2 + 3x + 2)/(2 - |x|)$

x	-2.1	-2.01	-2.001	-2.0001	-2.00001	-2.000001
$F(x)$	-1.1	-1.01	-1.001	-1.0001	-1.00001	-1.000001

x	-1.9	-1.99	-1.999	-1.9999	-1.99999	-1.999999
$F(x)$	$-.9$	$-.99$	$-.999$	$-.9999$	$-.99999$	$-.999999$

(b)



(c) $F(x) = \dfrac{x^2+3x+2}{2-|x|} = \begin{cases} \dfrac{(x+2)(x+1)}{2-x}, & x \geq 0 \\ \dfrac{(x+2)(x+1)}{2+x} = x+1, & x < 0 \text{ and } x \neq -2 \end{cases}$, and $\lim\limits_{x \to -2} (x+1) = -2+1 = -1.$

73. (a) $g(\theta) = (\sin\theta)/\theta$

θ	.1	.01	.001	.0001	.00001	.000001
$g(\theta)$	.998334	.999983	.999999	.999999	.999999	.999999

θ	$-.1$	$-.01$	$-.001$	$-.0001$	$-.00001$	$-.000001$
$g(\theta)$	.998334	.999983	.999999	.999999	.999999	.999999

$\lim\limits_{\theta \to 0} g(\theta) = 1$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b)



NOT TO SCALE

74. (a) $G(t) = (1 - \cos t)/t^2$

t	.1	.01	.001	.0001	.00001	.000001
$G(t)$	.499583	.499995	.499999	.5	.5	.5

t	−.1	−.01	−.001	−.0001	−.00001	−.000001
$G(t)$	.499583	.499995	.499999	.5	.5	.5

$\lim_{t\to 0} G(t) = 0.5$

(b)



Graph is NOT TO SCALE

75. (a) $f(x) = x^{1/(1-x)}$

x	.9	.99	.999	.9999	.99999	.999999
$f(x)$	.348678	.366032	.367695	.367861	.367877	.367879

x	1.1	1.01	1.001	1.0001	1.00001	1.000001
$f(x)$	.385543	.369711	.368063	.367897	.367881	.367878

$\lim_{x\to 1} f(x) \approx 0.36788$

(b)



Graph is NOT TO SCALE. Also, the intersection of the axes is not the origin: the axes intersect at the point (1, 2.71820).

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

76. (a) $f(x) = (3^x - 1)/x$

x	.1	.01	.001	.0001	.00001	.000001
$f(x)$	1.161231	1.104669	1.099215	1.098672	1.098618	1.098612

x	−.1	−.01	−.001	−.0001	−.00001	−.000001
$f(x)$	1.040415	1.092599	1.098009	1.098551	1.098606	1.098611

$\lim_{x\to 1} f(x) \approx 1.0986$

(b)



77. $\lim_{x\to c} f(x)$ exists at those points c where $\lim_{x\to c} x^4 = \lim_{x\to c} x^2$. Thus, $c^4 = c^2 \Rightarrow c^2(1-c^2) = 0 \Rightarrow c = 0, 1,$ or -1.
Moreover, $\lim_{x\to 0} f(x) = \lim_{x\to 0} x^2 = 0$ and $\lim_{x\to -1} f(x) = \lim_{x\to 1} f(x) = 1$.

78. Nothing can be concluded about the values of f, g, and h at $x = 2$. Yes, $f(2)$ could be 0. Since the conditions of the sandwich theorem are satisfied, $\lim_{x\to 2} f(x) = -5 \neq 0$.

79. $1 = \lim_{x\to 4} \frac{f(x)-5}{x-2} = \frac{\lim_{x\to 4} f(x) - \lim_{x\to 4} 5}{\lim_{x\to 4} x - \lim_{x\to 4} 2} = \frac{\lim_{x\to 4} f(x) - 5}{4-2} \Rightarrow \lim_{x\to 4} f(x) - 5 = 2(1) \Rightarrow \lim_{x\to 4} f(x) = 2 + 5 = 7.$

80. (a) $1 = \lim_{x\to -2} \frac{f(x)}{x^2} = \frac{\lim_{x\to -2} f(x)}{\lim_{x\to -2} x^2} = \frac{\lim_{x\to -2} f(x)}{4} \Rightarrow \lim_{x\to -2} f(x) = 4.$

(b) $1 = \lim_{x\to -2} \frac{f(x)}{x^2} = \left[\lim_{x\to -2} \frac{f(x)}{x}\right]\left[\lim_{x\to -2} \frac{1}{x}\right] = \left[\lim_{x\to -2} \frac{f(x)}{x}\right]\left(\frac{1}{-2}\right) \Rightarrow \lim_{x\to -2} \frac{f(x)}{x} = -2.$

81. (a) $0 = 3\cdot 0 = \left[\lim_{x\to 2} \frac{f(x)-5}{x-2}\right]\left[\lim_{x\to 2} (x-2)\right] = \lim_{x\to 2}\left[\left(\frac{f(x)-5}{x-2}\right)(x-2)\right] = \lim_{x\to 2} [f(x)-5]$
$= \lim_{x\to 2} f(x) - 5 \Rightarrow \lim_{x\to 2} f(x) = 5.$

(b) $0 = 4\cdot 0 = \left[\lim_{x\to 2} \frac{f(x)-5}{x-2}\right]\left[\lim_{x\to 2} (x-2)\right] \Rightarrow \lim_{x\to 2} f(x) = 5$ as in part (a).

82. (a) $0 = 1\cdot 0 = \left[\lim_{x\to 0} \frac{f(x)}{x^2}\right]\left[\lim_{x\to 0} x\right]^2 = \left[\lim_{x\to 0} \frac{f(x)}{x^2}\right]\left[\lim_{x\to 0} x^2\right] = \lim_{x\to 0}\left[\frac{f(x)}{x^2}\cdot x^2\right] = \lim_{x\to 0} f(x).$
That is, $\lim_{x\to 0} f(x) = 0$.

(b) $0 = 1\cdot 0 = \left[\lim_{x\to 0} \frac{f(x)}{x^2}\right]\left[\lim_{x\to 0} x\right] = \lim_{x\to 0}\left[\frac{f(x)}{x^2}\cdot x\right] = \lim_{x\to 0} \frac{f(x)}{x}$. That is, $\lim_{x\to 0} \frac{f(x)}{x} = 0.$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

83. (a) $\lim_{x\to 0} x\sin\frac{1}{x} = 0$



(b) $-1 \le \sin\frac{1}{x} \le 1$ for $x \ne 0$:

$x > 0 \Rightarrow -x \le x\sin\frac{1}{x} \le x \Rightarrow \lim_{x\to 0} x\sin\frac{1}{x} = 0$ by the sandwich theorem;

$x < 0 \Rightarrow -x \ge x\sin\frac{1}{x} \ge x \Rightarrow \lim_{x\to 0} x\sin\frac{1}{x} = 0$ by the sandwich theorem.

84. (a) $\lim_{x\to 0} x^2\cos\left(\frac{1}{x^3}\right) = 0$



(b) $-1 \le \cos\left(\frac{1}{x^3}\right) \le 1$ for $x \ne 0 \Rightarrow -x^2 \le x^2\cos\left(\frac{1}{x^3}\right) \le x^2 \Rightarrow \lim_{x\to 0} x^2\cos\left(\frac{1}{x^3}\right) = 0$ by the sandwich theorem since $\lim_{x\to 0} x^2 = 0$.

85–90. Example CAS commands:

Maple:

```
f := x -> (x^4 – 16)/(x – 2);
x0 := 2;
plot( f(x), x = x0-1..x0+1, color = black,
      title = "Section 2.2, #85(a)" );
limit( f(x), x = x0 );
```

In Exercise 87, note that the standard cube root, x^(1/3), is not defined for x<0 in many CASs. This can be overcome in Maple by entering the function as f := x -> (surd(x+1, 3) – 1)/x.

Mathematica: (assigned function and values for x0 and h may vary)

```
Clear[f, x]
f[x_]:=(x^3 – x^2 – 5x – 3)/(x + 1)^2
x0= –1; h = 0.1;
Plot[f[x], {x, x0 – h, x0 + h}]
Limit[f[x], x → x0]
```

2.3 THE PRECISE DEFINITION OF A LIMIT

1. 

Step 1: $|x-5| < \delta \Rightarrow -\delta < x-5 < \delta \Rightarrow -\delta+5 < x < \delta+5$

Step 2: $\delta+5 = 7 \Rightarrow \delta = 2$, or $-\delta+5 = 1 \Rightarrow \delta = 4$.

The value of δ which assures $|x-5| < \delta \Rightarrow 1 < x < 7$ is the smaller value, $\delta = 2$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

2.

Step 1: $|x-2|<\delta \Rightarrow -\delta<x-2<\delta \Rightarrow -\delta+2<x<\delta+2$

Step 2: $-\delta+2=1 \Rightarrow \delta=1$, or $\delta+2=7 \Rightarrow \delta=5$.

The value of δ which assures $|x-2|<\delta \Rightarrow 1<x<7$ is the smaller value, $\delta=1$.

3.

Step 1: $|x-(-3)|<\delta \Rightarrow -\delta<x+3<\delta \Rightarrow -\delta-3<x<\delta-3$

Step 2: $-\delta-3=-\frac{7}{2} \Rightarrow \delta=\frac{1}{2}$, or $\delta-3=-\frac{1}{2} \Rightarrow \delta=\frac{5}{2}$.

The value of δ which assures $|x-(-3)|<\delta \Rightarrow -\frac{7}{2}<x<-\frac{1}{2}$ is the smaller value, $\delta=\frac{1}{2}$.

4.



Step 1: $\left|x-\left(-\frac{3}{2}\right)\right|<\delta \Rightarrow -\delta<x+\frac{3}{2}<\delta \Rightarrow -\delta-\frac{3}{2}<x<\delta-\frac{3}{2}$

Step 2: $-\delta-\frac{3}{2}=-\frac{7}{2} \Rightarrow \delta=2$, or $\delta-\frac{3}{2}=-\frac{1}{2} \Rightarrow \delta=1$.

The value of δ which assures $\left|x-\left(-\frac{3}{2}\right)\right|<\delta \Rightarrow -\frac{7}{2}<x<-\frac{1}{2}$ is the smaller value, $\delta=1$.

5.



Step 1: $\left|x-\frac{1}{2}\right|<\delta \Rightarrow -\delta<x-\frac{1}{2}<\delta \Rightarrow -\delta+\frac{1}{2}<x<\delta+\frac{1}{2}$

Step 2: $-\delta+\frac{1}{2}=\frac{4}{9} \Rightarrow \delta=\frac{1}{18}$, or $\delta+\frac{1}{2}=\frac{4}{7} \Rightarrow \delta=\frac{1}{14}$.

The value of δ which assures $\left|x-\frac{1}{2}\right|<\delta \Rightarrow \frac{4}{9}<x<\frac{4}{7}$ is the smaller value, $\delta=\frac{1}{18}$.

6.



Step 1: $|x-3|<\delta \Rightarrow -\delta<x-3<\delta \Rightarrow -\delta+3<x<\delta+3$

Step 2: $-\delta+3=2.7591 \Rightarrow \delta=0.2409$, or $\delta+3=3.2391 \Rightarrow \delta=0.2391$.

The value of δ which assures $|x-3|<\delta \Rightarrow 2.7591<x<3.2391$ is the smaller value, $\delta=0.2391$.

7. Step 1: $|x-5|<\delta \Rightarrow -\delta<x-5<\delta \Rightarrow -\delta+5<x<\delta+5$

Step 2: From the graph, $-\delta+5=4.9 \Rightarrow \delta=0.1$, or $\delta+5=5.1 \Rightarrow \delta=0.1$; thus $\delta=0.1$ in either case.

8. Step 1: $|x-(-3)|<\delta \Rightarrow -\delta<x+3<\delta \Rightarrow -\delta-3<x<\delta-3$

Step 2: From the graph, $-\delta-3=-3.1 \Rightarrow \delta=0.1$, or $\delta-3=-2.9 \Rightarrow \delta=0.1$; thus $\delta=0.1$.

9. Step 1: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow -\delta+1<x<\delta+1$

Step 2: From the graph, $-\delta+1=\frac{9}{16} \Rightarrow \delta=\frac{7}{16}$, or $\delta+1=\frac{25}{16} \Rightarrow \delta=\frac{9}{16}$; thus $\delta=\frac{7}{16}$.

10. Step 1: $|x-3|<\delta \Rightarrow -\delta<x-3<\delta \Rightarrow -\delta+3<x<\delta+3$

Step 2: From the graph, $-\delta+3=2.61 \Rightarrow \delta=0.39$, or $\delta+3=3.41 \Rightarrow \delta=0.41$; thus $\delta=0.39$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

11. Step 1: $|x-2|<\delta \Rightarrow -\delta<x-2<\delta \Rightarrow -\delta+2<x<\delta+2$
Step 2: From the graph, $-\delta+2=\sqrt{3} \Rightarrow \delta=2-\sqrt{3}\approx 0.2679$, or $\delta+2=\sqrt{5} \Rightarrow \delta=\sqrt{5}-2\approx 0.2361$; thus $\delta=\sqrt{5}-2$.

12. Step 1: $|x-(-1)|<\delta \Rightarrow -\delta<x+1<\delta \Rightarrow -\delta-1<x<\delta-1$
Step 2: From the graph, $-\delta-1=-\frac{\sqrt{5}}{2} \Rightarrow \delta=\frac{\sqrt{5}-2}{2}\approx 0.118$ or $\delta-1=-\frac{\sqrt{3}}{2} \Rightarrow \delta=\frac{2-\sqrt{3}}{2}\approx 0.1340$; thus $\delta=\frac{\sqrt{5}-2}{2}$.

13. Step 1: $|x-(-1)|<\delta \Rightarrow -\delta<x+1<\delta \Rightarrow -\delta-1<x<\delta-1$
Step 2: From the graph, $-\delta-1=-\frac{16}{9} \Rightarrow \delta=\frac{7}{9}\approx 0.77$, or $\delta-1=-\frac{16}{25} \Rightarrow \frac{9}{25}=0.36$; thus $\delta=\frac{9}{25}=0.36$.

14. Step 1: $\left|x-\frac{1}{2}\right|<\delta \Rightarrow -\delta<x-\frac{1}{2}<\delta \Rightarrow -\delta+\frac{1}{2}<x<\delta+\frac{1}{2}$
Step 2: From the graph, $-\delta+\frac{1}{2}=\frac{1}{2.01} \Rightarrow \delta=\frac{1}{2}-\frac{1}{2.01}\approx 0.00248$, or $\delta+\frac{1}{2}=\frac{1}{1.99} \Rightarrow \delta=\frac{1}{1.99}-\frac{1}{2}\approx 0.00251$; thus $\delta=0.00248$.

15. Step 1: $|(x+1)-5|<0.01 \Rightarrow |x-4|<0.01 \Rightarrow -0.01<x-4<0.01 \Rightarrow 3.99<x<4.01$
Step 2: $|x-4|<\delta \Rightarrow -\delta<x-4<\delta \Rightarrow -\delta+4<x<\delta+4 \Rightarrow \delta=0.01$.

16. Step 1: $|(2x-2)-(-6)|<0.02 \Rightarrow |2x+4|<0.02 \Rightarrow -0.02<2x+4<0.02$
$\Rightarrow -4.02<2x<-3.98 \Rightarrow -2.01<x<-1.99$
Step 2: $|x-(-2)|<\delta \Rightarrow -\delta<x+2<\delta \Rightarrow -\delta-2<x<\delta-2 \Rightarrow \delta=0.01$.

17. Step 1: $|\sqrt{x+1}-1|<0.1 \Rightarrow -0.1<\sqrt{x+1}-1<0.1 \Rightarrow 0.9<\sqrt{x+1}<1.1 \Rightarrow 0.81<x+1<1.21$
$\Rightarrow -0.19<x<0.21$
Step 2: $|x-0|<\delta \Rightarrow -\delta<x<\delta$. Then, $-\delta=-0.19 \Rightarrow \delta=0.19$ or $\delta=0.21$; thus, $\delta=0.19$.

18. Step 1: $|\sqrt{x}-\frac{1}{2}|<0.1 \Rightarrow -0.1<\sqrt{x}-\frac{1}{2}<0.1 \Rightarrow 0.4<\sqrt{x}<0.6 \Rightarrow 0.16<x<0.36$
Step 2: $\left|x-\frac{1}{4}\right|<\delta \Rightarrow -\delta<x-\frac{1}{4}<\delta \Rightarrow$
Then $-\delta+\frac{1}{4}=0.16 \Rightarrow \delta=0.09$ or $\delta+\frac{1}{4}=0.36 \Rightarrow \delta=0.11$; thus $\delta=0.09$.

19. Step 1: $|\sqrt{19-x}-3|<1 \Rightarrow -1<\sqrt{19-x}-3<1 \Rightarrow 2<\sqrt{19-x}<4 \Rightarrow 4<19-x<16$
$\Rightarrow -4>x-19>-16 \Rightarrow 15>x>3$ or $3<x<15$
Step 2: $|x-10|<\delta \Rightarrow -\delta<x-10<\delta \Rightarrow -\delta+10<x<\delta+10$.
Then $-\delta+10=3 \Rightarrow \delta=7$, or $\delta+10=15 \Rightarrow \delta=5$; thus $\delta=5$.

20. Step 1: $|\sqrt{x-7}-4|<1 \Rightarrow -1<\sqrt{x-7}-4<1 \Rightarrow 3<\sqrt{x-7}<5 \Rightarrow 9<x-7<25 \Rightarrow 16<\mathrm{x}<32$
Step 2: $|x-23|<\delta \Rightarrow -\delta<x-23<\delta \Rightarrow -\delta+23<\mathrm{x}<\delta+23$.
Then $-\delta+23=16 \Rightarrow \delta=7$, or $\delta+23=32 \Rightarrow \delta=9$; thus $\delta=7$.

21. Step 1: $\left|\frac{1}{x}-\frac{1}{4}\right|<0.05 \Rightarrow -0.05<\frac{1}{x}-\frac{1}{4}<0.05 \Rightarrow 0.2<\frac{1}{x}<0.3 \Rightarrow \frac{10}{2}>x>\frac{10}{3}$ or $\frac{10}{3}<x<5$.
Step 2: $|x-4|<\delta \Rightarrow -\delta<x-4<\delta \Rightarrow -\delta+4<x<\delta+4$.
Then $-\delta+4=\frac{10}{3}$ or $\delta=\frac{2}{3}$, or $\delta+4=5$ or $\delta=1$; thus $\delta=\frac{2}{3}$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

22. Step 1: $|x^2-3|<0.1 \Rightarrow -0.1<x^2-3<0.1 \Rightarrow 2.9<x^2<3.1 \Rightarrow \sqrt{2.9}<x<\sqrt{3.1}$
Step 2: $|x-\sqrt{3}|<\delta \Rightarrow -\delta<x-\sqrt{3}<\delta \Rightarrow -\delta+\sqrt{3}<x<\delta+\sqrt{3}$.
Then $-\delta+\sqrt{3}=\sqrt{2.9} \Rightarrow \delta=\sqrt{3}-\sqrt{2.9}\approx 0.0291$, or $\delta+\sqrt{3}=\sqrt{3.1} \Rightarrow \delta=\sqrt{3.1}-\sqrt{3}\approx 0.0286$; thus $\delta=0.0286$

23. Step 1: $|x^2-4|<0.5 \Rightarrow -0.5<x^2-4<0.5 \Rightarrow 3.5<x^2<4.5 \Rightarrow \sqrt{3.5}<|x|<\sqrt{4.5} \Rightarrow -\sqrt{4.5}<x<-\sqrt{3.5}$, for x near -2.
Step 2: $|x-(-2)|<\delta \Rightarrow -\delta<x+2<\delta \Rightarrow -\delta-2<x<\delta-2$.
Then $-\delta-2=-\sqrt{4.5} \Rightarrow \delta=\sqrt{4.5}-2\approx 0.1213$, or $\delta-2=-\sqrt{3.5} \Rightarrow \delta=2-\sqrt{3.5}\approx 0.1292$; thus $\delta=\sqrt{4.5}-2\approx 0.12$.

24. Step 1: $\left|\frac{1}{x}-(-1)\right|<0.1 \Rightarrow -0.1<\frac{1}{x}+1<0.1 \Rightarrow -\frac{11}{10}<\frac{1}{x}<-\frac{9}{10} \Rightarrow -\frac{10}{11}>x>-\frac{10}{9}$ or $-\frac{10}{9}<x<-\frac{10}{11}$.
Step 2: $|x-(-1)|<\delta \Rightarrow -\delta<x+1<\delta \Rightarrow -\delta-1<x<\delta-1$.
Then $-\delta-1=-\frac{10}{9} \Rightarrow \delta=\frac{1}{9}$, or $\delta-1=-\frac{10}{11} \Rightarrow \delta=\frac{1}{11}$; thus $\delta=\frac{1}{11}$.

25. Step 1: $|(x^2-5)-11|<1 \Rightarrow |x^2-16|<1 \Rightarrow -1<x^2-16<1 \Rightarrow 15<x^2<17 \Rightarrow \sqrt{15}<x<\sqrt{17}$.
Step 2: $|x-4|<\delta \Rightarrow -\delta<x-4<\delta \Rightarrow -\delta+4<x<\delta+4$.
Then $-\delta+4=\sqrt{15} \Rightarrow \delta=4-\sqrt{15}\approx 0.1270$, or $\delta+4=\sqrt{17} \Rightarrow \delta=\sqrt{17}-4\approx 0.1231$; thus $\delta=\sqrt{17}-4\approx 0.12$.

26. Step 1: $\left|\frac{120}{x}-5\right|<1 \Rightarrow -1<\frac{120}{x}-5<1 \Rightarrow 4<\frac{120}{x}<6 \Rightarrow \frac{1}{4}>\frac{x}{120}>\frac{1}{6} \Rightarrow 30>x>20$ or $20<x<30$.
Step 2: $|x-24|<\delta \Rightarrow -\delta<x-24<\delta \Rightarrow -\delta+24<x<\delta+24$.
Then $-\delta+24=20 \Rightarrow \delta=4$, or $\delta+24=30 \Rightarrow \delta=6$; thus $\Rightarrow \delta=4$.

27. Step 1: $|mx-2m|<0.03 \Rightarrow -0.03<mx-2m<0.03 \Rightarrow -0.03+2m<mx<0.03+2m \Rightarrow 2-\frac{0.03}{m}<x<2+\frac{0.03}{m}$.
Step 2: $|x-2|<\delta \Rightarrow -\delta<x-2<\delta \Rightarrow -\delta+2<x<\delta+2$.
Then $-\delta+2=2-\frac{0.03}{m} \Rightarrow \delta=\frac{0.03}{m}$, or $\delta+2=2+\frac{0.03}{m} \Rightarrow \delta=\frac{0.03}{m}$. In either case, $\delta=\frac{0.03}{m}$.

28. Step 1: $|mx-3m|<c \Rightarrow -c<mx-3m<c \Rightarrow -c+3m<mx<c+3m \Rightarrow 3-\frac{c}{m}<x<3+\frac{c}{m}$
Step 2: $|x-3|<\delta \Rightarrow -\delta<x-3<\delta \Rightarrow -\delta+3<x<\delta+3$.
Then $-\delta+3=3-\frac{c}{m} \Rightarrow \delta=\frac{c}{m}$, or $\delta+3=3+\frac{c}{m} \Rightarrow \delta=\frac{c}{m}$. In either case, $\delta=\frac{c}{m}$.

29. Step 1: $\left|(mx+b)-\left(\frac{m}{2}+b\right)\right|<c \Rightarrow -c<mx-\frac{m}{2}<c \Rightarrow -c+\frac{m}{2}<mx<c+\frac{m}{2} \Rightarrow \frac{1}{2}-\frac{c}{m}<x<\frac{1}{2}+\frac{c}{m}$.
Step 2: $\left|x-\frac{1}{2}\right|<\delta \Rightarrow -\delta<x-\frac{1}{2}<\delta \Rightarrow -\delta+\frac{1}{2}<x<\delta+\frac{1}{2}$.
Then $-\delta+\frac{1}{2}=\frac{1}{2}-\frac{c}{m} \Rightarrow \delta=\frac{c}{m}$, or $\delta+\frac{1}{2}=\frac{1}{2}+\frac{c}{m} \Rightarrow \delta=\frac{c}{m}$. In either case, $\delta=\frac{c}{m}$.

30. Step 1: $|(mx+b)-(m+b)|<0.05 \Rightarrow -0.05<mx-m<0.05 \Rightarrow -0.05+m<mx<0.05+m$
$\Rightarrow 1-\frac{0.05}{m}<x<1+\frac{0.05}{m}$.
Step 2: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow -\delta+1<x<\delta+1$.
Then $-\delta+1=1-\frac{0.05}{m} \Rightarrow \delta=\frac{0.05}{m}$, or $\delta+1=1+\frac{0.05}{m} \Rightarrow \delta=\frac{0.05}{m}$. In either case, $\delta=\frac{0.05}{m}$.

31. $\lim_{x\to 3}(3-2x)=3-2(3)=-3$
Step 1: $|(3-2x)-(-3)|<0.02 \Rightarrow -0.02<6-2x<0.02 \Rightarrow -6.02<-2x<-5.98 \Rightarrow 3.01>x>2.99$ or $2.99<x<3.01$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

Step 2: $0 < |x-3| < \delta \Rightarrow -\delta < x-3 < \delta \Rightarrow -\delta+3 < x < \delta+3$.
Then $-\delta+3 = 2.99 \Rightarrow \delta = 0.01$, or $\delta+3 = 3.01 \Rightarrow \delta = 0.01$; thus $\delta = 0.01$.

32. $\lim_{x\to -1}(-3x-2) = (-3)(-1)-2 = 1$
Step 1: $|(-3x-2)-1| < 0.03 \Rightarrow -0.03 < -3x-3 < 0.03 \Rightarrow 0.01 > x+1 > -0.01 \Rightarrow -1.01 < x < -0.99$.
Step 2: $|x-(-1)| < \delta \Rightarrow -\delta < x+1 < \delta \Rightarrow -\delta-1 < x < \delta-1$.
Then $-\delta-1 = -1.01 \Rightarrow \delta = 0.01$, or $\delta-1 = -0.99 \Rightarrow \delta = 0.01$; thus $\delta = 0.01$.

33. $\lim_{x\to 2}\frac{x^2-4}{x-2} = \lim_{x\to 2}\frac{(x+2)(x-2)}{(x-2)} = \lim_{x\to 2}(x+2) = 2+2 = 4,\ x \neq 2$
Step 1: $\left|\left(\frac{x^2-4}{x-2}\right)-4\right| < 0.05 \Rightarrow -0.05 < \frac{(x+2)(x-2)}{(x-2)} - 4 < 0.05 \Rightarrow 3.95 < x+2 < 4.05,\ x \neq 2$
$\Rightarrow 1.95 < x < 2.05,\ x \neq 2$.
Step 2: $|x-2| < \delta \Rightarrow -\delta < x-2 < \delta \Rightarrow -\delta+2 < x < \delta+2$.
Then $-\delta+2 = 1.95 \Rightarrow \delta = 0.05$, or $\delta+2 = 2.05 \Rightarrow \delta = 0.05$; thus $\delta = 0.05$.

34. $\lim_{x\to -5}\frac{x^2+6x+5}{x+5} = \lim_{x\to -5}\frac{(x+5)(x+1)}{(x+5)} = \lim_{x\to -5}(x+1) = -4,\ x \neq -5$.
Step 1: $\left|\left(\frac{x^2+6x+5}{x+5}\right)-(-4)\right| < 0.05 \Rightarrow -0.05 < \frac{(x+5)(x+1)}{(x+5)} + 4 < 0.05 \Rightarrow -4.05 < x+1 < -3.95,\ \ x \neq -5$
$\Rightarrow -5.05 < x < -4.95,\ \ x \neq -5$.
Step 2: $|x-(-5)| < \delta \Rightarrow -\delta < x+5 < \delta \Rightarrow -\delta-5 < x < \delta-5$.
Then $-\delta-5 = -5.05 \Rightarrow \delta = 0.05$, or $\delta-5 = -4.95 \Rightarrow \delta = 0.05$; thus $\delta = 0.05$.

35. $\lim_{x\to -3}\sqrt{1-5x} = \sqrt{1-5(-3)} = \sqrt{16} = 4$
Step 1: $\left|\sqrt{1-5x}-4\right| < 0.5 \Rightarrow -0.5 < \sqrt{1-5x}-4 < 0.5 \Rightarrow 3.5 < \sqrt{1-5x} < 4.5 \Rightarrow 12.25 < 1-5x < 20.25$
$\Rightarrow 11.25 < -5x < 19.25 \Rightarrow -3.85 < x < 2.25$.
Step 2: $|x-(-3)| < \delta \Rightarrow -\delta < x+3 < \delta \Rightarrow -\delta-3 < x < \delta-3$.
Then $-\delta-3 = -3.85 \Rightarrow \delta = 0.85$, or $\delta-3 = -2.25 \Rightarrow 0.75$; thus $\delta = 0.75$.

36. $\lim_{x\to 2}\frac{4}{x} = \frac{4}{2} = 2$
Step 1: $\left|\frac{4}{x}-2\right| < 0.4 \Rightarrow -0.4 < \frac{4}{x}-2 < 0.4 \Rightarrow 1.6 < \frac{4}{x} < 2.4 \Rightarrow \frac{10}{16} > \frac{x}{4} > \frac{10}{24} \Rightarrow \frac{10}{4} > x > \frac{10}{6}$ or $\frac{5}{3} < x < \frac{5}{2}$.
Step 2: $|x-2| < \delta \Rightarrow -\delta < x-2 < \delta \Rightarrow -\delta+2 < x < \delta+2$.
Then $-\delta+2 = \frac{5}{3} \Rightarrow \delta = \frac{1}{3}$, or $\delta+2 = \frac{5}{2} \Rightarrow \delta = \frac{1}{2}$; thus $\delta = \frac{1}{3}$.

37. Step 1: $|(9-x)-5| < \epsilon \Rightarrow -\epsilon < 4-x < \epsilon \Rightarrow -\epsilon-4 < -x < \epsilon-4 \Rightarrow \epsilon+4 > x > 4-\epsilon \Rightarrow 4-\epsilon < x < 4+\epsilon$.
Step 2: $|x-4| < \delta \Rightarrow -\delta < x-4 < \delta \Rightarrow -\delta+4 < x < \delta+4$.
Then $-\delta+4 = -\epsilon+4 \Rightarrow \delta = \epsilon$, or $\delta+4 = \epsilon+4 \Rightarrow \delta = \epsilon$. Thus choose $\delta = \epsilon$.

38. Step 1: $|(3x-7)-2| < \epsilon \Rightarrow -\epsilon < 3x-9 < \epsilon \Rightarrow 9-\epsilon < 3x < 9+\epsilon \Rightarrow 3-\frac{\epsilon}{3} < x < 3+\frac{\epsilon}{3}$.
Step 2: $|x-3| < \delta \Rightarrow -\delta < x-3 < \delta \Rightarrow -\delta+3 < x < \delta+3$.
Then $-\delta+3 = 3-\frac{\epsilon}{3} \Rightarrow \delta = \frac{\epsilon}{3}$, or $\delta+3 = 3+\frac{\epsilon}{3} \Rightarrow \delta = \frac{\epsilon}{3}$. Thus choose $\delta = \frac{\epsilon}{3}$.

39. Step 1: $\left|\sqrt{x-5}-2\right| < \epsilon \Rightarrow -\epsilon < \sqrt{x-5}-2 < \epsilon \Rightarrow 2-\epsilon < \sqrt{x-5} < 2+\epsilon \Rightarrow (2-\epsilon)^2 < x-5 < (2+\epsilon)^2$
$\Rightarrow (2-\epsilon)^2+5 < x < (2+\epsilon)^2+5$.
Step 2: $|x-9| < \delta \Rightarrow -\delta < x-9 < \delta \Rightarrow -\delta+9 < x < \delta+9$.
Then $-\delta+9 = \epsilon^2-4\epsilon+9 \Rightarrow \delta = 4\epsilon-\epsilon^2$, or $\delta+9 = \epsilon^2+4\epsilon+9 \Rightarrow \delta = 4\epsilon+\epsilon^2$. Thus choose the smaller distance, $\delta = 4\epsilon-\epsilon^2$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

40. Step 1: $\left|\sqrt{4-x}-2\right|<\epsilon \Rightarrow -\epsilon<\sqrt{4-x}-2<\epsilon \Rightarrow 2-\epsilon<\sqrt{4-x}<2+\epsilon \Rightarrow (2-\epsilon)^2<4-x<(2+\epsilon)^2$
$\Rightarrow -(2+\epsilon)^2<x-4<-(2-\epsilon)^2 \Rightarrow -(2+\epsilon)^2+4<x<-(2-\epsilon)^2+4.$
Step 2: $|x-0|<\delta \Rightarrow -\delta<x<\delta.$
Then $-\delta=-(2+\epsilon)^2+4=-\epsilon^2-4\epsilon \Rightarrow \delta=4\epsilon+\epsilon^2$, or $\delta=-(2-\epsilon)^2+4=4\epsilon-\epsilon^2$. Thus choose the smaller distance, $\delta=4\epsilon-\epsilon^2$.

41. Step 1: For $x\neq 1$, $\left|x^2-1\right|<\epsilon \Rightarrow -\epsilon<x^2-1<\epsilon \Rightarrow 1-\epsilon<x^2<1+\epsilon \Rightarrow \sqrt{1-\epsilon}<|x|<\sqrt{1+\epsilon}$
$\Rightarrow \sqrt{1-\epsilon}<x<\sqrt{1+\epsilon}$ near $x=1$.
Step 2: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow -\delta+1<x<\delta+1.$
Then $-\delta+1=\sqrt{1-\epsilon} \Rightarrow \delta=1-\sqrt{1-\epsilon}$, or $\delta+1=\sqrt{1+\epsilon} \Rightarrow \delta=\sqrt{1+\epsilon}-1$. Choose
$\delta=\min\left\{1-\sqrt{1-\epsilon},\sqrt{1+\epsilon}-1\right\}$, that is, the smaller of the two distances.

42. Step 1: For $x\neq -2$, $\left|x^2-4\right|<\epsilon \Rightarrow -\epsilon<x^2-4<\epsilon \Rightarrow 4-\epsilon<x^2<4+\epsilon \Rightarrow \sqrt{4-\epsilon}<|x|<\sqrt{4+\epsilon} \Rightarrow -\sqrt{4+\epsilon}<x$
$<-\sqrt{4-\epsilon}$ near $x=-2$.
Step 2: $|x-(-2)|<\delta \Rightarrow -\delta<x+2<\delta \Rightarrow -\delta-2<x<\delta-2.$
Then $-\delta-2=-\sqrt{4+\epsilon} \Rightarrow \delta=\sqrt{4+\epsilon}-2$, or $\delta-2=-\sqrt{4-\epsilon} \Rightarrow \delta=2-\sqrt{4-\epsilon}$. Choose
$\delta=\min\left\{\sqrt{4+\epsilon}-2, 2-\sqrt{4-\epsilon}\right\}$.

43. Step 1: $\left|\frac{1}{x}-1\right|<\epsilon \Rightarrow -\epsilon<\frac{1}{x}-1<\epsilon \Rightarrow 1-\epsilon<\frac{1}{x}<1+\epsilon \Rightarrow \frac{1}{1+\epsilon}<x<\frac{1}{1-\epsilon}.$
Step 2: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow 1-\delta<x<1+\delta.$
Then $1-\delta=\frac{1}{1+\epsilon} \Rightarrow \delta=1-\frac{1}{1+\epsilon}=\frac{\epsilon}{1+\epsilon}$, or $1+\delta=\frac{1}{1-\epsilon} \Rightarrow \delta=\frac{1}{1-\epsilon}-1=\frac{\epsilon}{1-\epsilon}$.
Choose $\delta=\frac{\epsilon}{1+\epsilon}$, the smaller of the two distances.

44. Step 1: $\left|\frac{1}{x^2}-\frac{1}{3}\right|<\epsilon \Rightarrow -\epsilon<\frac{1}{x^2}-\frac{1}{3}<\epsilon \Rightarrow \frac{1}{3}-\epsilon<\frac{1}{x^2}<\frac{1}{3}+\epsilon \Rightarrow \frac{1-3\epsilon}{3}<\frac{1}{x^2}<\frac{1+3\epsilon}{3}$
$\Rightarrow \frac{3}{1-3\epsilon}>x^2>\frac{3}{1+3\epsilon} \Rightarrow \sqrt{\frac{3}{1+3\epsilon}}<|x|<\sqrt{\frac{3}{1-3\epsilon}}$, or $\sqrt{\frac{3}{1+3\epsilon}}<x<\sqrt{\frac{3}{1-3\epsilon}}$ for x near $\sqrt{3}$.
Step 2: $\left|x-\sqrt{3}\right|<\delta \Rightarrow -\delta<x-\sqrt{3}<\delta \Rightarrow \sqrt{3}-\delta<x<\sqrt{3}+\delta.$
Then $\sqrt{3}-\delta=\sqrt{\frac{3}{1+3\epsilon}} \Rightarrow \delta=\sqrt{3}-\sqrt{\frac{3}{1+3\epsilon}}$, or $\sqrt{3}+\delta=\sqrt{\frac{3}{1-3\epsilon}} \Rightarrow \delta=\sqrt{\frac{3}{1-3\epsilon}}-\sqrt{3}$.
Choose $\delta=\min\left\{\sqrt{3}-\sqrt{\frac{3}{1+3\epsilon}}, \sqrt{\frac{3}{1-3\epsilon}}-\sqrt{3}\right\}$.

45. Step 1: $\left|\left(\frac{x^2-9}{x+3}\right)-(-6)\right|<\epsilon \Rightarrow -\epsilon<(x-3)+6<\epsilon,\ x\neq -3 \Rightarrow -\epsilon<x+3<\epsilon \Rightarrow -\epsilon-3<x<\epsilon-3.$
Step 2: $|x-(-3)|<\delta \Rightarrow -\delta<x+3<\delta \Rightarrow -\delta-3<x<\delta-3.$
Then $-\delta-3=-\epsilon-3 \Rightarrow \delta=\epsilon$, or $\delta-3=\epsilon-3 \Rightarrow \delta=\epsilon$. Choose $\delta=\epsilon$.

46. Step 1: $\left|\left(\frac{x^2-1}{x-1}\right)-2\right|<\epsilon \Rightarrow -\epsilon<(x+1)-2<\epsilon,\ x\neq 1 \Rightarrow 1-\epsilon<x<1+\epsilon.$
Step 2: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow 1-\delta<x<1+\delta.$
Then $1-\delta=1-\epsilon \Rightarrow \delta=\epsilon$, or $1+\delta=1+\epsilon \Rightarrow \delta=\epsilon$. Choose $\delta=\epsilon$.

47. Step 1: $x<1$: $|(4-2x)-2|<\epsilon \Rightarrow 0<2-2x<\epsilon$ since $x<1$. Thus, $1-\frac{\epsilon}{2}<x<0$;
$x\geq 1$: $|(6x-4)-2|<\epsilon \Rightarrow 0\leq 6x-6<\epsilon$ since $x\geq 1$. Thus, $1\leq x<1+\frac{\epsilon}{6}$.
Step 2: $|x-1|<\delta \Rightarrow -\delta<x-1<\delta \Rightarrow 1-\delta<x<1+\delta.$
Then $1-\delta=1-\frac{\epsilon}{2} \Rightarrow \delta=\frac{\epsilon}{2}$, or $1+\delta=1+\frac{\epsilon}{6} \Rightarrow \delta=\frac{\epsilon}{6}$. Choose $\delta=\frac{\epsilon}{6}$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

48. Step 1: $x < 0$: $|2x - 0| < \epsilon \Rightarrow -\epsilon < 2x < 0 \Rightarrow -\frac{\epsilon}{2} < x < 0$;
$x \geq 0$: $\left|\frac{x}{2} - 0\right| < \epsilon \Rightarrow 0 \leq x < 2\epsilon$.
Step 2: $|x - 0| < \delta \Rightarrow -\delta < x < \delta$.
Then $-\delta = -\frac{\epsilon}{2} \Rightarrow \delta = \frac{\epsilon}{2}$, or $\delta = 2\epsilon \Rightarrow \delta = 2\epsilon$. Choose $\delta = \frac{\epsilon}{2}$.

49. By the figure, $-x \leq x \sin\frac{1}{x} \leq x$ for all $x > 0$ and $-x \geq x \sin\frac{1}{x} \geq x$ for $x < 0$. Since $\lim_{x\to 0}(-x) = \lim_{x\to 0} x = 0$, then by the sandwich theorem, in either case, $\lim_{x\to 0} x \sin\frac{1}{x} = 0$.

50. By the figure, $-x^2 \leq x^2 \sin\frac{1}{x} \leq x^2$ for all x except possibly at $x = 0$. Since $\lim_{x\to 0}(-x^2) = \lim_{x\to 0} x^2 = 0$, then by the sandwich theorem, $\lim_{x\to 0} x^2 \sin\frac{1}{x} = 0$.

51. As x approaches the value 0, the values of $g(x)$ approach k. Thus for every number $\epsilon > 0$, there exists a $\delta > 0$ such that $0 < |x - 0| < \delta \Rightarrow |g(x) - k| < \epsilon$.

52. Write $x = h + c$. Then $0 < |x - c| < \delta \Leftrightarrow -\delta < x - c < \delta$, $x \neq c \Leftrightarrow -\delta < (h + c) - c < \delta$, $h + c \neq c \Leftrightarrow -\delta < h < \delta$, $h \neq 0 \Leftrightarrow 0 < |h - 0| < \delta$.
Thus, $\lim_{x\to c} f(x) = L \Leftrightarrow$ for any $\epsilon > 0$, there exists $\delta > 0$ such that $|f(x) - L| < \epsilon$ whenever $0 < |x - c| < \delta \Leftrightarrow$ $|f(h + c) - L| < \epsilon$ whenever $0 < |h - 0| < \delta \Leftrightarrow \lim_{h\to 0} f(h + c) = L$.

53. Let $f(x) = x^2$. The function values do get closer to -1 as x approaches 0, but $\lim_{x\to 0} f(x) = 0$, not -1. The function $f(x) = x^2$ never gets arbitrarily close to -1 for x near 0.

54. Let $f(x) = \sin x$, $L = \frac{1}{2}$, and $x_0 = 0$. There exists a value of x (namely $x = \frac{\pi}{6}$) for which $\left|\sin x - \frac{1}{2}\right| < \epsilon$ for any given $\epsilon > 0$. However, $\lim_{x\to 0} \sin x = 0$, not $\frac{1}{2}$. The wrong statement does not require x to be arbitrarily close to x_0.
As another example, let $g(x) = \sin\frac{1}{x}$, $L = \frac{1}{2}$, and $x_0 = 0$. We can choose infinitely many values of x near 0 such that $\sin\frac{1}{x} = \frac{1}{2}$ as you can see from the accompanying figure. However, $\lim_{x\to 0} \sin\frac{1}{x}$ fails to exist. The wrong statement does not require all values of x arbitrarily close to $x_0 = 0$ to lie within $\epsilon > 0$ of $L = \frac{1}{2}$. Again you can see from the figure that there are also infinitely many values of x near 0 such that $\sin\frac{1}{x} = 0$. If we choose $\epsilon < \frac{1}{4}$ we cannot satisfy the inequality $\left|\sin\frac{1}{x} - \frac{1}{2}\right| < \epsilon$ for all values of x sufficiently near $x_0 = 0$.



55. $|A - 9| \leq 0.01 \Rightarrow -0.01 \leq \pi\left(\frac{x}{2}\right)^2 - 9 \leq 0.01 \Rightarrow 8.99 \leq \frac{\pi x^2}{4} \leq 9.01 \Rightarrow \frac{4}{\pi}(8.99) \leq x^2 \leq \frac{4}{\pi}(9.01)$
$\Rightarrow 2\sqrt{\frac{8.99}{\pi}} \leq x \leq 2\sqrt{\frac{9.01}{\pi}}$ or $3.384 \leq x \leq 3.387$. To be safe, the left endpoint was rounded up and the right endpoint was rounded down.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

56. $V = RI \Rightarrow \frac{V}{R} = I \Rightarrow \left|\frac{V}{R} - 5\right| \le 0.1 \Rightarrow -0.1 \le \frac{120}{R} - 5 \le 0.1 \Rightarrow 4.9 \le \frac{120}{R} \le 5.1 \Rightarrow \frac{10}{49} \ge \frac{R}{120} \ge \frac{10}{51}$
$\Rightarrow \frac{(120)(10)}{51} \le R \le \frac{(120)(10)}{49} \Rightarrow 23.53 \le R \le 24.48.$

To be safe, the left endpoint was rounded up and the right endpoint was rounded down.

57. (a) $-\delta < x-1 < 0 \Rightarrow 1-\delta < x < 1 \Rightarrow f(x) = x$. Then $|f(x)-2| = |x-2| = 2-x > 2-1 = 1$. That is, $|f(x)-2| \ge 1 \ge \frac{1}{2}$ no matter how small δ is taken when $1-\delta < x < 1 \Rightarrow \lim_{x\to 1} f(x) \ne 2$.

(b) $0 < x-1 < \delta \Rightarrow 1 < x < 1+\delta \Rightarrow f(x) = x+1$. Then $|f(x)-1| = |(x+1)-1| = |x| = x > 1$. That is, $|f(x)-1| \ge 1$ no matter how small δ is taken when $1 < x < 1+\delta \Rightarrow \lim_{x\to 1} f(x) \ne 1$.

(c) $-\delta < x-1 < 0 \Rightarrow 1-\delta < x < 1 \Rightarrow f(x) = x$. Then $|f(x)-1.5| = |x-1.5| = 1.5-x > 1.5-1 = 0.5$.
Also, $0 < x-1 < \delta \Rightarrow 1 < x < 1+\delta \Rightarrow f(x) = x+1$. Then $|f(x)-1.5| = |(x+1)-1.5| = |x-0.5|$
$= x-0.5 > 1-0.5 = 0.5$. Thus, no matter how small δ is taken, there exists a value of x such that $-\delta < x-1 < \delta$ but $|f(x)-1.5| \ge \frac{1}{2} \Rightarrow \lim_{x\to 1} f(x) \ne 1.5$.

58. (a) For $2 < x < 2+\delta \Rightarrow h(x) = 2 \Rightarrow |h(x)-4| = 2$. Thus for $\epsilon < 2$, $|h(x)-4| \ge \epsilon$ whenever $2 < x < 2+\delta$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 2} h(x) \ne 4$.

(b) For $2 < x < 2+\delta \Rightarrow h(x) = 2 \Rightarrow |h(x)-3| = 1$. Thus for $\epsilon < 1$, $|h(x)-3| \ge \epsilon$ whenever $2 < x < 2+\delta$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 2} h(x) \ne 3$.

(c) For $2-\delta < x < 2 \Rightarrow h(x) = x^2$ so $|h(x)-2| = |x^2-2|$. No matter how small $\delta > 0$ is chosen, x^2 is close to 4 when x is near 2 and to the left on the real line $\Rightarrow |x^2-2|$ will be close to 2. Thus if $\epsilon < 1$, $|h(x)-2| \ge \epsilon$ whenever $2-\delta < x < 2$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 2} h(x) \ne 2$.

59. (a) For $3-\delta < x < 3 \Rightarrow f(x) > 4.8 \Rightarrow |f(x)-4| \ge 0.8$. Thus for $\epsilon < 0.8$, $|f(x)-4| \ge \epsilon$ whenever $3-\delta < x < 3$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 3} f(x) \ne 4$.

(b) For $3 < x < 3+\delta \Rightarrow f(x) < 3 \Rightarrow |f(x)-4.8| \ge 1.8$. Thus for $\epsilon < 1.8$, $|f(x)-4.8| \ge \epsilon$ whenever $3 < x < 3+\delta$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 3} f(x) \ne 4.8$.

(c) For $3-\delta < x < 3 \Rightarrow f(x) > 4.8 \Rightarrow |f(x)-3| \ge 1.8$. Again, for $\epsilon < 1.8$, $|f(x)-3| \ge \epsilon$ whenever $3-\delta < x < 3$ no matter how small we choose $\delta > 0 \Rightarrow \lim_{x\to 3} f(x) \ne 3$.

60. (a) No matter how small we choose $\delta > 0$, for x near -1 satisfying $-1-\delta < x < -1+\delta$, the values of $g(x)$ are near $1 \Rightarrow |g(x)-2|$ is near 1. Then, for $\epsilon = \frac{1}{2}$ we have $|g(x)-2| \ge \frac{1}{2}$ for some x satisfying $-1-\delta < x < -1+\delta$, or $0 < |x+1| < \delta \Rightarrow \lim_{x\to -1} g(x) \ne 2$.

(b) Yes, $\lim_{x\to -1} g(x) = 1$ because from the graph we can find a $\delta > 0$ such that $|g(x)-1| < \epsilon$ if $0 < |x-(-1)| < \delta$.

61–66. Example CAS commands (values of del may vary for a specified eps):

Maple:

```
f := x -> (x^4-81)/(x-3); x0 := 3;
plot( f(x), x=x0-1..x0+1, color=black,                    # (a)
      title="Section 2.3, #61(a)" );
L := limit( f(x), x=x0 );                                 # (b)
epsilon := 0.2;                                           # (c)
plot( [f(x), L-epsilon, L+epsilon], x=x0-0.01..x0+0.01,
      color=black, linestyle=[1,3,3], title="Section 2.3, #61(c)" );
```

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

```
q := fsolve( abs( f(x)-L ) = epsilon, x=x0-1..x0+1 );          # (d)
delta := abs(x0-q);
plot( [f(x), L-epsilon, L+epsilon], x=x0-delta..x0+delta, color=black, title="Section 2.3, #61(d)" );
for eps in [0.1, 0.005, 0.001 ] do                              # (e)
q := fsolve( abs( f(x)-L ) = eps, x=x0-1..x0+1 );
delta := abs(x0-q);
head := sprintf("Section 2.3, #61(e)\n epsilon = %5f, delta = %5f \n", eps, delta );
print(plot( [f(x), L-eps, L+eps], x=x0-delta..x0+delta,
color=black, linestyle=[1,3,3], title=head ));
end do:
```

Mathematica (assigned function and values for x0, eps and del may vary):

```
Clear[f, x]
y1: = L − eps; y2: = L + eps; x0 = 1;
f[x_]: = (3x^2 − (7x + 1)Sqrt[x] + 5)/(x − 1)
Plot[f[x], {x, x0 − 0.2, x0 + 0.2}]
L: = Limit[f[x], x → x0]
eps = 0.1; del = 0.2;
Plot[{f[x], y1, y2}, {x, x0 − del, x0 + del}, PlotRange → {L − 2eps, L + 2eps}]
```

2.4 ONE-SIDED LIMITS

1. (a) True (b) True (c) False (d) True
   (e) True (f) True (g) False (h) False
   (i) False (j) False (k) True (l) False

2. (a) True (b) False (c) False (d) True
   (e) True (f) True (g) True (h) True
   (i) True (j) False (k) True

3. (a) $\lim\limits_{x\to 2^+} f(x) = \frac{2}{2} + 1 = 2,\ \lim\limits_{x\to 2^-} f(x) = 3 - 2 = 1$
   (b) No, $\lim\limits_{x\to 2} f(x)$ does not exist because $\lim\limits_{x\to 2^+} f(x) \neq \lim\limits_{x\to 2^-} f(x)$
   (c) $\lim\limits_{x\to 4^-} f(x) = \frac{4}{2} + 1 = 3,\ \lim\limits_{x\to 4^+} f(x) = \frac{4}{2} + 1 = 3$
   (d) Yes, $\lim\limits_{x\to 4} f(x) = 3$ because $3 = \lim\limits_{x\to 4^-} f(x) = \lim\limits_{x\to 4^+} f(x)$

4. (a) $\lim\limits_{x\to 2^+} f(x) = \frac{2}{2} = 1,\ \lim\limits_{x\to 2^-} f(x) = 3 - 2 = 1,\ f(2) = 2$
   (b) Yes, $\lim\limits_{x\to 2} f(x) = 1$ because $1 = \lim\limits_{x\to 2^+} f(x) = \lim\limits_{x\to 2^-} f(x)$
   (c) $\lim\limits_{x\to -1^-} f(x) = 3 - (-1) = 4,\ \lim\limits_{x\to -1^+} f(x) = 3 - (-1) = 4$
   (d) Yes, $\lim\limits_{x\to -1} f(x) = 4$ because $4 = \lim\limits_{x\to -1^-} f(x) = \lim\limits_{x\to -1^+} f(x)$

5. (a) No, $\lim\limits_{x\to 0^+} f(x)$ does not exist since $\sin\left(\frac{1}{x}\right)$ does not approach any single value as x approaches 0
   (b) $\lim\limits_{x\to 0^-} f(x) = \lim\limits_{x\to 0^-} 0 = 0$
   (c) $\lim\limits_{x\to 0} f(x)$ does not exist because $\lim\limits_{x\to 0^+} f(x)$ does not exist

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

6. (a) Yes, $\lim_{x\to 0^+} g(x) = 0$ by the sandwich theorem since $-\sqrt{x} \le g(x) \le \sqrt{x}$ when $x > 0$

   (b) No, $\lim_{x\to 0^-} g(x)$ does not exist since $\sqrt{x}$ is not defined for $x < 0$

   (c) Yes, $\lim_{x\to 0} g(x) = \lim_{x\to 0^+} g(x) = 0$ since $x = 0$ is a boundary point of the domain

7. (a)



   (b) $\lim_{x\to 1^-} f(x) = 1 = \lim_{x\to 1^+} f(x)$

   (c) Yes, $\lim_{x\to 1} f(x) = 1$ since the right-hand and left-hand limits exist and equal 1

8. (a)



   (b) $\lim_{x\to 1^+} f(x) = 0 = \lim_{x\to 1^-} f(x)$

   (c) Yes, $\lim_{x\to 1} f(x) = 0$ since the right-hand and left-hand limits exist and equal 0

9. (a) domain: $0 \le x \le 2$
   range: $0 < y \le 1$ and $y = 2$

   (b) $\lim_{x\to c} f(x)$ exists for c belonging to $(0, 1) \cup (1, 2)$

   (c) $x = 2$

   (d) $x = 0$



10. (a) domain: $-\infty < x < \infty$
    range: $-1 \le y \le 1$

    (b) $\lim_{x\to c} f(x)$ exists for c belonging to
    $(-\infty, -1) \cup (-1, 1) \cup (1, \infty)$

    (c) none

    (d) none



11. $\lim_{x\to -0.5^-} \sqrt{\frac{x+2}{x-1}} = \sqrt{\frac{-0.5+2}{-0.5+1}} = \sqrt{\frac{3/2}{1/2}} = \sqrt{3}$

12. $\lim_{x\to 1^+} \sqrt{\frac{x-1}{x+2}} = \sqrt{\frac{1-1}{1+2}} = \sqrt{0} = 0$

13. $\lim_{x\to -2^+} \left(\frac{x}{x+1}\right)\left(\frac{2x+5}{x^2+x}\right) = \left(\frac{-2}{-2+1}\right)\left(\frac{2(-2)+5}{(-2)^2+(-2)}\right) = (2)\left(\frac{1}{2}\right) = 1$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

14. $\lim\limits_{x\to 1^-}\left(\frac{1}{x+1}\right)\left(\frac{x+6}{x}\right)\left(\frac{3-x}{7}\right)=\left(\frac{1}{1+1}\right)\left(\frac{1+6}{1}\right)\left(\frac{3-1}{7}\right)=\left(\frac{1}{2}\right)\left(\frac{7}{1}\right)\left(\frac{2}{7}\right)=1$

15. $\lim\limits_{h\to 0^+}\frac{\sqrt{h^2+4h+5}-\sqrt{5}}{h}=\lim\limits_{h\to 0^+}\left(\frac{\sqrt{h^2+4h+5}-\sqrt{5}}{h}\right)\left(\frac{\sqrt{h^2+4h+5}+\sqrt{5}}{\sqrt{h^2+4h+5}+\sqrt{5}}\right)=\lim\limits_{h\to 0^+}\frac{(h^2+4h+5)-5}{h\left(\sqrt{h^2+4h+5}+\sqrt{5}\right)}$

$=\lim\limits_{h\to 0^+}\frac{h(h+4)}{h\left(\sqrt{h^2+4h+5}+\sqrt{5}\right)}=\frac{0+4}{\sqrt{5}+\sqrt{5}}=\frac{2}{\sqrt{5}}$

16. $\lim\limits_{h\to 0^-}\frac{\sqrt{6}-\sqrt{5h^2+11h+6}}{h}=\lim\limits_{h\to 0^-}\left(\frac{\sqrt{6}-\sqrt{5h^2+11h+6}}{h}\right)\left(\frac{\sqrt{6}+\sqrt{5h^2+11h+6}}{\sqrt{6}+\sqrt{5h^2+11h+6}}\right)$

$=\lim\limits_{h\to 0^-}\frac{6-(5h^2+11h+6)}{h\left(\sqrt{6}+\sqrt{5h^2+11h+6}\right)}=\lim\limits_{h\to 0^-}\frac{-h(5h+11)}{h\left(\sqrt{6}+\sqrt{5h^2+11h+6}\right)}=\frac{-(0+11)}{\sqrt{6}+\sqrt{6}}=-\frac{11}{2\sqrt{6}}$

17. (a) $\lim\limits_{x\to -2^+}(x+3)\frac{|x+2|}{x+2}=\lim\limits_{x\to -2^+}(x+3)\frac{(x+2)}{(x+2)}$ $(|x+2|=(x+2)$ for $x>-2)$

$=\lim\limits_{x\to -2^+}(x+3)=((-2)+3)=1$

(b) $\lim\limits_{x\to -2^-}(x+3)\frac{|x+2|}{x+2}=\lim\limits_{x\to -2^-}(x+3)\left[\frac{-(x+2)}{(x+2)}\right]$ $(|x+2|=-(x+2)$ for $x<-2)$

$=\lim\limits_{x\to -2^-}(x+3)(-1)=-(-2+3)=-1$

18. (a) $\lim\limits_{x\to 1^+}\frac{\sqrt{2x}(x-1)}{|x-1|}=\lim\limits_{x\to -1^+}\frac{\sqrt{2x}(x-1)}{(x-1)}$ $(|x-1|=x-1$ for $x>1)$

$=\lim\limits_{x\to -1^+}\sqrt{2x}=\sqrt{2}$

(b) $\lim\limits_{x\to 1^-}\frac{\sqrt{2x}(x-1)}{|x-1|}=\lim\limits_{x\to 1^-}\frac{\sqrt{2x}(x-1)}{-(x-1)}$ $(|x-1|=-(x-1)$ for $x<1)$

$=\lim\limits_{x\to 1^-}-\sqrt{2x}=-\sqrt{2}$

19. (a) If $0<x<\frac{\pi}{2}$, then $\sin x>0$, so that $\lim\limits_{x\to 0^+}\frac{|\sin x|}{\sin x}=\lim\limits_{x\to 0^+}\frac{\sin x}{\sin x}=\lim\limits_{x\to 0^+}1=1$

(b) If $\frac{-\pi}{2}<x<0$, then $\sin x<0$, so that $\lim\limits_{x\to 0^-}\frac{|\sin x|}{\sin x}=\lim\limits_{x\to 0^-}\frac{-\sin x}{\sin x}=\lim\limits_{x\to 0^-}-1=-1$

20. (a) If $0<x<\frac{\pi}{2}$, then $\cos x<1$, so that $\lim\limits_{x\to 0^+}\frac{1-\cos x}{|\cos x-1|}=\lim\limits_{x\to 0^+}\frac{1-\cos x}{-(\cos x-1)}=\lim\limits_{x\to 0^+}\frac{1-\cos x}{1-\cos x}=\lim\limits_{x\to 0^+}1=1$

(b) If $\frac{-\pi}{2}<x<0$, then $\cos x<1$, so that $\lim\limits_{x\to 0^-}\frac{\cos x-1}{|\cos x-1|}=\lim\limits_{x\to 0^-}\frac{\cos x-1}{-(\cos x-1)}=\lim\limits_{x\to 0^-}-1=-1$

21. (a) $\lim\limits_{\theta\to 3^+}\frac{\lfloor\theta\rfloor}{\theta}=\frac{3}{3}=1$ (b) $\lim\limits_{\theta\to 3^-}\frac{\lfloor\theta\rfloor}{\theta}=\frac{2}{3}$

22. (a) $\lim\limits_{t\to 4^+}(t-\lfloor t\rfloor)=4-4=0$ (b) $\lim\limits_{t\to 4^-}(t-\lfloor t\rfloor)=4-3=1$

23. $\lim\limits_{\theta\to 0}\frac{\sin\sqrt{2}\theta}{\sqrt{2}\theta}=\lim\limits_{x\to 0}\frac{\sin x}{x}=1$ (where $x=\sqrt{2}\theta$)

24. $\lim\limits_{t\to 0}\frac{\sin kt}{t}=\lim\limits_{t\to 0}\frac{k\sin kt}{kt}=\lim\limits_{\theta\to 0}\frac{k\sin\theta}{\theta}=k\lim\limits_{\theta\to 0}\frac{\sin\theta}{\theta}=k\cdot 1=k$ (where $\theta=kt$)

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

25. $\lim_{y\to 0}\frac{\sin 3y}{4y}=\frac{1}{4}\lim_{y\to 0}\frac{3\sin 3y}{3y}=\frac{3}{4}\lim_{y\to 0}\frac{\sin 3y}{3y}=\frac{3}{4}\lim_{\theta\to 0}\frac{\sin\theta}{\theta}=\frac{3}{4}$ (where $\theta=3y$)

26. $\lim_{h\to 0^-}\frac{h}{\sin 3h}=\lim_{h\to 0^-}\left(\frac{1}{3}\cdot\frac{3h}{\sin 3h}\right)=\frac{1}{3}\lim_{h\to 0^-}\frac{1}{\left(\frac{\sin 3h}{3h}\right)}=\frac{1}{3}\left(\frac{1}{\lim_{\theta\to 0^-}\frac{\sin\theta}{\theta}}\right)=\frac{1}{3}\cdot 1=\frac{1}{3}$ (where $\theta=3h$)

27. $\lim_{x\to 0}\frac{\tan 2x}{x}=\lim_{x\to 0}\frac{\left(\frac{\sin 2x}{\cos 2x}\right)}{x}=\lim_{x\to 0}\frac{\sin 2x}{x\cos 2x}=\left(\lim_{x\to 0}\frac{1}{\cos 2x}\right)\left(\lim_{x\to 0}\frac{2\sin 2x}{2x}\right)=1\cdot 2=2$

28. $\lim_{t\to 0}\frac{2t}{\tan t}=2\lim_{t\to 0}\frac{t}{\left(\frac{\sin t}{\cos t}\right)}=2\lim_{t\to 0}\frac{t\cos t}{\sin t}=2\left(\lim_{t\to 0}\cos t\right)\left(\frac{1}{\lim_{t\to 0}\frac{\sin t}{t}}\right)=2\cdot 1\cdot 1=2$

29. $\lim_{x\to 0}\frac{x\csc 2x}{\cos 5x}=\lim_{x\to 0}\left(\frac{x}{\sin 2x}\cdot\frac{1}{\cos 5x}\right)=\left(\frac{1}{2}\lim_{x\to 0}\frac{2x}{\sin 2x}\right)\left(\lim_{x\to 0}\frac{1}{\cos 5x}\right)=\left(\frac{1}{2}\cdot 1\right)(1)=\frac{1}{2}$

30. $\lim_{x\to 0}6x^2(\cot x)(\csc 2x)=\lim_{x\to 0}\frac{6x^2\cos x}{\sin x\sin 2x}=\lim_{x\to 0}\left(3\cos x\cdot\frac{x}{\sin x}\cdot\frac{2x}{\sin 2x}\right)=3\cdot 1\cdot 1=3$

31. $\lim_{x\to 0}\frac{x+x\cos x}{\sin x\cos x}=\lim_{x\to 0}\left(\frac{x}{\sin x\cos x}+\frac{x\cos x}{\sin x\cos x}\right)=\lim_{x\to 0}\left(\frac{x}{\sin x}\cdot\frac{1}{\cos x}\right)+\lim_{x\to 0}\frac{x}{\sin x}$

$=\lim_{x\to 0}\left(\frac{1}{\frac{\sin x}{x}}\right)\cdot\lim_{x\to 0}\left(\frac{1}{\cos x}\right)+\lim_{x\to 0}\left(\frac{1}{\frac{\sin x}{x}}\right)=(1)(1)+1=2$

32. $\lim_{x\to 0}\frac{x^2-x+\sin x}{2x}=\lim_{x\to 0}\left(\frac{x}{2}-\frac{1}{2}+\frac{1}{2}\left(\frac{\sin x}{x}\right)\right)=0-\frac{1}{2}+\frac{1}{2}(1)=0$

33. $\lim_{\theta\to 0}\frac{1-\cos\theta}{\sin 2\theta}=\lim_{\theta\to 0}\frac{(1-\cos\theta)(1+\cos\theta)}{(2\sin\theta\cos\theta)(1+\cos\theta)}=\lim_{\theta\to 0}\frac{1-\cos^2\theta}{(2\sin\theta\cos\theta)(1+\cos\theta)}=\lim_{\theta\to 0}\frac{\sin^2\theta}{(2\sin\theta\cos\theta)(1+\cos\theta)}$

$=\lim_{\theta\to 0}\frac{\sin\theta}{(2\cos\theta)(1+\cos\theta)}=\frac{0}{(2)(2)}=0$

34. $\lim_{x\to 0}\frac{x-x\cos x}{\sin^2 3x}=\lim_{x\to 0}\frac{x(1-\cos x)}{\sin^2 3x}=\lim_{x\to 0}\frac{\frac{x(1-\cos x)}{9x^2}}{\frac{\sin^2 3x}{9x^2}}=\lim_{x\to 0}\frac{\frac{1-\cos x}{9x}}{\left(\frac{\sin 3x}{3x}\right)^2}=\frac{\frac{1}{9}\lim_{x\to 0}\left(\frac{1-\cos x}{x}\right)}{\lim_{x\to 0}\left(\frac{\sin 3x}{3x}\right)^2}=\frac{\frac{1}{9}(0)}{1^2}=0$

35. $\lim_{t\to 0}\frac{\sin(1-\cos t)}{1-\cos t}=\lim_{\theta\to 0}\frac{\sin\theta}{\theta}=1$ since $\theta=1-\cos t\to 0$ as $t\to 0$

36. $\lim_{h\to 0}\frac{\sin(\sin h)}{\sin h}=\lim_{\theta\to 0}\frac{\sin\theta}{\theta}=1$ since $\theta=\sin h\to 0$ as $h\to 0$

37. $\lim_{\theta\to 0}\frac{\sin\theta}{\sin 2\theta}=\lim_{\theta\to 0}\left(\frac{\sin\theta}{\sin 2\theta}\cdot\frac{2\theta}{2\theta}\right)=\frac{1}{2}\lim_{\theta\to 0}\left(\frac{\sin\theta}{\theta}\cdot\frac{2\theta}{\sin 2\theta}\right)=\frac{1}{2}\cdot 1\cdot 1=\frac{1}{2}$

38. $\lim_{x\to 0}\frac{\sin 5x}{\sin 4x}=\lim_{x\to 0}\left(\frac{\sin 5x}{\sin 4x}\cdot\frac{4x}{5x}\cdot\frac{5}{4}\right)=\frac{5}{4}\lim_{x\to 0}\left(\frac{\sin 5x}{5x}\cdot\frac{4x}{\sin 4x}\right)=\frac{5}{4}\cdot 1\cdot 1=\frac{5}{4}$

39. $\lim_{\theta\to 0}\theta\cos\theta=0\cdot 1=0$

40. $\lim_{\theta\to 0}\sin\theta\cot 2\theta=\lim_{\theta\to 0}\sin\theta\frac{\cos 2\theta}{\sin 2\theta}=\lim_{\theta\to 0}\sin\theta\frac{\cos 2\theta}{2\sin\theta\cos\theta}=\lim_{\theta\to 0}\frac{\cos 2\theta}{2\cos\theta}=\frac{1}{2}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

41. $\lim_{x\to 0}\frac{\tan 3x}{\sin 8x}=\lim_{x\to 0}\left(\frac{\sin 3x}{\cos 3x}\cdot\frac{1}{\sin 8x}\right)=\lim_{x\to 0}\left(\frac{\sin 3x}{\cos 3x}\cdot\frac{1}{\sin 8x}\cdot\frac{8x}{3x}\cdot\frac{3}{8}\right)$
$=\frac{3}{8}\lim_{x\to 0}\left(\frac{1}{\cos 3x}\right)\left(\frac{\sin 3x}{3x}\right)\left(\frac{8x}{\sin 8x}\right)=\frac{3}{8}\cdot 1\cdot 1\cdot 1=\frac{3}{8}$

42. $\lim_{y\to 0}\frac{\sin 3y\cot 5y}{y\cot 4y}=\lim_{y\to 0}\frac{\sin 3y\sin 4y\cos 5y}{y\cos 4y\sin 5y}=\lim_{y\to 0}\left(\frac{\sin 3y}{y}\right)\left(\frac{\sin 4y}{\cos 4y}\right)\left(\frac{\cos 5y}{\sin 5y}\right)\left(\frac{3\cdot 4\cdot 5y}{3\cdot 4\cdot 5y}\right)$
$=\lim_{y\to 0}\left(\frac{\sin 3y}{3y}\right)\left(\frac{\sin 4y}{4y}\right)\left(\frac{5y}{\sin 5y}\right)\left(\frac{\cos 5y}{\cos 4y}\right)\left(\frac{3\cdot 4}{5}\right)=1\cdot 1\cdot 1\cdot 1\cdot\frac{12}{5}=\frac{12}{5}$

43. $\lim_{\theta\to 0}\frac{\tan\theta}{\theta^2\cot 3\theta}=\lim_{\theta\to 0}\frac{\frac{\sin\theta}{\cos\theta}}{\theta^2\frac{\cos 3\theta}{\sin 3\theta}}=\lim_{\theta\to 0}\frac{\sin\theta\sin 3\theta}{\theta^2\cos\theta\cos 3\theta}=\lim_{\theta\to 0}\left(\frac{\sin\theta}{\theta}\right)\left(\frac{\sin 3\theta}{3\theta}\right)\left(\frac{3}{\cos\theta\cos 3\theta}\right)=(1)(1)\left(\frac{3}{1\cdot 1}\right)=3$

44. $\lim_{\theta\to 0}\frac{\theta\cot 4\theta}{\sin^2\theta\cot^2 2\theta}=\lim_{\theta\to 0}\frac{\theta\frac{\cos 4\theta}{\sin 4\theta}}{\sin^2\theta\frac{\cos^2 2\theta}{\sin^2 2\theta}}=\lim_{\theta\to 0}\frac{\theta\cos 4\theta\sin^2 2\theta}{\sin^2\theta\cos^2 2\theta\sin 4\theta}=\lim_{\theta\to 0}\frac{\theta\cos 4\theta(2\sin\theta\cos\theta)^2}{\sin^2\theta\cos^2 2\theta\sin 4\theta}=\lim_{\theta\to 0}\frac{\theta\cos 4\theta(4\sin^2\theta\cos^2\theta)}{\sin^2\theta\cos^2 2\theta\sin 4\theta}$
$=\lim_{\theta\to 0}\frac{4\theta\cos 4\theta\cos^2\theta}{\cos^2 2\theta\sin 4\theta}=\lim_{\theta\to 0}\left(\frac{4\theta}{\sin 4\theta}\right)\left(\frac{\cos 4\theta\cos^2\theta}{\cos^2 2\theta}\right)=\lim_{\theta\to 0}\left(\frac{1}{\frac{\sin 4\theta}{4\theta}}\right)\left(\frac{\cos 4\theta\cos^2\theta}{\cos^2 2\theta}\right)=\left(\frac{1}{1}\right)\left(\frac{1\cdot 1^2}{1^2}\right)=1$

45. $\lim_{x\to 0}\frac{1-\cos 3x}{2x}=\lim_{x\to 0}\frac{1-\cos 3x}{2x}\cdot\frac{1+\cos 3x}{1+\cos 3x}=\lim_{x\to 0}\frac{1-\cos^2 3x}{2x(1+\cos 3x)}=\lim_{x\to 0}\frac{\sin^2 3x}{2x(1+\cos 3x)}=\lim_{x\to 0}\frac{3}{2}\cdot\frac{\sin 3x}{3x}\cdot\frac{\sin 3x}{1+\cos 3x}$
$=\lim_{\theta\to 0}\frac{3}{2}\cdot\frac{\sin\theta}{\theta}\cdot\frac{\sin\theta}{1+\cos\theta}=\frac{3}{2}(1)\left(\frac{0}{1+1}\right)=0\quad(\text{where }\theta=3x)$

46. $\lim_{x\to 0}\frac{\cos^2 x-\cos x}{x^2}=\lim_{x\to 0}\frac{\cos x(\cos x-1)}{x^2}=\lim_{x\to 0}\frac{\cos x(\cos x-1)}{x^2}\cdot\frac{\cos x+1}{\cos x+1}=\lim_{x\to 0}\frac{\cos x(\cos^2 x-1)}{x^2(\cos x+1)}=\lim_{x\to 0}\frac{\cos x\cdot(-\sin^2 x)}{x^2(\cos x+1)}$
$=\lim_{x\to 0}\left\{-\frac{\sin x}{x}\cdot\frac{\sin x}{x}\cdot\frac{\cos x}{\cos x+1}\right\}=-(1)(1)\cdot\frac{1}{1+1}=\frac{-1}{2}$

47. Yes. If $\lim_{x\to a^+}f(x)=L=\lim_{x\to a^-}f(x)$, then $\lim_{x\to a}f(x)=L$. If $\lim_{x\to a^+}f(x)\neq\lim_{x\to a^-}f(x)$, then $\lim_{x\to a}f(x)$ does not exist.

48. Since $\lim_{x\to c}f(x)=L$ if and only if $\lim_{x\to c^+}f(x)=L$ and $\lim_{x\to c^-}f(x)=L$, then $\lim_{x\to c}f(x)$ can be found by calculating $\lim_{x\to c^+}f(x)$.

49. If f is an odd function of x, then $f(-x)=-f(x)$. Given $\lim_{x\to 0^+}f(x)=3$, then $\lim_{x\to 0^-}f(x)=-3$.

50. If f is an even function of x, then $f(-x)=f(x)$. Given $\lim_{x\to 2^-}f(x)=7$ then $\lim_{x\to -2^+}f(x)=7$. However, nothing can be said about $\lim_{x\to -2^-}f(x)$ because we don't know $\lim_{x\to 2^+}f(x)$.

51. $I=(5,5+\delta)\Rightarrow 5<x<5+\delta$. Also, $\sqrt{x-5}<\epsilon\Rightarrow x-5<\epsilon^2\Rightarrow x<5+\epsilon^2$. Choose $\delta=\epsilon^2\Rightarrow\lim_{x\to 5^+}\sqrt{x-5}=0$.

52. $I=(4-\delta,4)\Rightarrow 4-\delta<x<4$. Also, $\sqrt{4-x}<\epsilon\Rightarrow 4-x<\epsilon^2\Rightarrow x>4-\epsilon^2$. Choose $\delta=\epsilon^2\Rightarrow\lim_{x\to 4^-}\sqrt{4-x}=0$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

53. As $x \to 0^-$ the number x is always negative. Thus, $\left|\frac{x}{|x|} - (-1)\right| < \epsilon \Rightarrow \left|\frac{x}{-x} + 1\right| < \epsilon \Rightarrow 0 < \epsilon$ which is always true independent of the value of x. Hence we can choose any $\delta > 0$ with $-\delta < x < 0 \Rightarrow \lim\limits_{x \to 0^-} \frac{x}{|x|} = -1$.

54. Since $x \to 2^+$ we have $x > 2$ and $|x-2| = x-2$. Then, $\left|\frac{x-2}{|x-2|} - 1\right| = \left|\frac{x-2}{x-2} - 1\right| < \epsilon \Rightarrow 0 < \epsilon$ which is always true so long as $x > 2$. Hence we can choose any $\delta > 0$, and thus $2 < x < 2 + \delta \Rightarrow \left|\frac{x-2}{|x-2|} - 1\right| < \epsilon$. Thus, $\lim\limits_{x \to -2^+} \frac{x-2}{|x-2|} = 1$.

55. (a) $\lim\limits_{x \to 400^+} \lfloor x \rfloor = 400$. Just observe that if $400 < x < 401$, then $\lfloor x \rfloor = 400$. Thus if we choose $\delta = 1$, we have for any number $\epsilon > 0$ that $400 < x < 400 + \delta \Rightarrow |\lfloor x \rfloor - 400| = |400 - 400| = 0 < \epsilon$.

(b) $\lim\limits_{x \to 400^-} \lfloor x \rfloor = 399$. Just observe that if $399 < x < 400$ then $\lfloor x \rfloor = 399$. Thus if we choose $\delta = 1$, we have for any number $\epsilon > 0$ that $400 - \delta < x < 400 \Rightarrow |\lfloor x \rfloor - 399| = |399 - 399| = 0 < \epsilon$.

(c) Since $\lim\limits_{x \to 400^+} \lfloor x \rfloor \neq \lim\limits_{x \to 400^-} \lfloor x \rfloor$ we conclude that $\lim\limits_{x \to 400} \lfloor x \rfloor$ does not exist.

56. (a) $\lim\limits_{x \to 0^+} f(x) = \lim\limits_{x \to 0^+} \sqrt{x} = \sqrt{0} = 0$; $|\sqrt{x} - 0| < \epsilon \Rightarrow -\epsilon < \sqrt{x} < \epsilon \Rightarrow 0 < x < \epsilon^2$ for x positive. Choose $\delta = \epsilon^2$ $\Rightarrow \lim\limits_{x \to 0^+} f(x) = 0$.

(b) $\lim\limits_{x \to 0^-} f(x) = \lim\limits_{x \to 0^-} x^2 \sin\left(\frac{1}{x}\right) = 0$ by the sandwich theorem since $-x^2 \le x^2 \sin\left(\frac{1}{x}\right) \le x^2$ for all $x \neq 0$. Since $|x^2 - 0| = |-x^2 - 0| = x^2 < \epsilon$ whenever $|x| < \sqrt{\epsilon}$, we choose $\delta = \sqrt{\epsilon}$ and obtain $\left|x^2 \sin\left(\frac{1}{x}\right) - 0\right| < \epsilon$ if $-\delta < x < 0$.

(c) The function f has limit 0 at $x_0 = 0$ since both the right-hand and left-hand limits exist and equal 0.

2.5 CONTINUITY

1. No, discontinuous at $x = 2$, not defined at $x = 2$

2. No, discontinuous at $x = 3, 1 = \lim\limits_{x \to 3^-} g(x) \neq g(3) = 1.5$

3. Continuous on $[-1, 3]$

4. No, discontinuous at $x = 1, 1.5 = \lim\limits_{x \to 1^-} k(x) \neq \lim\limits_{x \to 1^+} k(x) = 0$

5. (a) Yes
   (b) Yes, $\lim\limits_{x \to -1^+} f(x) = 0$
   (c) Yes
   (d) Yes

6. (a) Yes, $f(1) = 1$
   (b) Yes, $\lim\limits_{x \to 1} f(x) = 2$
   (c) No
   (d) No

7. (a) No
   (b) No

8. $[-1, 0) \cup (0, 1) \cup (1, 2) \cup (2, 3)$

9. $f(2) = 0$, since $\lim\limits_{x \to 2^-} f(x) = -2(2) + 4 = 0 = \lim\limits_{x \to 2^+} f(x)$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

10. $f(1)$ should be changed to $2 = \lim_{x\to 1} f(x)$

11. Nonremovable discontinuity at $x = 1$ because $\lim_{x\to 1} f(x)$ fails to exist ($\lim_{x\to 1^-} f(x) = 1$ and $\lim_{x\to 1^+} f(x) = 0$). Removable discontinuity at $x = 0$ by assigning the number $\lim_{x\to 0} f(x) = 0$ to be the value of $f(0)$ rather than $f(0) = 1$.

12. Nonremovable discontinuity at $x = 1$ because $\lim_{x\to 1} f(x)$ fails to exist ($\lim_{x\to 1^-} f(x) = 2$ and $\lim_{x\to 1^+} f(x) = 1$). Removable discontinuity at $x = 2$ by assigning the number $\lim_{x\to 2} f(x) = 1$ to be the value of $f(2)$ rather than $f(2) = 2$.

13. Discontinuous only when $x - 2 = 0 \Rightarrow x = 2$

14. Discontinuous only when $(x+2)^2 = 0 \Rightarrow x = -2$

15. Discontinuous only when $x^2 - 4x + 3 = 0 \Rightarrow (x-3)(x-1) = 0 \Rightarrow x = 3$ or $x = 1$

16. Discontinuous only when $x^2 - 3x - 10 = 0 \Rightarrow (x-5)(x+2) = 0 \Rightarrow x = 5$ or $x = -2$

17. Continuous everywhere. ($|x-1| + \sin x$ defined for all x; limits exist and are equal to function values.)

18. Continuous everywhere. ($|x| + 1 \neq 0$ for all x; limits exist and are equal to function values.)

19. Discontinuous only at $x = 0$

20. Discontinuous at odd integer multiples of $\frac{\pi}{2}$, i.e., $x = (2n-1)\frac{\pi}{2}$, n an integer, but continuous at all other x.

21. Discontinuous when $2x$ is an integer multiple of π, i.e., $2x = n\pi$, n an integer $\Rightarrow x = \frac{n\pi}{2}$, n an integer, but continuous at all other x.

22. Discontinuous when $\frac{\pi x}{2}$ is an odd integer multiple of $\frac{\pi}{2}$, i.e., $\frac{\pi x}{2} = (2n-1)\frac{\pi}{2}$, n an integer $\Rightarrow x = 2n-1$, n an integer (i.e., x is an odd integer). Continuous everywhere else.

23. Discontinuous at odd integer multiples of $\frac{\pi}{2}$, i.e., $x = (2n-1)\frac{\pi}{2}$, n an integer, but continuous at all other x.

24. Continuous everywhere since $-1 \le \sin x \le 1 \Rightarrow 0 \le \sin^2 x \le 1 \Rightarrow 1 + \sin^2 x \ge 1$; limits exist and are equal to the function values.

25. Discontinuous when $2x + 3 < 0$ or $x < -\frac{3}{2} \Rightarrow$ continuous on the interval $\left[-\frac{3}{2}, \infty\right)$.

26. Discontinuous when $3x - 1 < 0$ or $x < \frac{1}{3} \Rightarrow$ continuous on the interval $\left[\frac{1}{3}, \infty\right)$.

27. Continuous everywhere: $(2x-1)^{1/3}$ is defined for all x; limits exist and are equal to function values.

28. Continuous everywhere: $(2-x)^{1/5}$ is defined for all x; limits exist and are equal to function values.

29. Continuous everywhere since $\lim_{x\to 3} \frac{x^2-x-6}{x-3} = \lim_{x\to 3} \frac{(x-3)(x+2)}{x-3} = \lim_{x\to 3} (x+2) = 5 = g(3)$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

30. Discontinuous at $x = -2$ since $\lim_{x\to -2} f(x)$ does not exist while $f(-2) = 4$.

31. Discontinuous at $x = 1$; $\lim_{x\to 1^+} (x^2 + 2) = 3$, but $\lim_{x\to 1^-} e^x = e$, so that $\lim_{x\to 1} f(x)$ does not exist while $f(1) = e$; and $\lim_{x\to 0^-} (1 - x) = 1 = \lim_{x\to 0^+} e^x$, so that $\lim_{x\to 0} f(x) = 1 = f(0)$

32. Discontinuous at $x = \ln 2$, since $2 - e^x = 0 \Rightarrow e^x = 2 \Rightarrow \ln e^x = \ln 2 \Rightarrow x = \ln 2$

33. $\lim_{x\to\pi} \sin(x - \sin x) = \sin(\pi - \sin\pi) = \sin(\pi - 0) = \sin\pi = 0$, and function continuous at $x = \pi$.

34. $\lim_{t\to 0} \sin(\frac{\pi}{2}\cos(\tan t)) = \sin(\frac{\pi}{2}\cos(\tan(0))) = \sin\left(\frac{\pi}{2}\cos(0)\right) = \sin\left(\frac{\pi}{2}\right) = 1$, and function continuous at $t = 0$.

35. $\lim_{y\to 1} \sec(y\sec^2 y - \tan^2 y - 1) = \lim_{y\to 1} \sec(y\sec^2 y - \sec^2 y) = \lim_{y\to 1} \sec((y-1)\sec^2 y) = \sec((1-1)\sec^2 1) = \sec 0 = 1$, and function continuous at $y = 1$.

36. $\lim_{x\to 0} \tan\left[\frac{\pi}{4}\cos(\sin x^{1/3})\right] = \tan\left[\frac{\pi}{4}\cos(\sin(0))\right] = \tan\left(\frac{\pi}{4}\cos(0)\right) = \tan\left(\frac{\pi}{4}\right) = 1$, and function continuous at $x = 0$.

37. $\lim_{t\to 0} \cos\left[\frac{\pi}{\sqrt{19 - 3\sec 2t}}\right] = \cos\left[\frac{\pi}{\sqrt{19 - 3\sec 0}}\right] = \cos\frac{\pi}{\sqrt{16}} = \cos\frac{\pi}{4} = \frac{\sqrt{2}}{2}$, and function continuous at $t = 0$.

38. $\lim_{x\to \frac{\pi}{6}} \sqrt{\csc^2 x + 5\sqrt{3}\tan x} = \sqrt{\csc^2\left(\frac{\pi}{6}\right) + 5\sqrt{3}\tan\left(\frac{\pi}{6}\right)} = \sqrt{4 + 5\sqrt{3}\left(\frac{1}{\sqrt{3}}\right)} = \sqrt{9} = 3$, and function continuous at $x = \frac{\pi}{6}$.

39. $\lim_{x\to 0^+} \sin\left(\frac{\pi}{2}e^{\sqrt{x}}\right) = \sin\left(\frac{\pi}{2}e^0\right) = \sin\left(\frac{\pi}{2}\right) = 1$, and the function is continuous at $x = 0$.

40. $\lim_{x\to 1} \cos^{-1}\left(\ln\sqrt{x}\right) = \cos^{-1}\left(\ln\sqrt{1}\right) = \cos^{-1}(0) = \frac{\pi}{2}$, and the function is continuous at $x = 1$.

41. $g(x) = \frac{x^2-9}{x-3} = \frac{(x+3)(x-3)}{(x-3)} = x + 3,\ x \neq 3 \Rightarrow g(3) = \lim_{x\to 3} (x + 3) = 6$

42. $h(t) = \frac{t^2+3t-10}{t-2} = \frac{(t+5)(t-2)}{t-2} = t + 5,\ t \neq 2 \Rightarrow h(2) = \lim_{t\to 2} (t + 5) = 7$

43. $f(s) = \frac{s^3-1}{s^3-1} = \frac{(s^2+s+1)(s-1)}{(s+1)(s-1)} = \frac{s^2+s+1}{s+1},\ s \neq 1 \Rightarrow f(1) = \lim_{s\to 1}\left(\frac{s^2+s+1}{s+1}\right) = \frac{3}{2}$

44. $g(x) = \frac{x^2-16}{x^2-3x-4} = \frac{(x+4)(x-4)}{(x-4)(x+1)} = \frac{x+4}{x+1},\ x \neq 4 \Rightarrow g(4) = \lim_{x\to 4}\left(\frac{x+4}{x+1}\right) = \frac{8}{5}$

45. As defined, $\lim_{x\to 3^-} f(x) = (3)^2 - 1 = 8$ and $\lim_{x\to 3^+} (2a)(3) = 6a$. For $f(x)$ to be continuous we must have $6a = 8 \Rightarrow a = \frac{4}{3}$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

46. As defined, $\lim_{x\to -2^-} g(x) = -2$ and $\lim_{x\to -2^+} g(x) = b(-2)^2 = 4b$. For $g(x)$ to be continuous we must have $4b = -2 \Rightarrow b = -\frac{1}{2}$.

47. As defined, $\lim_{x\to 2^-} f(x) = 12$ and $\lim_{x\to 2^+} f(x) = a^2(2) - 2a = 2a^2 - 2a$. For $f(x)$ to be continuous we must have $12 = 2a^2 - 2a \Rightarrow a = 3$ or $a = -2$.

48. As defined, $\lim_{x\to 0^-} g(x) = \frac{0-b}{b+1} = \frac{-b}{b+1}$ and $\lim_{x\to 0^+} g(x) = (0)^2 + b = b$. For $g(x)$ to be continuous we must have $\frac{-b}{b+1} = b \Rightarrow b = 0$ or $b = -2$.

49. As defined, $\lim_{x\to -1^-} f(x) = -2$ and $\lim_{x\to -1^+} f(x) = a(-1) + b = -a + b$, and $\lim_{x\to 1^-} f(x) = a(1) + b = a + b$ and $\lim_{x\to 1^+} f(x) = 3$. For $f(x)$ to be continuous we must have $-2 = -a + b$ and $a + b = 3 \Rightarrow a = \frac{5}{2}$ and $b = \frac{1}{2}$.

50. As defined, $\lim_{x\to 0^-} g(x) = a(0) + 2b = 2b$ and $\lim_{x\to 0^+} g(x) = (0)^2 + 3a - b = 3a - b$, and $\lim_{x\to 2^-} g(x) = (2)^2 + 3a - b = 4 + 3a - b$ and $\lim_{x\to 0^+} g(x) = 3(2) - 5 = 1$. For $g(x)$ to be continuous we must have $2b = 3a - b$ and $4 + 3a - b = 1 \Rightarrow a = -\frac{3}{2}$ and $b = -\frac{3}{2}$.

51. The function can be extended: $f(0) \approx 2.3$.



52. The function cannot be extended to be continuous at $x = 0$. If $f(0) \approx 2.3$, it will be continuous from the right. Or if $f(0) \approx -2.3$, it will be continuous from the left.



53. The function cannot be extended to be continuous at $x = 0$. If $f(0) = 1$, it will be continuous from the right. Or if $f(0) = -1$, it will be continuous from the left.



54. The function can be extended: $f(0) \approx 7.39$.



Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

55. $f(x)$ is continuous on $[0, 1]$ and $f(0) < 0$, $f(1) > 0 \Rightarrow$ by the Intermediate Value Theorem $f(x)$ takes on every value between $f(0)$ and $f(1) \Rightarrow$ the equation $f(x) = 0$ has at least one solution between $x = 0$ and $x = 1$.



56. $\cos x = x \Rightarrow (\cos x) - x = 0$. If $x = -\frac{\pi}{2}$, $\cos\left(-\frac{\pi}{2}\right) - \left(-\frac{\pi}{2}\right) > 0$. If $x = \frac{\pi}{2}$, $\cos\left(\frac{\pi}{2}\right) - \frac{\pi}{2} < 0$. Thus $\cos x - x = 0$ for some x between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ according to the Intermediate Value Theorem, since the function $\cos x - x$ is continuous.

57. Let $f(x) = x^3 - 15x + 1$, which is continuous on $[-4, 4]$. Then $f(-4) = -3$, $f(-1) = 15$, $f(1) = -13$, and $f(4) = 5$. By the Intermediate Value Theorem, $f(x) = 0$ for some x in each of the intervals $-4 < x < -1$, $-1 < x < 1$, and $1 < x < 4$. That is, $x^3 - 15x + 1 = 0$ has three solutions in $[-4, 4]$. Since a polynomial of degree 3 can have at most 3 solutions, these are the only solutions.

58. Without loss of generality, assume that $a < b$. Then $F(x) = (x-a)^2(x-b)^2 + x$ is continuous for all values of x, so it is continuous on the interval $[a, b]$. Moreover $F(a) = a$ and $F(b) = b$. By the Intermediate Value Theorem, since $a < \frac{a+b}{2} < b$, there is a number c between a and b such that $F(x) = \frac{a+b}{2}$.

59. Answers may vary. Note that f is continuous for every value of x.
    (a) $f(0) = 10$, $f(1) = 1^3 - 8(1) + 10 = 3$. Since $3 < \pi < 10$, by the Intermediate Value Theorem, there exists a c so that $0 < c < 1$ and $f(c) = \pi$.
    (b) $f(0) = 10$, $f(-4) = (-4)^3 - 8(-4) + 10 = -22$. Since $-22 < -\sqrt{3} < 10$, by the Intermediate Value Theorem, there exists a c so that $-4 < c < 0$ and $f(c) = -\sqrt{3}$.
    (c) $f(0) = 10$, $f(1000) = (1000)^3 - 8(1000) + 10 = 999,992,010$. Since $10 < 5,000,000 < 999,992,010$, by the Intermediate Value Theorem, there exists a c so that $0 < c < 1000$ and $f(c) = 5,000,000$.

60. All five statements ask for the same information because of the intermediate value property of continuous functions.
    (a) A root of $f(x) = x^3 - 3x - 1$ is a point c where $f(c) = 0$.
    (b) The point where $y = x^3$ crosses $y = 3x + 1$ have the same y-coordinate, or $y = x^3 = 3x + 1 \Rightarrow f(x) = x^3 - 3x - 1 = 0$.
    (c) $x^3 - 3x = 1 \Rightarrow x^3 - 3x - 1 = 0$. The solutions to the equation are the roots of $f(x) = x^3 - 3x - 1$.
    (d) The points where $y = x^3 - 3x$ crosses $y = 1$ have common y-coordinates, or $y = x^3 - 3x = 1 \Rightarrow f(x) = x^3 - 3x - 1 = 0$.
    (e) The solutions of $x^3 - 3x - 1 = 0$ are those points where $f(x) = x^3 - 3x - 1$ has value 0.

61. Answers may vary. For example, $f(x) = \frac{\sin(x-2)}{x-2}$ is discontinuous at $x = 2$ because it is not defined there. However, the discontinuity can be removed because f has a limit (namely 1) as $x \to 2$.

62. Answers may vary. For example, $g(x) = \frac{1}{x+1}$ has a discontinuity at $x = -1$ because $\lim_{x \to -1} g(x)$ does not exist. $\left(\lim_{x \to -1^-} g(x) = -\infty \text{ and } \lim_{x \to -1^+} g(x) = +\infty.\right)$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

63. (a) Suppose x_0 is rational $\Rightarrow f(x_0) = 1$. Choose $\epsilon = \frac{1}{2}$. For any $\delta > 0$ there is an irrational number x (actually infinitely many) in the interval $(x_0 - \delta, x_0 + \delta) \Rightarrow f(x) = 0$. Then $0 < |x - x_0| < \delta$ but $|f(x) - f(x_0)| = 1 > \frac{1}{2} = \epsilon$, so $\lim_{x \to x_0} f(x)$ fails to exist $\Rightarrow f$ is discontinuous at x_0 rational.
On the other hand, x_0 irrational $\Rightarrow f(x_0) = 0$ and there is a rational number x in $(x_0 - \delta, x_0 + \delta) \Rightarrow f(x) = 1$. Again $\lim_{x \to x_0} f(x)$ fails to exist $\Rightarrow f$ is discontinuous at x_0 irrational. That is, f is discontinuous at every point.

(b) f is neither right-continuous nor left-continuous at any point x_0 because in every interval $(x_0 - \delta, x_0)$ or $(x_0, x_0 + \delta)$ there exist both rational and irrational real numbers. Thus neither limits $\lim_{x \to x_0^-} f(x)$ and $\lim_{x \to x_0^+} f(x)$ exist by the same arguments used in part (a).

64. Yes. Both $f(x) = x$ and $g(x) = x - \frac{1}{2}$ are continuous on $[0, 1]$. However $\frac{f(x)}{g(x)}$ is undefined at $x = \frac{1}{2}$ since $g\left(\frac{1}{2}\right) = 0 \Rightarrow \frac{f(x)}{g(x)}$ is discontinuous at $x = \frac{1}{2}$.

65. No. For instance, if $f(x) = 0$, $g(x) = \lceil x \rceil$, then $h(x) = 0\left(\lceil x \rceil\right) = 0$ is continuous at $x = 0$ and $g(x)$ is not.

66. Let $f(x) = \frac{1}{x-1}$ and $g(x) = x + 1$. Both functions are continuous at $x = 0$. The composition $f \circ g = f(g(x)) = \frac{1}{(x+1)-1} = \frac{1}{x}$ is discontinuous at $x = 0$, since it is not defined there. Theorem 10 requires that $f(x)$ be continuous at $g(0)$, which is not the case here since $g(0) = 1$ and f is undefined at 1.

67. Yes, because of the Intermediate Value Theorem. If $f(a)$ and $f(b)$ did have different signs then f would have to equal zero at some point between a and b since f is continuous on $[a, b]$.

68. Let $f(x)$ be the new position of point x and let $d(x) = f(x) - x$. The displacement function d is negative if x is the left-hand point of the rubber band and positive if x is the right-hand point of the rubber band. By the Intermediate Value Theorem, $d(x) = 0$ for some point in between. That is, $f(x) = x$ for some point x, which is then in its original position.

69. If $f(0) = 0$ or $f(1) = 1$, we are done (i.e., $c = 0$ or $c = 1$ in those cases). Then let $f(0) = a > 0$ and $f(1) = b < 1$ because $0 \le f(x) \le 1$. Define $g(x) = f(x) - x \Rightarrow g$ is continuous on $[0, 1]$. Moreover, $g(0) = f(0) - 0 = a > 0$ and $g(1) = f(1) - 1 = b - 1 < 0 \Rightarrow$ by the Intermediate Value Theorem there is a number c in $(0, 1)$ such that $g(c) = 0 \Rightarrow f(c) - c = 0$ or $f(c) = c$.

70. Let $\epsilon = \frac{|f(c)|}{2} > 0$. Since f is continuous at $x = c$ there is a $\delta > 0$ such that $|x - c| < \delta \Rightarrow |f(x) - f(c)| < \epsilon$ $\Rightarrow f(c) - \epsilon < f(x) < f(c) + \epsilon$.
If $f(c) > 0$, then $\epsilon = \frac{1}{2} f(c) \Rightarrow \frac{1}{2} f(c) < f(x) < \frac{3}{2} f(c) \Rightarrow f(x) > 0$ on the interval $(c - \delta, c + \delta)$.
If $f(c) < 0$, then $\epsilon = -\frac{1}{2} f(c) \Rightarrow \frac{3}{2} f(c) < f(x) < \frac{1}{2} f(c) \Rightarrow f(x) < 0$ on the interval $(c - \delta, c + \delta)$.





Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

71. By Exercise 52 in Section 2.3, we have $\lim_{x\to c} f(x) = L \Leftrightarrow \lim_{h\to 0} f(c+h) = L$.
Thus, $f(x)$ is continuous at $x = c \Leftrightarrow \lim_{x\to c} f(x) = f(c) \Leftrightarrow \lim_{h\to 0} f(c+h) = f(c)$.

72. By Exercise 71, it suffices to show that $\lim_{h\to 0} \sin(c+h) = \sin c$ and $\lim_{h\to 0} \cos(c+h) = \cos c$.

Now $\lim_{h\to 0} \sin(c+h) = \lim_{h\to 0} \left[(\sin c)(\cos h) + (\cos c)(\sin h)\right] = (\sin c)\left(\lim_{h\to 0} \cos h\right) + (\cos c)\left(\lim_{h\to 0} \sin h\right)$.

By Example 11 Section 2.2, $\lim_{h\to 0} \cos h = 1$ and $\lim_{h\to 0} \sin h = 0$. So $\lim_{h\to 0} \sin(c+h) = \sin c$ and thus $f(x) = \sin x$ is continuous at $x = c$. Similarly,

$\lim_{h\to 0} \cos(c+h) = \lim_{h\to 0} \left[(\cos c)(\cos h) - (\sin c)(\sin h)\right] = (\cos c)\left(\lim_{h\to 0} \cos h\right) - (\sin c)\left(\lim_{h\to 0} \sin h\right) = \cos c$. Thus, $g(x) = \cos x$ is continuous at $x = c$.

73. $x \approx 1.8794, -1.5321, -0.3473$

74. $x \approx 1.4516, -0.8547, 0.4030$

75. $x \approx 1.7549$

76. $x \approx 1.5596$

77. $x \approx 3.5156$

78. $x \approx -3.9058, 3.8392, 0.0667$

79. $x \approx 0.7391$

80. $x \approx -1.8955, 0, 1.8955$

2.6 LIMITS INVOLVING INFINITY; ASYMPTOTES OF GRAPHS

1. (a) $\lim_{x\to 2} f(x) = 0$
(b) $\lim_{x\to -3^+} f(x) = -2$
(c) $\lim_{x\to -3^-} f(x) = 2$
(d) $\lim_{x\to 3} f(x)$ = does not exist
(e) $\lim_{x\to 0^+} f(x) = -1$
(f) $\lim_{x\to 0^-} f(x) = +\infty$
(g) $\lim_{x\to 0} f(x)$ = does not exist
(h) $\lim_{x\to \infty} f(x) = 1$
(i) $\lim_{x\to -\infty} f(x) = 0$

2. (a) $\lim_{x\to 4} f(x) = 2$
(b) $\lim_{x\to 2^+} f(x) = -3$
(c) $\lim_{x\to 2^-} f(x) = 1$
(d) $\lim_{x\to 2} f(x)$ = does not exist
(e) $\lim_{x\to -3^+} f(x) = +\infty$
(f) $\lim_{x\to -3^-} f(x) = +\infty$
(g) $\lim_{x\to -3} f(x) = +\infty$
(h) $\lim_{x\to 0^+} f(x) = +\infty$
(i) $\lim_{x\to 0^-} f(x) = -\infty$
(j) $\lim_{x\to 0} f(x)$ = does not exist
(k) $\lim_{x\to \infty} f(x) = 0$
(l) $\lim_{x\to -\infty} f(x) = -1$

Note: In these exercises we use the result $\lim_{x\to \pm\infty} \frac{1}{x^{m/n}} = 0$ whenever $\frac{m}{n} > 0$. This result follows immediately from Theorem 8 and the power rule in Theorem 1: $\lim_{x\to \pm\infty} \left(\frac{1}{x^{m/n}}\right) = \lim_{x\to \pm\infty} \left(\frac{1}{x}\right)^{m/n} = \left(\lim_{x\to \pm\infty} \frac{1}{x}\right)^{m/n} = 0^{m/n} = 0$.

3. (a) -3
(b) -3

4. (a) π
(b) π

5. (a) $\frac{1}{2}$
(b) $\frac{1}{2}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

6. (a) $\frac{1}{8}$ (b) $\frac{1}{8}$

7. (a) $-\frac{5}{3}$ (b) $-\frac{5}{3}$

8. (a) $\frac{3}{4}$ (b) $\frac{3}{4}$

9. $-\frac{1}{x} \le \frac{\sin 2x}{x} \le \frac{1}{x} \Rightarrow \lim_{x\to\infty} \frac{\sin 2x}{x} = 0$ by the Sandwich Theorem

10. $-\frac{1}{3\theta} \le \frac{\cos\theta}{3\theta} \le \frac{1}{3\theta} \Rightarrow \lim_{\theta\to-\infty} \frac{\cos\theta}{3\theta} = 0$ by the Sandwich Theorem

11. $\lim_{t\to\infty} \frac{2-t+\sin t}{t+\cos t} = \lim_{t\to\infty} \frac{\frac{2}{t}-1+\left(\frac{\sin t}{t}\right)}{1+\left(\frac{\cos t}{t}\right)} = \frac{0-1+0}{1+0} = -1$

12. $\lim_{r\to\infty} \frac{r+\sin r}{2r+7-5\sin r} = \lim_{r\to\infty} \frac{1+\left(\frac{\sin r}{r}\right)}{2+\frac{7}{r}-5\left(\frac{\sin r}{r}\right)} = \lim_{r\to\infty} \frac{1+0}{2+0-0} = \frac{1}{2}$

13. (a) $\lim_{x\to\infty} \frac{2x+3}{5x+7} = \lim_{x\to\infty} \frac{2+\frac{3}{x}}{5+\frac{7}{x}} = \frac{2}{5}$ (b) $\frac{2}{5}$ (same process as part (a))

14. (a) $\lim_{x\to\infty} \frac{2x^3+7}{x^3-x^2+x+7} = \lim_{x\to\infty} \frac{2+\left(\frac{7}{x^3}\right)}{1-\frac{1}{x}+\frac{1}{x^2}+\frac{7}{x^3}} = 2$

(b) 2 (same process as part (a))

15. (a) $\lim_{x\to\infty} \frac{x+1}{x^2+3} = \lim_{x\to\infty} \frac{\frac{1}{x}+\frac{1}{x^2}}{1+\frac{3}{x^2}} = 0$ (b) 0 (same process as part (a))

16. (a) $\lim_{x\to\infty} \frac{3x+7}{x^2-2} = \lim_{x\to\infty} \frac{\frac{3}{x}+\frac{7}{x^2}}{1+\frac{2}{x^2}} = 0$ (b) 0 (same process as part (a))

17. (a) $\lim_{x\to\infty} \frac{7x^3}{x^3-3x^2+6x} = \lim_{x\to\infty} \frac{7}{1-\frac{3}{x}+\frac{9}{x^2}} = 7$ (b) 7 (same process as part (a))

18. (a) $\lim_{x\to\infty} \frac{9x^4+x}{2x^4+5x^2-x+6} = \lim_{x\to\infty} \frac{9+\frac{1}{x^3}}{2+\frac{5}{x^2}-\frac{1}{x^3}+\frac{6}{x^4}} = \frac{9}{2}$ (b) $\frac{9}{2}$ (same process as part (a))

19. (a) $\lim_{x\to\infty} \frac{10x^5+x^4+31}{x^6} = \lim_{x\to\infty} \frac{\frac{10}{x}+\frac{1}{x^2}+\frac{31}{x^6}}{1} = 0$ (b) 0 (same process as part (a))

20. (a) $\lim_{x\to\infty} \frac{x^3+7x^2-2}{x^2-x-1} = \lim_{x\to\infty} \frac{x+7-2x^{-1}}{1-x^{-1}-x^{-2}} = \infty$, since $x^{-n} \to 0$ and $x+7 \to \infty$.

(b) $\lim_{x\to-\infty} \frac{x^3+7x^2-2}{x^2-x-1} = \lim_{x\to-\infty} \frac{x+7-2x^{-1}}{1-x^{-1}-x^{-2}} = -\infty$, since $x^{-n} \to 0$ and $x+7 \to -\infty$.

21. (a) $\lim_{x\to\infty} \frac{3x^7+5x^2-1}{6x^3-7x+3} = \lim_{x\to\infty} \frac{3x^4+5x^{-1}-x^{-3}}{6-7x^{-2}+3x^{-3}} = \infty$, since $x^{-n} \to 0$ and $3x^4 \to \infty$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b) $\lim\limits_{x\to-\infty}\dfrac{3x^7+5x^2-1}{6x^3-7x+3}=\lim\limits_{x\to-\infty}\dfrac{3x^4+5x^{-1}-x^{-3}}{6-7x^{-2}+3x^{-3}}=\infty$, since $x^{-n}\to 0$ and $3x^4\to\infty$.

22. (a) $\lim\limits_{x\to\infty}\dfrac{5x^8-2x^3+9}{3+x-4x^5}=\lim\limits_{x\to\infty}\dfrac{5x^3-2x^{-2}+9x^{-5}}{3x^{-5}+x^{-4}-4}=-\infty$, since $x^{-n}\to 0$, $5x^3\to\infty$, and the denominator $\to -4$.

(b) $\lim\limits_{x\to-\infty}\dfrac{5x^8-2x^3+9}{3+x-4x^5}=\lim\limits_{x\to-\infty}\dfrac{5x^3-2x^{-2}+9x^{-5}}{3x^{-5}+x^{-4}-4}=\infty$, since $x^{-n}\to 0$, $5x^3\to-\infty$, and the denominator $\to -4$.

23. $\lim\limits_{x\to\infty}\sqrt{\dfrac{8x^2-3}{2x^2+x}}=\lim\limits_{x\to\infty}\sqrt{\dfrac{8-\frac{3}{x^2}}{2+\frac{1}{x}}}=\sqrt{\lim\limits_{x\to\infty}\dfrac{8-\frac{3}{x^2}}{2+\frac{1}{x}}}=\sqrt{\dfrac{8-0}{2+0}}=\sqrt{4}=2$

24. $\lim\limits_{x\to-\infty}\left(\dfrac{x^2+x-1}{8x^2-3}\right)^{1/3}=\lim\limits_{x\to-\infty}\left(\dfrac{1+\frac{1}{x}-\frac{1}{x^2}}{8-\frac{3}{x^2}}\right)^{1/3}=\left(\lim\limits_{x\to-\infty}\dfrac{1+\frac{1}{x}-\frac{1}{x^2}}{8-\frac{3}{x^2}}\right)^{1/3}=\left(\dfrac{1+0-0}{8-0}\right)^{1/3}=\left(\dfrac{1}{8}\right)^{1/3}=\dfrac{1}{2}$

25. $\lim\limits_{x\to-\infty}\left(\dfrac{1-x^3}{x^2-7x}\right)^5=\lim\limits_{x\to-\infty}\left(\dfrac{\frac{1}{x^2}-x}{1-\frac{7}{x}}\right)^5=\left(\lim\limits_{x\to-\infty}\dfrac{\frac{1}{x^2}-x}{1-\frac{7}{x}}\right)^5=\left(\dfrac{0+\infty}{1-0}\right)^5=\infty$

26. $\lim\limits_{x\to\infty}\sqrt{\dfrac{x^2-5x}{x^3+x-2}}=\lim\limits_{x\to\infty}\sqrt{\dfrac{\frac{1}{x}-\frac{5}{x^2}}{1+\frac{1}{x^2}-\frac{2}{x^3}}}=\sqrt{\lim\limits_{x\to\infty}\dfrac{\frac{1}{x}-\frac{5}{x^2}}{1+\frac{1}{x^2}-\frac{2}{x^3}}}=\sqrt{\dfrac{0-0}{1+0-0}}=\sqrt{0}=0$

27. $\lim\limits_{x\to\infty}\dfrac{2\sqrt{x}+x^{-1}}{3x-7}=\lim\limits_{x\to\infty}\dfrac{\left(\frac{2}{x^{1/2}}\right)+\left(\frac{1}{x^2}\right)}{3-\frac{7}{x}}=0$

28. $\lim\limits_{x\to\infty}\dfrac{2+\sqrt{x}}{2-\sqrt{x}}=\lim\limits_{x\to\infty}\dfrac{\left(\frac{2}{x^{1/2}}\right)+1}{\left(\frac{2}{x^{1/2}}\right)-1}=-1$

29. $\lim\limits_{x\to-\infty}\dfrac{\sqrt[3]{x}-\sqrt[5]{x}}{\sqrt[3]{x}+\sqrt[5]{x}}=\lim\limits_{x\to-\infty}\dfrac{1-x^{(1/5)-(1/3)}}{1+x^{(1/5)-(1/3)}}=\lim\limits_{x\to-\infty}\dfrac{1-\left(\frac{1}{x^{2/15}}\right)}{1+\left(\frac{1}{x^{2/15}}\right)}=1$

30. $\lim\limits_{x\to\infty}\dfrac{x^{-1}+x^{-4}}{x^{-2}-x^{-3}}=\lim\limits_{x\to\infty}\dfrac{x+\frac{1}{x^2}}{1-\frac{1}{x}}=\infty$

31. $\lim\limits_{x\to\infty}\dfrac{2x^{5/3}-x^{1/3}+7}{x^{8/5}+3x+\sqrt{x}}=\lim\limits_{x\to\infty}\dfrac{2x^{1/15}-\frac{1}{x^{19/15}}+\frac{7}{x^{8/5}}}{1+\frac{3}{x^{3/5}}+\frac{1}{x^{11/10}}}=\infty$

32. $\lim\limits_{x\to-\infty}\dfrac{\sqrt[3]{x}-5x+3}{2x+x^{2/3}-4}=\lim\limits_{x\to-\infty}\dfrac{\frac{1}{x^{2/3}}-5+\frac{3}{x}}{2+\frac{1}{x^{1/3}}-\frac{4}{x}}=-\dfrac{5}{2}$

33. $\lim\limits_{x\to\infty}\dfrac{\sqrt{x^2+1}}{x+1}=\lim\limits_{x\to\infty}\dfrac{\sqrt{x^2+1}/\sqrt{x^2}}{(x+1)/\sqrt{x^2}}=\lim\limits_{x\to\infty}\dfrac{\sqrt{(x^2+1)/x^2}}{(x+1)/x}=\lim\limits_{x\to\infty}\dfrac{\sqrt{1+1/x^2}}{(1+1/x)}=\dfrac{\sqrt{1+0}}{(1+0)}=1$

34. $\lim\limits_{x\to-\infty}\dfrac{\sqrt{x^2+1}}{x+1}=\lim\limits_{x\to-\infty}\dfrac{\sqrt{x^2+1}/\sqrt{x^2}}{(x+1)/\sqrt{x^2}}=\lim\limits_{x\to-\infty}\dfrac{\sqrt{(x^2+1)/x^2}}{(x+1)/(-x)}=\lim\limits_{x\to\infty}\dfrac{\sqrt{1+1/x^2}}{(-1-1/x)}=\dfrac{\sqrt{1+0}}{(-1-0)}=-1$

35. $\lim\limits_{x\to\infty}\dfrac{x-3}{\sqrt{4x^2+25}}=\lim\limits_{x\to\infty}\dfrac{(x-3)/\sqrt{x^2}}{\sqrt{4x^2+25}/\sqrt{x^2}}=\lim\limits_{x\to\infty}\dfrac{(x-3)/x}{\sqrt{(4x^2+25)/x^2}}=\lim\limits_{x\to\infty}\dfrac{(1-3/x)}{\sqrt{4+25/x^2}}=\dfrac{(1-0)}{\sqrt{4+0}}=\dfrac{1}{2}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

36. $\lim\limits_{x\to-\infty}\dfrac{4-3x^3}{\sqrt{x^6+9}}=\lim\limits_{x\to-\infty}\dfrac{(4-3x^3)/\sqrt{x^6}}{\sqrt{x^6+9}/\sqrt{x^6}}=\lim\limits_{x\to-\infty}\dfrac{(4-3x^3)/(-x^3)}{\sqrt{(x^6+9)/x^6}}=\lim\limits_{x\to\infty}\dfrac{(-4/x^3+3)}{\sqrt{1+9/x^6}}=\dfrac{(0+3)}{\sqrt{1+0}}=3$

37. $\lim\limits_{x\to0^+}\dfrac{1}{3x}=\infty$ $\left(\dfrac{\text{positive}}{\text{positive}}\right)$

38. $\lim\limits_{x\to0^-}\dfrac{5}{2x}=-\infty$ $\left(\dfrac{\text{positive}}{\text{negative}}\right)$

39. $\lim\limits_{x\to2^-}\dfrac{3}{x-2}=-\infty$ $\left(\dfrac{\text{positive}}{\text{negative}}\right)$

40. $\lim\limits_{x\to3^+}\dfrac{1}{x-3}=\infty$ $\left(\dfrac{\text{positive}}{\text{positive}}\right)$

41. $\lim\limits_{x\to-8^+}\dfrac{2x}{x+8}=-\infty$ $\left(\dfrac{\text{negative}}{\text{positive}}\right)$

42. $\lim\limits_{x\to-5^-}\dfrac{3x}{2x+10}=\infty$ $\left(\dfrac{\text{negative}}{\text{negative}}\right)$

43. $\lim\limits_{x\to7}\dfrac{4}{(x-7)^2}=\infty$ $\left(\dfrac{\text{positive}}{\text{positive}}\right)$

44. $\lim\limits_{x\to0}\dfrac{-1}{x^2(x+1)}=-\infty$ $\left(\dfrac{\text{negative}}{\text{positive}\cdot\text{positive}}\right)$

45. (a) $\lim\limits_{x\to0^+}\dfrac{2}{3x^{1/3}}=\infty$ (b) $\lim\limits_{x\to0^-}\dfrac{2}{3x^{1/3}}=-\infty$

46. (a) $\lim\limits_{x\to0^+}\dfrac{2}{x^{1/5}}=\infty$ (b) $\lim\limits_{x\to0^-}\dfrac{2}{x^{1/5}}=-\infty$

47. $\lim\limits_{x\to0}\dfrac{4}{x^{2/5}}=\lim\limits_{x\to0}\dfrac{4}{(x^{1/5})^2}=\infty$

48. $\lim\limits_{x\to0}\dfrac{1}{x^{2/3}}=\lim\limits_{x\to0}\dfrac{1}{(x^{1/3})^2}=\infty$

49. $\lim\limits_{x\to\left(\frac{\pi}{2}\right)^-}\tan x=\infty$

50. $\lim\limits_{x\to\left(\frac{-\pi}{2}\right)^+}\sec x=\infty$

51. $\lim\limits_{\theta\to0^-}(1+\csc\theta)=-\infty$

52. $\lim\limits_{\theta\to0^+}(2-\cot\theta)=-\infty$ and $\lim\limits_{\theta\to0^-}(2-\cot\theta)=\infty$, so the limit does not exist

53. (a) $\lim\limits_{x\to2^+}\dfrac{1}{x^2-4}=\lim\limits_{x\to2^+}\dfrac{1}{(x+2)(x-2)}=\infty$ $\left(\dfrac{1}{\text{positive}\cdot\text{positive}}\right)$

(b) $\lim\limits_{x\to2^-}\dfrac{1}{x^2-4}=\lim\limits_{x\to2^-}\dfrac{1}{(x+2)(x-2)}=-\infty$ $\left(\dfrac{1}{\text{positive}\cdot\text{negative}}\right)$

(c) $\lim\limits_{x\to-2^+}\dfrac{1}{x^2-4}=\lim\limits_{x\to-2^+}\dfrac{1}{(x+2)(x-2)}=-\infty$ $\left(\dfrac{1}{\text{positive}\cdot\text{negative}}\right)$

(d) $\lim\limits_{x\to-2^-}\dfrac{1}{x^2-4}=\lim\limits_{x\to-2^-}\dfrac{1}{(x+2)(x-2)}=\infty$ $\left(\dfrac{1}{\text{negative}\cdot\text{negative}}\right)$

54. (a) $\lim\limits_{x\to1^+}\dfrac{x}{x^2-1}=\lim\limits_{x\to1^+}\dfrac{x}{(x+1)(x-1)}=\infty$ $\left(\dfrac{\text{positive}}{\text{positive}\cdot\text{positive}}\right)$

(b) $\lim\limits_{x\to1^-}\dfrac{x}{x^2-1}=\lim\limits_{x\to1^-}\dfrac{x}{(x+1)(x-1)}=-\infty$ $\left(\dfrac{\text{positive}}{\text{positive}\cdot\text{negative}}\right)$

(c) $\lim\limits_{x\to-1^+}\dfrac{x}{x^2-1}=\lim\limits_{x\to-1^+}\dfrac{x}{(x+1)(x-1)}=\infty$ $\left(\dfrac{\text{negative}}{\text{positive}\cdot\text{negative}}\right)$

(d) $\lim\limits_{x\to-1^-}\dfrac{x}{x^2-1}=\lim\limits_{x\to-1^-}\dfrac{x}{(x+1)(x-1)}=-\infty$ $\left(\dfrac{\text{negative}}{\text{negative}\cdot\text{negative}}\right)$

55. (a) $\lim\limits_{x\to0^+}\left(\dfrac{x^2}{2}-\dfrac{1}{x}\right)=0+\lim\limits_{x\to0^+}\dfrac{1}{-x}=-\infty$ $\left(\dfrac{1}{\text{negative}}\right)$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(b) $\lim_{x\to 0^-}\left(\frac{x^2}{2}-\frac{1}{x}\right)=0+\lim_{x\to 0^-}\frac{1}{-x}=\infty \quad \left(\frac{1}{\text{positive}}\right)$

(c) $\lim_{x\to\sqrt[3]{2}}\left(\frac{x^2}{2}-\frac{1}{x}\right)=\frac{2^{2/3}}{2}-\frac{1}{2^{1/3}}=2^{-1/3}-2^{-1/3}=0$

(d) $\lim_{x\to -1}\left(\frac{x^2}{2}-\frac{1}{x}\right)=\frac{1}{2}-\left(\frac{1}{-1}\right)=\frac{3}{2}$

56. (a) $\lim_{x\to -2^+}\frac{x^2-1}{2x+4}=\infty \quad \left(\frac{\text{positive}}{\text{positive}}\right)$ (b) $\lim_{x\to -2^-}\frac{x^2-1}{2x+4}=-\infty \quad \left(\frac{\text{positive}}{\text{negative}}\right)$

(c) $\lim_{x\to 1^+}\frac{x^2-1}{2x+4}=\lim_{x\to 1^+}\frac{(x+1)(x-1)}{2x+4}=\frac{2\cdot 0}{2+4}=0$

(d) $\lim_{x\to 0^-}\frac{x^2-1}{2x+4}=\frac{-1}{4}$

57. (a) $\lim_{x\to 0^+}\frac{x^2-3x+2}{x^3-2x^2}=\lim_{x\to 0^+}\frac{(x-2)(x-1)}{x^2(x-2)}=-\infty \quad \left(\frac{\text{negative}\cdot\text{negative}}{\text{positive}\cdot\text{negative}}\right)$

(b) $\lim_{x\to 2^+}\frac{x^2-3x+2}{x^3-2x^2}=\lim_{x\to 2^+}\frac{(x-2)(x-1)}{x^2(x-2)}=\lim_{x\to 2^+}\frac{x-1}{x^2}=\frac{1}{4}, x\neq 2$

(c) $\lim_{x\to 2^-}\frac{x^2-3x+2}{x^3-2x^2}=\lim_{x\to 2^-}\frac{(x-2)(x-1)}{x^2(x-2)}=\lim_{x\to 2^-}\frac{x-1}{x^2}=\frac{1}{4}, x\neq 2$

(d) $\lim_{x\to 2}\frac{x^2-3x+2}{x^3-2x^2}=\lim_{x\to 2}\frac{(x-2)(x-1)}{x^2(x-2)}=\lim_{x\to 2}\frac{x-1}{x^2}=\frac{1}{4}, x\neq 2$

(e) $\lim_{x\to 0}\frac{x^2-3x+2}{x^3-2x^2}=\lim_{x\to 0}\frac{(x-2)(x-1)}{x^2(x-2)}=-\infty \quad \left(\frac{\text{negative}\cdot\text{negative}}{\text{positive}\cdot\text{negative}}\right)$

58. (a) $\lim_{x\to 2^+}\frac{x^2-3x+2}{x^3-4x}=\lim_{x\to 2^+}\frac{(x-2)(x-1)}{x(x-2)(x+2)}=\lim_{x\to 2^+}\frac{(x-1)}{x(x+2)}=\frac{1}{2(4)}=\frac{1}{8}$

(b) $\lim_{x\to -2^+}\frac{x^2-3x+2}{x^3-4x}=\lim_{x\to -2^+}\frac{(x-2)(x-1)}{x(x-2)(x+2)}=\lim_{x\to -2^+}\frac{(x-1)}{x(x+2)}=\infty \quad \left(\frac{\text{negative}}{\text{negative}\cdot\text{positive}}\right)$

(c) $\lim_{x\to 0^-}\frac{x^2-3x+2}{x^3-4x}=\lim_{x\to 0^-}\frac{(x-2)(x-1)}{x(x-2)(x+2)}=\lim_{x\to 0^-}\frac{(x-1)}{x(x+2)}=\infty \quad \left(\frac{\text{negative}}{\text{negative}\cdot\text{positive}}\right)$

(d) $\lim_{x\to 1^+}\frac{x^2-3x+2}{x^3-4x}=\lim_{x\to 1^+}\frac{(x-2)(x-1)}{x(x-2)(x+2)}=\lim_{x\to 1^+}\frac{(x-1)}{x(x+2)}=\frac{0}{(1)(3)}=0$

(e) $\lim_{x\to 0^+}\frac{x-1}{x(x+2)}=-\infty \quad \left(\frac{\text{negative}}{\text{positive}\cdot\text{positive}}\right)$

and $\lim_{x\to 0^-}\frac{x-1}{x(x+2)}=\infty \quad \left(\frac{\text{negative}}{\text{negative}\cdot\text{positive}}\right)$

so the function has no limit as $x\to 0$.

59. (a) $\lim_{t\to 0^+}\left[2-\frac{3}{t^{1/3}}\right]=-\infty$ (b) $\lim_{t\to 0^-}\left[2-\frac{3}{t^{1/3}}\right]=\infty$

60. (a) $\lim_{t\to 0^+}\left[\frac{1}{t^{3/5}}+7\right]=\infty$ (b) $\lim_{t\to 0^-}\left[\frac{1}{t^{3/5}}+7\right]=-\infty$

61. (a) $\lim_{x\to 0^+}\left[\frac{1}{x^{2/3}}+\frac{2}{(x-1)^{2/3}}\right]=\infty$ (b) $\lim_{x\to 0^-}\left[\frac{1}{x^{2/3}}+\frac{2}{(x-1)^{2/3}}\right]=\infty$

(c) $\lim_{x\to 1^+}\left[\frac{1}{x^{2/3}}+\frac{2}{(x-1)^{2/3}}\right]=\infty$ (d) $\lim_{x\to 1^-}\left[\frac{1}{x^{2/3}}+\frac{2}{(x-1)^{2/3}}\right]=\infty$

62. (a) $\lim_{x\to 0^+}\left[\frac{1}{x^{1/3}}-\frac{1}{(x-1)^{4/3}}\right]=\infty$ (b) $\lim_{x\to 0^-}\left[\frac{1}{x^{1/3}}-\frac{1}{(x-1)^{4/3}}\right]=-\infty$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(c) $\lim_{x\to 1^+}\left[\frac{1}{x^{1/3}}-\frac{1}{(x-1)^{4/3}}\right]=-\infty$

(d) $\lim_{x\to 1^-}\left[\frac{1}{x^{1/3}}-\frac{1}{(x-1)^{4/3}}\right]=-\infty$

63. $y=\frac{1}{x-1}$



64. $y=\frac{1}{x+1}$



65. $y=\frac{1}{2x+4}$



66. $y=\frac{-3}{x-3}$



67. $y=\frac{x+3}{x+2}=1+\frac{1}{x+2}$



68. $y=\frac{2x}{x+1}=2-\frac{2}{x+1}$



69. domain $=(-\infty,\infty)$; y in range and $y=4+\frac{3x^2}{x^2+1}$, $0\le\frac{3x^2}{x^2+1}<3$ and $\lim_{x\to\pm\infty}\frac{3x^2}{x^2+1}=3\Rightarrow$ range $=[4,7)$; horizontal asymptote is $y=7$.

70. domain $=(-\infty,-1)\cup(-1,1)\cup(1,\infty)$; y in range and $y=\frac{2x}{x^2-1}$; if $x=0$, then $y=0$, $\lim_{x\to\pm\infty}\frac{2x}{x^2-1}=0$, $\lim_{x\to 1^+}\frac{2x}{x^2-1}=+\infty$, and $\lim_{x\to 1^-}\frac{2x}{x^2-1}=-\infty$; $\lim_{x\to -1^+}\frac{2x}{x^2-1}=\infty$ and $\lim_{x\to -1^-}\frac{2x}{x^2-1}=-\infty\Rightarrow$ range $=(-\infty,\infty)$; horizontal asymptote is $y=0$; vertical asymptotes are $x=-1$, $x=1$

71. domain $=(-\infty,\infty)$; y in range and $y=\frac{8-e^x}{2+e^x}$; if $x=\ln 8$, then $y=0$, $-1<\frac{8-e^x}{2+e^x}<4$, $\lim_{x\to\infty}\frac{8-e^x}{2+e^x}=-1$, and $\lim_{x\to-\infty}\frac{8-e^x}{2+e^x}=4\Rightarrow$ range $=(-1,4)$; horizontal asymptotes are $y=-1$, $y=4$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

72. domain $= (-\infty, \infty)$; y in range and $y = \dfrac{4e^x + e^{2x}}{e^x + e^{2x}}$, $1 < \dfrac{4e^x + e^{2x}}{e^x + e^{2x}} < 4$, $\lim\limits_{x \to \infty} \dfrac{4e^x + e^{2x}}{e^x + e^{2x}} = 1$, and $\lim\limits_{x \to -\infty} \dfrac{4e^x + e^{2x}}{e^x + e^{2x}} = 4 \Rightarrow$ range $= (1, 4)$; horizontal asymptotes are $y = 1$, $y = 4$

73. domain $= (-\infty, 0) \cup (0, \infty)$; y in range and $y = \dfrac{\sqrt{x^2+4}}{x}$, if $x > 0$, then $1 < \dfrac{\sqrt{x^2+4}}{x}$, $\lim\limits_{x \to 0^+} \dfrac{\sqrt{x^2+4}}{x} = \infty$, and $\lim\limits_{x \to \infty} \dfrac{\sqrt{x^2+4}}{x} = 1$; if $x < 0$, then $\dfrac{\sqrt{x^2+4}}{x} < -1$, $\lim\limits_{x \to 0^-} \dfrac{\sqrt{x^2+4}}{x} = -\infty$, and $\lim\limits_{x \to -\infty} \dfrac{\sqrt{x^2+4}}{x} = -1 \Rightarrow$ range $= (-\infty, -1) \cup (1, \infty)$; horizontal asymptotes are $y = -1$, $y = 1$; vertical asymptote is $x = 0$

74. domain $= (-\infty, 2) \cup (2, \infty)$; y in range and $y = \dfrac{x^3}{x^3 - 8}$, $\lim\limits_{x \to 2^+} \dfrac{x^3}{x^3-8} = +\infty$, $\lim\limits_{x \to 2^-} \dfrac{x^3}{x^3-8} = -\infty$, $\lim\limits_{x \to \pm\infty} \dfrac{x^3}{x^3-8} = 1$, and $\dfrac{x^3}{x^3-8} \neq 1 \Rightarrow$ range $= (-\infty, 1) \cup (1, \infty)$; horizontal asymptote is $y = 1$; vertical asymptote is $x = 2$

75. Here is one possibility.



76. Here is one possibility.



77. Here is one possibility.



78. Here is one possibility.



79. Here is one possibility.



80. Here is one possibility.



Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

81. Here is one possibility.



82. Here is one possibility.



83. Yes. If $\lim_{x\to\infty} \frac{f(x)}{g(x)} = 2$ then the ratio the polynomials' leading coefficients is 2, so $\lim_{x\to-\infty} \frac{f(x)}{g(x)} = 2$ as well.

84. Yes, it can have a horizontal or oblique asymptote.

85. At most 1 horizontal asymptote: If $\lim_{x\to\infty} \frac{f(x)}{g(x)} = L$, then the ratio of the polynomials' leading coefficients is L, so $\lim_{x\to-\infty} \frac{f(x)}{g(x)} = L$ as well.

86. $\lim_{x\to\infty} \left(\sqrt{x+9} - \sqrt{x+4}\right) = \lim_{x\to\infty} \left[\sqrt{x+9} - \sqrt{x+4}\right] \cdot \left[\frac{\sqrt{x+9}+\sqrt{x+4}}{\sqrt{x+9}+\sqrt{x+4}}\right] = \lim_{x\to\infty} \frac{(x+9)-(x+4)}{\sqrt{x+9}+\sqrt{x+4}}$

$= \lim_{x\to\infty} \frac{5}{\sqrt{x+9}+\sqrt{x+4}} = \lim_{x\to\infty} \frac{\frac{5}{\sqrt{x}}}{\sqrt{1+\frac{9}{x}}+\sqrt{1+\frac{4}{x}}} = \frac{0}{1+1} = 0$

87. $\lim_{x\to\infty} \left(\sqrt{x^2+25} - \sqrt{x^2-1}\right) = \lim_{x\to\infty} \left[\sqrt{x^2+25} - \sqrt{x^2-1}\right] \cdot \left[\frac{\sqrt{x^2+25}+\sqrt{x^2-1}}{\sqrt{x^2+25}+\sqrt{x^2-1}}\right] = \lim_{x\to\infty} \frac{(x^2+25)-(x^2-1)}{\sqrt{x^2+25}+\sqrt{x^2-1}}$

$= \lim_{x\to\infty} \frac{26}{\sqrt{x^2+25}+\sqrt{x^2-1}} = \lim_{x\to\infty} \frac{\frac{26}{x}}{\sqrt{1+\frac{25}{x^2}}+\sqrt{1-\frac{1}{x^2}}} = \frac{0}{1+1} = 0$

88. $\lim_{x\to-\infty} \left(\sqrt{x^2+3} + x\right) = \lim_{x\to-\infty} \left[\sqrt{x^2+3} + x\right] \cdot \left[\frac{\sqrt{x^2+3}-x}{\sqrt{x^2+3}-x}\right] = \lim_{x\to-\infty} \frac{(x^2+3)-(x^2)}{\sqrt{x^2+3}-x} = \lim_{x\to-\infty} \frac{3}{\sqrt{x^2+3}-x}$

$= \lim_{x\to-\infty} \frac{\frac{3}{\sqrt{x^2}}}{\sqrt{1+\frac{3}{x^2}}-\frac{x}{\sqrt{x^2}}} = \lim_{x\to-\infty} \frac{-\frac{3}{x}}{\sqrt{1+\frac{3}{x^2}}+1} = \frac{0}{1+1} = 0$

89. $\lim_{x\to-\infty} \left(2x + \sqrt{4x^2+3x-2}\right) = \lim_{x\to-\infty} \left[2x + \sqrt{4x^2+3x-2}\right] \cdot \left[\frac{2x-\sqrt{4x^2+3x-2}}{2x-\sqrt{4x^2+3x-2}}\right] = \lim_{x\to-\infty} \frac{(4x^2)-(4x^2+3x-2)}{2x-\sqrt{4x^2+3x-2}}$

$= \lim_{x\to-\infty} \frac{-3x+2}{2x-\sqrt{4x^2+3x-2}} = \lim_{x\to-\infty} \frac{\frac{-3x+2}{\sqrt{x^2}}}{\frac{2x}{\sqrt{x^2}}-\sqrt{4+\frac{3}{x}-\frac{2}{x^2}}} = \lim_{x\to-\infty} \frac{\frac{-3x+2}{-x}}{\frac{2x}{-x}-\sqrt{4+\frac{3}{x}-\frac{2}{x^2}}}$

$= \lim_{x\to-\infty} \frac{3-\frac{2}{x}}{-2-\sqrt{4+\frac{3}{x}-\frac{2}{x^2}}} = \frac{3-0}{-2-2} = -\frac{3}{4}$

90. $\lim_{x\to\infty} \left(\sqrt{9x^2-x} - 3x\right) = \lim_{x\to\infty} \left[\sqrt{9x^2-x} - 3x\right] \cdot \left[\frac{\sqrt{9x^2-x}+3x}{\sqrt{9x^2-x}+3x}\right] = \lim_{x\to\infty} \frac{(9x^2-x)-(9x^2)}{\sqrt{9x^2-x}+3x} = \lim_{x\to\infty} \frac{-x}{\sqrt{9x^2-x}+3x}$

$= \lim_{x\to\infty} \frac{-\frac{x}{x}}{\sqrt{\frac{9x^2}{x^2}-\frac{x}{x^2}}+\frac{3x}{x}} = \lim_{x\to\infty} \frac{-1}{\sqrt{9-\frac{1}{x}}+3} = \frac{-1}{3+3} = -\frac{1}{6}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

91. $\lim_{x\to\infty}\left(\sqrt{x^2+3x}-\sqrt{x^2-2x}\right)=\lim_{x\to\infty}\left[\sqrt{x^2+3x}-\sqrt{x^2-2x}\right]\cdot\left[\frac{\sqrt{x^2+3x}+\sqrt{x^2-2x}}{\sqrt{x^2+3x}+\sqrt{x^2-2x}}\right]=\lim_{x\to\infty}\frac{(x^2+3x)-(x^2-2x)}{\sqrt{x^2+3x}+\sqrt{x^2-2x}}$
$=\lim_{x\to\infty}\frac{5x}{\sqrt{x^2+3x}+\sqrt{x^2-2x}}=\lim_{x\to\infty}\frac{5}{\sqrt{1+\frac{3}{x}}+\sqrt{1-\frac{2}{x}}}=\frac{5}{1+1}=\frac{5}{2}$

92. $\lim_{x\to\infty}\sqrt{x^2+x}-\sqrt{x^2-x}=\lim_{x\to\infty}\left[\sqrt{x^2+x}-\sqrt{x^2-x}\right]\cdot\left[\frac{\sqrt{x^2+x}+\sqrt{x^2-x}}{\sqrt{x^2+x}+\sqrt{x^2-x}}\right]=\lim_{x\to\infty}\frac{(x^2+x)-(x^2-x)}{\sqrt{x^2+x}+\sqrt{x^2-x}}=\lim_{x\to\infty}\frac{2x}{\sqrt{x^2+x}+\sqrt{x^2-x}}$
$=\lim_{x\to\infty}\frac{2}{\sqrt{1+\frac{1}{x}}+\sqrt{1-\frac{1}{x}}}=\frac{2}{1+1}=1$

93. For any $\epsilon>0$, take $N=1$. Then for all $x>N$ we have that $|f(x)-k|=|k-k|=0<\epsilon$.

94. For any $\epsilon>0$, take $N=1$. Then for all $y<-N$ we have that $|f(x)-k|=|k-k|=0<\epsilon$.

95. For every real number $-B<0$, we must find a $\delta>0$ such that for all x, $0<|x-0|<\delta\Rightarrow\frac{-1}{x^2}<-B$.
Now, $-\frac{1}{x^2}<-B<0\Leftrightarrow\frac{1}{x^2}>B>0\Leftrightarrow x^2<\frac{1}{B}\Leftrightarrow|x|<\frac{1}{\sqrt{B}}$. Choose $\delta=\frac{1}{\sqrt{B}}$, then $0<|x|<\delta\Rightarrow|x|<\frac{1}{\sqrt{B}}$
$\Rightarrow\frac{-1}{x^2}<-B$ so that $\lim_{x\to0}-\frac{1}{x^2}=-\infty$.

96. For every real number $B>0$, we must find a $\delta>0$ such that for all x, $0<|x-0|<\delta\Rightarrow\frac{1}{|x|}>B$. Now,
$\frac{1}{|x|}>B>0\Leftrightarrow|x|<\frac{1}{B}$. Choose $\delta=\frac{1}{B}$. Then $0<|x-0|<\delta\Rightarrow|x|<\frac{1}{B}\Rightarrow\frac{1}{|x|}>B$ so that $\lim_{x\to0}\frac{1}{|x|}=\infty$.

97. For every real number $-B<0$, we must find a $\delta>0$ such that for all x, $0<|x-3|<\delta\Rightarrow\frac{-2}{(x-3)^2}<-B$. Now,
$\frac{-2}{(x-3)^2}<-B<0\Leftrightarrow\frac{2}{(x-3)^2}>B>0\Leftrightarrow\frac{(x-3)^2}{2}<\frac{1}{B}\Leftrightarrow(x-3)^2<\frac{2}{B}\Leftrightarrow0<|x-3|<\sqrt{\frac{2}{B}}$. Choose $\delta=\sqrt{\frac{2}{B}}$, then
$0<|x-3|<\delta\Rightarrow\frac{-2}{(x-3)^2}<-B<0$ so that $\lim_{x\to3}\frac{-2}{(x-3)^2}=-\infty$.

98. For every real number $B>0$, we must find a $\delta>0$ such that for all x, $0<|x-(-5)|<\delta\Rightarrow\frac{1}{(x+5)^2}>B$.
Now, $\frac{1}{(x+5)^2}>B>0\Leftrightarrow(x+5)^2<\frac{1}{B}\Leftrightarrow|x+5|<\frac{1}{\sqrt{B}}$. Choose $\delta=\frac{1}{\sqrt{B}}$. Then $0<|x-(-5)|<\delta$
$\Rightarrow|x+5|<\frac{1}{\sqrt{B}}\Rightarrow\frac{1}{(x+5)^2}>B$ so that $\lim_{x\to-5}\frac{1}{(x+5)^2}=\infty$.

99. (a) We say that $f(x)$ approaches infinity as x approaches x_0 from the left, and write $\lim_{x\to x_0^-}f(x)=\infty$, if for every positive number B, there exists a corresponding number $\delta>0$ such that for all x, $x_0-\delta<x<x_0\Rightarrow f(x)>B$.
(b) We say that $f(x)$ approaches minus infinity as x approaches x_0 from the right, and write $\lim_{x\to x_0^+}f(x)=-\infty$, if for every positive number B (or negative number $-B$) there exists a corresponding number $\delta>0$ such that for all x, $x_0<x<x_0+\delta\Rightarrow f(x)<-B$.
(c) We say that $f(x)$ approaches minus infinity as x approaches x_0 from the left, and write $\lim_{x\to x_0^-}f(x)=-\infty$, if for every positive number B (or negative number $-B$) there exists a corresponding number $\delta>0$ such that for all x, $x_0-\delta<x<x_0\Rightarrow f(x)<-B$.

100. For $B>0$, $\frac{1}{x}>B>0\Leftrightarrow x<\frac{1}{B}$. Choose $\delta=\frac{1}{B}$. Then $0<x<\delta\Rightarrow0<x<\frac{1}{B}\Rightarrow\frac{1}{x}>B$ so that $\lim_{x\to0^+}\frac{1}{x}=\infty$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

101. For $B > 0$, $\frac{1}{x} < -B < 0 \Leftrightarrow -\frac{1}{x} > B > 0 \Leftrightarrow -x < \frac{1}{B} \Leftrightarrow -\frac{1}{B} < x$. Choose $\delta = \frac{1}{B}$. Then $-\delta < x < 0 \Rightarrow -\frac{1}{B} < x$ $\Rightarrow \frac{1}{x} < -B$ so that $\lim_{x \to 0^-} \frac{1}{x} = -\infty$.

102. For $B > 0$, $\frac{1}{x-2} < -B \Leftrightarrow -\frac{1}{x-2} > B \Leftrightarrow -(x-2) < \frac{1}{B} \Leftrightarrow x - 2 > -\frac{1}{B} \Leftrightarrow x > 2 - \frac{1}{B}$. Choose $\delta = \frac{1}{B}$. Then $2 - \delta < x < 2 \Rightarrow -\delta < x - 2 < 0 \Rightarrow -\frac{1}{B} < x - 2 < 0 \Rightarrow \frac{1}{x-2} < -B < 0$ so that $\lim_{x \to 2^-} \frac{1}{x-2} = -\infty$.

103. For $B > 0$, $\frac{1}{x-2} > B \Leftrightarrow 0 < x - 2 < \frac{1}{B}$. Choose $\delta = \frac{1}{B}$. Then $2 < x < 2 + \delta \Rightarrow 0 < x - 2 < \delta \Rightarrow 0 < x - 2 < \frac{1}{B}$ $\Rightarrow \frac{1}{x-2} > B > 0$ so that $\lim_{x \to 2^+} \frac{1}{x-2} = \infty$.

104. For $B > 0$ and $0 < x < 1$, $\frac{1}{1-x^2} > B \Leftrightarrow 1 - x^2 < \frac{1}{B} \Leftrightarrow (1-x)(1+x) < \frac{1}{B}$. Now $\frac{1+x}{2} < 1$ since $x < 1$. Choose $\delta < \frac{1}{2B}$. Then $1 - \delta < x < 1 \Rightarrow -\delta < x - 1 < 0 \Rightarrow 1 - x < \delta < \frac{1}{2B} \Rightarrow (1-x)(1+x) < \frac{1}{B}\left(\frac{1+x}{2}\right) < \frac{1}{B} \Rightarrow \frac{1}{1-x^2} > B$ for $0 < x < 1$ and x near $1 \Rightarrow \lim_{x \to 1^-} \frac{1}{1-x^2} = \infty$.

105. $y = \frac{x^2}{x-1} = x + 1 + \frac{1}{x-1}$



106. $y = \frac{x^2+1}{x-1} = x + 1 + \frac{2}{x-1}$



107. $y = \frac{x^2-4}{x-1} = x + 1 - \frac{3}{x-1}$



108. $y = \frac{x^2-1}{2x+4} = \frac{1}{2}x - 1 + \frac{3}{2x+4}$



109. $y = \frac{x^2-1}{x} = x - \frac{1}{x}$



110. $y = \frac{x^3+1}{x^2} = x + \frac{1}{x^2}$



Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

111. $y = \dfrac{x}{\sqrt{4-x^2}}$



112. $y = \dfrac{-1}{\sqrt{4-x^2}}$



113. $y = x^{2/3} + \dfrac{1}{x^{1/3}}$



114. $y = \sin\left(\dfrac{\pi}{x^2+1}\right)$



115. (a) $y \to \infty$ (see accompanying graph)
(b) $y \to \infty$ (see accompanying graph)
(c) cusps at $x = \pm 1$ (see accompanying graph)



116. (a) $y \to 0$ and a cusp at $x = 0$ (see the accompanying graph)
(b) $y \to \frac{3}{2}$ (see accompanying graph)
(c) a vertical asymptote at $x = 1$ and contains the point $\left(-1, \dfrac{3}{2\sqrt[3]{4}}\right)$ (see accompanying graph)



Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

CHAPTER 2 PRACTICE EXERCISES

1. At $x=-1$: $\lim_{x\to -1^-} f(x) = \lim_{x\to -1^+} f(x) = 1$
$\Rightarrow \lim_{x\to -1} f(x) = 1 = f(-1)$
$\Rightarrow f$ is continuous at $x=-1$.



At $x=0$: $\lim_{x\to 0^-} f(x) = \lim_{x\to 0^+} f(x) = 0$
$\Rightarrow \lim_{x\to 0} f(x) = 0.$
But $f(0) = 1 \neq \lim_{x\to 0} f(x)$
$\Rightarrow f$ is discontinuous at $x=0$.
If we define $f(0)=0$, then the discontinuity at $x=0$ is removable.

At $x=1$: $\lim_{x\to 1^-} f(x) = -1$ and $\lim_{x\to 1^+} f(x) = 1$
$\Rightarrow \lim_{x\to 1} f(x)$ does not exist
$\Rightarrow f$ is discontinuous at $x=1$.

2. At $x=-1$: $\lim_{x\to -1^-} f(x) = 0$ and $\lim_{x\to -1^+} f(x) = -1$
$\Rightarrow \lim_{x\to -1} f(x)$ does not exist
$\Rightarrow f$ is discontinuous at $x=-1$.



At $x=0$: $\lim_{x\to 0^-} f(x) = -\infty$ and $\lim_{x\to 0^+} f(x) = \infty$
$\Rightarrow \lim_{x\to 0} f(x)$ does not exist
$\Rightarrow f$ is discontinuous at $x=0$.

At $x=1$: $\lim_{x\to 1^-} f(x) = \lim_{x\to 1^+} f(x) = 1 \Rightarrow \lim_{x\to 1} f(x) = 1.$
But $f(1) = 0 \neq \lim_{x\to 1} f(x)$
$\Rightarrow f$ is discontinuous at $x=1$.
If we define $f(1)=1$, then the discontinuity at $x=1$ is removable.

3. (a) $\lim_{t\to t_0} (3f(t)) = 3 \lim_{t\to t_0} f(t) = 3(-7) = -21$

(b) $\lim_{t\to t_0} (f(t))^2 = \left(\lim_{t\to t_0} f(t)\right)^2 = (-7)^2 = 49$

(c) $\lim_{t\to t_0} (f(t)\cdot g(t)) = \lim_{t\to t_0} f(t) \cdot \lim_{t\to t_0} g(t) = (-7)(0) = 0$

(d) $\lim_{t\to t_0} \dfrac{f(t)}{g(t)-7} = \dfrac{\lim_{t\to t_0} f(t)}{\lim_{t\to t_0} (g(t)-7)} = \dfrac{\lim_{t\to t_0} f(t)}{\lim_{t\to t_0} g(t) - \lim_{t\to t_0} 7} = \dfrac{-7}{0-7} = 1$

(e) $\lim_{t\to t_0} \cos(g(t)) = \cos\left(\lim_{t\to t_0} g(t)\right) = \cos 0 = 1$

(f) $\lim_{t\to t_0} |f(t)| = \left|\lim_{t\to t_0} f(t)\right| = |-7| = 7$

(g) $\lim_{t\to t_0} (f(t)+g(t)) = \lim_{t\to t_0} f(t) + \lim_{t\to t_0} g(t) = -7 + 0 = -7$

(h) $\lim_{t\to t_0} \left(\dfrac{1}{f(t)}\right) = \dfrac{1}{\lim_{t\to t_0} f(t)} = \dfrac{1}{-7} = -\dfrac{1}{7}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

4. (a) $\lim_{x\to 0} -g(x) = -\lim_{x\to 0} g(x) = -\sqrt{2}$

(b) $\lim_{x\to 0} (g(x)\cdot f(x)) = \lim_{x\to 0} g(x)\cdot \lim_{x\to 0} f(x) = \left(\sqrt{2}\right)\left(\frac{1}{2}\right) = \frac{\sqrt{2}}{2}$

(c) $\lim_{x\to 0} (f(x)+g(x)) = \lim_{x\to 0} f(x) + \lim_{x\to 0} g(x) = \frac{1}{2} + \sqrt{2}$

(d) $\lim_{x\to 0} \frac{1}{f(x)} = \frac{1}{\lim_{x\to 0} f(x)} = \frac{1}{\frac{1}{2}} = 2$

(e) $\lim_{x\to 0} (x + f(x)) = \lim_{x\to 0} x + \lim_{x\to 0} f(x) = 0 + \frac{1}{2} = \frac{1}{2}$

(f) $\lim_{x\to 0} \frac{f(x)\cdot\cos x}{x-1} = \frac{\lim_{x\to 0} f(x)\cdot \lim_{x\to 0} \cos x}{\lim_{x\to 0} x - \lim_{x\to 0} 1} = \frac{\left(\frac{1}{2}\right)(1)}{0-1} = -\frac{1}{2}$

5. Since $\lim_{x\to 0} x = 0$ we must have that $\lim_{x\to 0} (4 - g(x)) = 0$. Otherwise, if $\lim_{x\to 0} (4 - g(x))$ is a finite positive number, we would have $\lim_{x\to 0^-} \left[\frac{4-g(x)}{x}\right] = -\infty$ and $\lim_{x\to 0^+} \left[\frac{4-g(x)}{x}\right] = \infty$ so the limit could not equal 1 as $x \to 0$. Similar reasoning holds if $\lim_{x\to 0} (4 - g(x))$ is a finite negative number. We conclude that $\lim_{x\to 0} g(x) = 4$.

6. $2 = \lim_{x\to -4}\left[x \lim_{x\to 0} g(x)\right] = \lim_{x\to -4} x \cdot \lim_{x\to -4}\left[\lim_{x\to 0} g(x)\right] = -4 \lim_{x\to -4}\left[\lim_{x\to 0} g(x)\right] = -4 \lim_{x\to 0} g(x)$ (since $\lim_{x\to 0} g(x)$ is a constant) $\Rightarrow \lim_{x\to 0} g(x) = \frac{2}{-4} = -\frac{1}{2}$.

7. (a) $\lim_{x\to c} f(x) = \lim_{x\to c} x^{1/3} = c^{1/3} = f(c)$ for every real number $c \Rightarrow f$ is continuous on $(-\infty, \infty)$.

(b) $\lim_{x\to c} g(x) = \lim_{x\to c} x^{3/4} = c^{3/4} = g(c)$ for every nonnegative real number $c \Rightarrow g$ is continuous on $[0, \infty)$.

(c) $\lim_{x\to c} h(x) = \lim_{x\to c} x^{-2/3} = \frac{1}{c^{2/3}} = h(c)$ for every nonzero real number $c \Rightarrow h$ is continuous on $(-\infty, 0)$ and $(-\infty, \infty)$.

(d) $\lim_{x\to c} k(x) = \lim_{x\to c} x^{-1/6} = \frac{1}{c^{1/6}} = k(c)$ for every positive real number $c \Rightarrow k$ is continuous on $(0, \infty)$

8. (a) $\bigcup_{n\in I} \left(\left(n - \frac{1}{2}\right)\pi, \left(n + \frac{1}{2}\right)\pi\right)$, where I = the set of all integers.

(b) $\bigcup_{n\in I} (n\pi,\ (n+1)\pi)$, where I = the set of all integers.

(c) $(-\infty,\ \pi) \cup (\pi,\ \infty)$

(d) $(-\infty,\ 0) \cup (0,\ \infty)$

9. (a) $\lim_{x\to 0} \frac{x^2-4x+4}{x^3+5x^2-14x} = \lim_{x\to 0} \frac{(x-2)(x-2)}{x(x+7)(x-2)} = \lim_{x\to 0} \frac{x-2}{x(x+7)}$, $x \neq 2$; the limit does not exist because $\lim_{x\to 0^-} \frac{x-2}{x(x+7)} = \infty$ and $\lim_{x\to 0^+} \frac{x-2}{x(x+7)} = -\infty$

(b) $\lim_{x\to 2} \frac{x^2-4x+4}{x^3+5x^2-14x} = \lim_{x\to 2} \frac{(x-2)(x-2)}{x(x+7)(x-2)} = \lim_{x\to 2} \frac{x-2}{x(x+7)}$, $x \neq 2$, and $\lim_{x\to 2} \frac{x-2}{x(x+7)} = \frac{0}{2(9)} = 0$

10. (a) $\lim_{x\to 0} \frac{x^2+x}{x^5+2x^4+x^3} = \lim_{x\to 0} \frac{x(x+1)}{x^3(x^2+2x+1)} = \lim_{x\to 0} \frac{x+1}{x^2(x+1)(x+1)} = \lim_{x\to 0} \frac{1}{x^2(x+1)}$, $x \neq 0$ and $x \neq -1$.
Now $\lim_{x\to 0^-} \frac{1}{x^2(x+1)} = \infty$ and $\lim_{x\to 0^+} \frac{1}{x^2(x+1)} = \infty \Rightarrow \lim_{x\to 0} \frac{x^2+x}{x^5+2x^4+x^3} = \infty$.

(b) $\lim_{x\to -1} \frac{x^2+x}{x^5+2x^4+x^3} = \lim_{x\to -1} \frac{x(x+1)}{x^3(x^2+2x+1)} = \lim_{x\to -1} \frac{1}{x^2(x+1)}$, $x \neq 0$ and $x \neq -1$. The limit does not exist because $\lim_{x\to -1^-} \frac{1}{x^2(x+1)} = -\infty$ and $\lim_{x\to -1^+} \frac{1}{x^2(x+1)} = \infty$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

11. $\lim\limits_{x\to 1}\dfrac{1-\sqrt{x}}{1-x}=\lim\limits_{x\to 1}\dfrac{1-\sqrt{x}}{(1-\sqrt{x})(1+\sqrt{x})}=\lim\limits_{x\to 1}\dfrac{1}{1+\sqrt{x}}=\dfrac{1}{2}$

12. $\lim\limits_{x\to a}\dfrac{x^2-a^2}{x^4-a^4}=\lim\limits_{x\to a}\dfrac{(x^2-a^2)}{(x^2+a^2)(x^2-a^2)}=\lim\limits_{x\to a}\dfrac{1}{x^2+a^2}=\dfrac{1}{2a^2}$

13. $\lim\limits_{h\to 0}\dfrac{(x+h)^2-x^2}{h}=\lim\limits_{h\to 0}\dfrac{(x^2+2hx+h^2)-x^2}{h}=\lim\limits_{h\to 0}\ (2x+h)=2x$

14. $\lim\limits_{x\to 0}\dfrac{(x+h)^2-x^2}{h}=\lim\limits_{x\to 0}\dfrac{(x^2+2hx+h^2)-x^2}{h}=\lim\limits_{x\to 0}\ (2x+h)=h$

15. $\lim\limits_{x\to 0}\dfrac{\frac{1}{2+x}-\frac{1}{2}}{x}=\lim\limits_{x\to 0}\dfrac{2-(2+x)}{2x(2+x)}=\lim\limits_{x\to 0}\ \dfrac{-1}{4+2x}=-\dfrac{1}{4}$

16. $\lim\limits_{x\to 0}\dfrac{(2+x)^3-8}{x}=\lim\limits_{x\to 0}\dfrac{(x^3+6x^2+12x+8)-8}{x}=\lim\limits_{x\to 0}\ (x^2+6x+12)=12$

17. $\lim\limits_{x\to 1}\dfrac{x^{1/3}-1}{\sqrt{x}-1}=\lim\limits_{x\to 1}\dfrac{(x^{1/3}-1)}{(\sqrt{x}-1)}\cdot\dfrac{(x^{2/3}+x^{1/3}+1)(\sqrt{x}+1)}{(\sqrt{x}+1)(x^{2/3}+x^{1/3}+1)}=\lim\limits_{x\to 1}\dfrac{(x-1)(\sqrt{x}+1)}{(x-1)(x^{2/3}+x^{1/3}+1)}=\lim\limits_{x\to 1}\dfrac{\sqrt{x}+1}{x^{2/3}+x^{1/3}+1}=\dfrac{1+1}{1+1+1}=\dfrac{2}{3}$

18. $\lim\limits_{x\to 64}\dfrac{x^{2/3}-16}{\sqrt{x}-8}=\lim\limits_{x\to 64}\dfrac{(x^{1/3}-4)(x^{1/3}+4)}{\sqrt{x}-8}=\lim\limits_{x\to 64}\dfrac{(x^{1/3}-4)(x^{1/3}+4)}{\sqrt{x}-8}\cdot\dfrac{(x^{2/3}+4x^{1/3}+16)(\sqrt{x}+8)}{(\sqrt{x}+8)(x^{2/3}+4x^{1/3}+16)}$
$=\lim\limits_{x\to 64}\dfrac{(x-64)\,(x^{1/3}+4)\,(\sqrt{x}+8)}{(x-64)\,(x^{2/3}+4x^{1/3}+16)}=\lim\limits_{x\to 64}\dfrac{(x^{1/3}+4)\,(\sqrt{x}+8)}{x^{2/3}+4x^{1/3}+16}=\dfrac{(4+4)\,(8+8)}{16+16+16}=\dfrac{8}{3}$

19. $\lim\limits_{x\to 0}\dfrac{\tan 2x}{\tan \pi x}=\lim\limits_{x\to 0}\dfrac{\sin\ 2x}{\cos\ 2x}\cdot\dfrac{\cos \pi x}{\sin \pi x}=\lim\limits_{x\to 0}\left(\dfrac{\sin\ 2x}{2x}\right)\left(\dfrac{\cos \pi x}{\cos\ 2x}\right)\left(\dfrac{\pi x}{\sin \pi x}\right)\left(\dfrac{2x}{\pi x}\right)=1\cdot 1\cdot 1\cdot\dfrac{2}{\pi}=\dfrac{2}{\pi}$

20. $\lim\limits_{x\to \pi^-}\csc x=\lim\limits_{x\to \pi^-}\dfrac{1}{\sin x}=\infty$

21. $\lim\limits_{x\to \pi}\sin\left(\dfrac{x}{2}+\sin x\right)=\sin\left(\dfrac{\pi}{2}+\sin \pi\right)=\sin\left(\dfrac{\pi}{2}\right)=1$

22. $\lim\limits_{x\to \pi}\cos^2(x-\tan x)=\cos^2(\pi-\tan \pi)\ =\ \cos^2(\pi)=(-1)^2=1$

23. $\lim\limits_{x\to 0}\dfrac{8x}{3\sin\ x-x}=\lim\limits_{x\to 0}\dfrac{8}{3\frac{\sin x}{x}-1}=\dfrac{8}{3(1)-1}=4$

24. $\lim\limits_{x\to 0}=\dfrac{\cos\ 2x-1}{\sin x}=\lim\limits_{x\to 0}\left(\dfrac{\cos\ 2x-1}{\sin x}\cdot\dfrac{\cos\ 2x+1}{\cos\ 2x+1}\right)=\lim\limits_{x\to 0}\dfrac{\cos^2\ 2x-1}{\sin x(\cos\ 2x+1)}=\lim\limits_{x\to 0}\dfrac{-\sin^2 2x}{\sin\ x(\cos\ 2x+1)}$
$=\lim\limits_{x\to 0}\dfrac{-4\sin x\cos^2 x}{\cos\ 2x+1}=\dfrac{-4(0)(1)^2}{1+1}=0$

25. Let $x=t-3\Rightarrow\lim\limits_{t\to 3^+}\ln(t-3)=\lim\limits_{x\to 0^+}\ln x=-\infty$

26. $\lim\limits_{t\to 1}t^2\ln\left(2-\sqrt{t}\right)=\ln 1=0$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

27. $-1 \le \cos\left(\frac{\pi}{\theta}\right) \le 1 \Rightarrow e^{-1} \le e^{\cos(\pi/\theta)} \le e^{1} \Rightarrow \sqrt{\theta}e^{-1} \le \sqrt{\theta}e^{\cos(\pi/\theta)} \le \sqrt{\theta}e \Rightarrow \lim_{\theta \to 0^+} \sqrt{\theta}e^{\cos(\pi/\theta)} = 0$ by the Sandwich Theorem

28. $\lim_{z \to 0^+} \frac{2e^{1/z}}{e^{1/z}+1} = \lim_{z \to 0^+} \frac{2}{1+e^{1/z}} = \frac{2}{1+0} = 2$

29. $\lim_{x \to 0^+} [4\,g(x)]^{1/3} = 2 \Rightarrow \left[\lim_{x \to 0^+} 4\,g(x)\right]^{1/3} = 2 \Rightarrow \lim_{x \to 0^+} 4\,g(x) = 8$, since $2^3 = 8$. Then $\lim_{x \to 0^+} g(x) = 2$.

30. $\lim_{x \to \sqrt{5}} \frac{1}{x+g(x)} = 2 \Rightarrow \lim_{x \to \sqrt{5}} (x + g(x)) = \frac{1}{2} \Rightarrow \sqrt{5} + \lim_{x \to \sqrt{5}} g(x) = \frac{1}{2} \Rightarrow \lim_{x \to \sqrt{5}} g(x) = \frac{1}{2} - \sqrt{5}$

31. $\lim_{x \to 1} \frac{3x^2+1}{g(x)} = \infty \Rightarrow \lim_{x \to 1} g(x) = 0$ since $\lim_{x \to 1} (3x^2 + 1) = 4$

32. $\lim_{x \to -2} \frac{5-x^2}{\sqrt{g(x)}} = 0 \Rightarrow \lim_{x \to -2} g(x) = \infty$ since $\lim_{x \to -2} (5 - x^2) = 1$

33. (a) $f(-1) = -1$ and $f(2) = 5 \Rightarrow f$ has a root between -1 and 2 by the Intermediate Value Theorem.
(b), (c) root is 1.32471795724

34. (a) $f(-2) = -2$ and $f(0) = 2 \Rightarrow f$ has a root between -2 and 0 by the Intermediate Value Theorem.
(b), (c) root is -1.76929235424

35. At $x = -1$: $\lim_{x \to -1^-} f(x) = \lim_{x \to -1^-} \frac{x(x^2-1)}{|x^2-1|}$
$= \lim_{x \to -1^-} \frac{x(x^2-1)}{x^2-1} = \lim_{x \to -1^-} x = -1$, and
$\lim_{x \to -1^+} f(x) = \lim_{x \to -1^+} \frac{x(x^2-1)}{|x^2-1|} = \lim_{x \to -1^+} \frac{x(x^2-1)}{-(x^2-1)}$
$= \lim_{x \to -1} (-x) = -(-1) = 1$. Since $\lim_{x \to -1^-} f(x) \ne \lim_{x \to -1^+} f(x) \Rightarrow \lim_{x \to -1} f(x)$ does not exist, the function f cannot be extended to a continuous function at $x = -1$.



At $x = 1$: $\lim_{x \to 1^-} f(x) = \lim_{x \to 1^-} \frac{x(x^2-1)}{|x^2-1|} = \lim_{x \to 1^-} \frac{x(x^2-1)}{-(x^2-1)} = \lim_{x \to 1^-} (-x) = -1$, and $\lim_{x \to 1^+} f(x) = \lim_{x \to 1^+} \frac{x(x^2-1)}{|x^2-1|}$
$= \lim_{x \to 1^+} \frac{x(x^2-1)}{x^2-1} = \lim_{x \to 1^+} x = 1$.
Again $\lim_{x \to 1} f(x)$ does not exist so f cannot be extended to a continuous function at $x = 1$ either.

36. The discontinuity at $x = 0$ of $f(x) = \sin\left(\frac{1}{x}\right)$ is nonremovable because $\lim_{x \to 0} \sin\frac{1}{x}$ does not exist.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

37. Yes, f does have a continuous extension at $a = 1$: define $f(1) = \lim_{x\to 1} \frac{x-1}{x-\sqrt[4]{x}} = \frac{4}{3}$.



$f(x) = \dfrac{x-1}{x-\sqrt[4]{x}}, \quad a = 1$

38. Yes, g does have a continuous extension at $a = \frac{\pi}{2}$: $g\left(\frac{\pi}{2}\right) = \lim_{\theta\to\frac{\pi}{2}} \frac{5\cos\theta}{4\theta - 2\pi} = -\frac{5}{4}$.



39. From the graph we see that $\lim_{t\to 0^-} h(t) \neq \lim_{t\to 0^+} h(t)$ so h cannot be extended to a continuous function at $a = 0$.



$h(t) = (1 + |t|)^{1/t}, \quad a = 0$

40. From the graph we see that $\lim_{x\to 0^-} k(x) \neq \lim_{x\to 0^+} k(x)$ so k cannot be extended to a continuous function at $a = 0$.



41. $\lim_{x\to\infty} \frac{2x+3}{5x+7} = \lim_{x\to\infty} \frac{2+\frac{3}{x}}{5+\frac{7}{x}} = \frac{2+0}{5+0} = \frac{2}{5}$

42. $\lim_{x\to-\infty} \frac{2x^2+3}{5x^2+7} = \lim_{x\to-\infty} \frac{2+\frac{3}{x^2}}{5+\frac{7}{x^2}} = \frac{2+0}{5+0} = \frac{2}{5}$

43. $\lim_{x\to-\infty} \frac{x^2-4x+8}{3x^3} = \lim_{x\to-\infty} \left(\frac{1}{3x} - \frac{4}{3x^2} + \frac{8}{3x^3}\right) = 0 - 0 + 0 = 0$

44. $\lim_{x\to\infty} \frac{1}{x^2-7x+1} = \lim_{x\to\infty} \frac{\frac{1}{x^2}}{1-\frac{7}{x}+\frac{1}{x^2}} = \frac{0}{1-0+0} = 0$

45. $\lim_{x\to-\infty} \frac{x^2-7x}{x+1} = \lim_{x\to-\infty} \frac{x-7}{1+\frac{1}{x}} = -\infty$

46. $\lim_{x\to\infty} \frac{x^4+x^3}{12x^3+128} = \lim_{x\to-\infty} \frac{x+1}{12+\frac{128}{x^3}} = \infty$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

47. $\lim_{x\to\infty}\frac{\sin x}{\lfloor x\rfloor}\le\lim_{x\to\infty}\frac{1}{\lfloor x\rfloor}=0$ since $\lfloor x\rfloor\to\infty$ as $x\to\infty\Rightarrow\lim_{x\to\infty}\frac{\sin x}{\lfloor x\rfloor}=0.$

48. $\lim_{\theta\to\infty}\frac{\cos\theta-1}{\theta}\le\lim_{\theta\to\infty}\frac{2}{\theta}=0\Rightarrow\lim_{\theta\to\infty}\frac{\cos\theta-1}{\theta}=0.$

49. $\lim_{x\to\infty}\frac{x+\sin x+2\sqrt{x}}{x+\sin x}=\lim_{x\to\infty}\frac{1+\frac{\sin x}{x}+\frac{2}{\sqrt{x}}}{1+\frac{\sin x}{x}}=\frac{1+0+0}{1+0}=1$

50. $\lim_{x\to\infty}\frac{x^{2/3}+x^{-1}}{x^{2/3}+\cos^2 x}=\lim_{x\to\infty}\left(\frac{1+x^{-5/3}}{1+\frac{\cos^2 x}{x^{2/3}}}\right)=\frac{1+0}{1+0}=1$

51. $\lim_{x\to\infty}e^{1/x}\cos\left(\frac{1}{x}\right)=e^0\cdot\cos(0)=1\cdot1=1$

52. $\lim_{x\to\infty}\ln\left(1+\frac{1}{t}\right)=\ln 1=0$

53. $\lim_{x\to-\infty}\tan^{-1}x=-\frac{\pi}{2}$

54. $\lim_{t\to-\infty}e^{3t}\sin^{-1}\left(\frac{1}{t}\right)=0\cdot\sin^{-1}(0)=0\cdot0=0$

55. (a) $y=\frac{x^2+4}{x-3}$ is undefined at $x=3$: $\lim_{x\to3^-}\frac{x^2+4}{x-3}=-\infty$ and $\lim_{x\to3^+}\frac{x^2+4}{x-3}=+\infty,$ thus $x=3$ is a vertical asymptote.

(b) $y=\frac{x^2-x-2}{x^2-2x+1}$ is undefined at $x=1$: $\lim_{x\to1^-}\frac{x^2-x-2}{x^2-2x+1}=-\infty$ and $\lim_{x\to1^+}\frac{x^2-x-2}{x^2-2x+1}=-\infty,$ thus $x=1$ is a vertical asymptote.

(c) $y=\frac{x^2+x-6}{x^2+2x-8}$ is undefined at $x=2$ and -4: $\lim_{x\to2}\frac{x^2+x-6}{x^2+2x-8}=\lim_{x\to2}\frac{x+3}{x+4}=\frac{5}{6};$ $\lim_{x\to-4^-}\frac{x^2+x-6}{x^2+2x-8}=\lim_{x\to-4^-}\frac{x+3}{x+4}=\infty$ $\lim_{x\to-4^+}\frac{x^2+x-6}{x^2+2x-8}=\lim_{x\to-4^+}\frac{x+3}{x+4}=-\infty.$ Thus $x=-4$ is a vertical asymptote.

56. (a) $y=\frac{1-x^2}{x^2+1}$: $\lim_{x\to\infty}\frac{1-x^2}{x^2+1}=\lim_{x\to\infty}\frac{\frac{1}{x^2}-1}{1+\frac{1}{x^2}}=\frac{-1}{1}=-1$ and $\lim_{x\to-\infty}\frac{1-x^2}{x^2+1}=\lim_{x\to-\infty}\frac{\frac{1}{x^2}-1}{1+\frac{1}{x^2}}=\frac{-1}{1}=-1,$ thus $y=-1$ is a horizontal asymptote.

(b) $y=\frac{\sqrt{x}+4}{\sqrt{x}+4}$: $\lim_{x\to\infty}\frac{\sqrt{x}+4}{\sqrt{x}+4}=\lim_{x\to\infty}\frac{1+\frac{4}{\sqrt{x}}}{\sqrt{1+\frac{4}{x}}}=\frac{1+0}{\sqrt{1+0}}=1,$ thus $y=1$ is a horizontal asymptote.

(c) $y=\frac{\sqrt{x^2+4}}{x}$: $\lim_{x\to\infty}\frac{\sqrt{x^2+4}}{x}=\lim_{x\to\infty}\frac{\sqrt{1+\frac{4}{x^2}}}{1}=\frac{\sqrt{1+0}}{1}=1$ and $\lim_{x\to-\infty}\frac{\sqrt{x^2+4}}{x}=\lim_{x\to-\infty}\frac{\sqrt{1+\frac{4}{x^2}}}{\frac{x}{\sqrt{x^2}}}$

$=\lim_{x\to-\infty}\frac{\sqrt{1+\frac{4}{x^2}}}{\frac{x}{-x}}=\lim_{x\to-\infty}\frac{\sqrt{1+\frac{4}{x^2}}}{-1}=\frac{\sqrt{1+0}}{-1}=\frac{1}{-1}=-1,$

thus $y=1$ and $y=-1$ are horizontal asymptotes.

(d) $y=\sqrt{\frac{x^2+9}{9x^2+1}}$: $\lim_{x\to\infty}\sqrt{\frac{x^2+9}{9x^2+1}}=\lim_{x\to\infty}\sqrt{\frac{1+\frac{9}{x^2}}{9+\frac{1}{x^2}}}=\sqrt{\frac{1+0}{9+0}}=\frac{1}{3}$ and $\lim_{x\to-\infty}\sqrt{\frac{x^2+9}{9x^2+1}}=\lim_{x\to-\infty}\sqrt{\frac{1+\frac{9}{x^2}}{9+\frac{1}{x^2}}}=\sqrt{\frac{1+0}{9+0}}=\frac{1}{3},$ thus $y=\frac{1}{3}$ is a horizontal asymptote.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

57. domain $= [-4, 2) \cup (2, 4]$; y in range and $y = \dfrac{\sqrt{16-x^2}}{x-2}$, if $x = \pm 4$, then $y = 0$, $\lim\limits_{x \to 2^+} \dfrac{\sqrt{16-x^2}}{x-2} = \infty$, and $\lim\limits_{x \to 2^-} \dfrac{\sqrt{16-x^2}}{x-2} = -\infty$, $\Rightarrow$ range $= (-\infty, \infty)$

58. Since $\lim\limits_{x \to b^{\pm}} \dfrac{\sqrt{ax^2+4}}{x-b} = \pm\infty \Rightarrow$ vertical asymptote is $x = b$; $\lim\limits_{x \to \infty} \dfrac{\sqrt{ax^2+4}}{x-b} = \lim\limits_{x \to \infty} \dfrac{|x|}{x-b} \cdot \sqrt{a + \dfrac{4}{x^2}}$

$= \lim\limits_{x \to \infty} \dfrac{x}{x-b} \sqrt{a + \dfrac{4}{x^2}} = \sqrt{a} \Rightarrow$ horizontal asymptote is $y = \sqrt{a}$, $\lim\limits_{x \to -\infty} \dfrac{\sqrt{ax^2+4}}{x-b}$

$= \lim\limits_{x \to -\infty} \dfrac{|x|}{x-b} \cdot \sqrt{a + \dfrac{4}{x^2}} = \lim\limits_{x \to -\infty} \dfrac{-x}{x-b} \cdot \sqrt{a + \dfrac{4}{x^2}} = -\sqrt{a} \Rightarrow$ horizontal asymptote is $y = -\sqrt{a}$

CHAPTER 2 ADDITIONAL AND ADVANCED EXERCISES

1. (a)

x	0.1	0.01	0.001	0.0001	0.00001
x^x	0.7943	0.9550	0.9931	0.9991	0.9999

Apparently, $\lim\limits_{x \to 0^+} x^x = 1$

(b)



2. (a)

x	10	100	1000
$\left(\frac{1}{x}\right)^{1/(\ln x)}$	0.3679	0.3679	0.3679

Apparently, $\lim\limits_{x \to \infty} \left(\dfrac{1}{x}\right)^{1/(\ln x)} = 0.3678 = \dfrac{1}{e}$

(b)



Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

3. $\lim_{v\to c^-} L = \lim_{v\to c^-} L_0\sqrt{1-\frac{v^2}{c^2}} = L_0\sqrt{1-\frac{\lim_{v\to c^-} v^2}{c^2}} = L_0\sqrt{1-\frac{c^2}{c^2}} = 0$

The left-hand limit was needed because the function L is undefined if $v > c$ (the rocket cannot move faster than the speed of light).

4. (a) $\left|\frac{\sqrt{x}}{2}-1\right| < 0.2 \Rightarrow -0.2 < \frac{\sqrt{x}}{2}-1 < 0.2 \Rightarrow 0.8 < \frac{\sqrt{x}}{2} < 1.2 \Rightarrow 1.6 < \sqrt{x} < 2.4 \Rightarrow 2.56 < x < 5.76.$

(b) $\left|\frac{\sqrt{x}}{2}-1\right| < 0.1 \Rightarrow -0.1 < \frac{\sqrt{x}}{2}-1 < 0.1 \Rightarrow 0.9 < \frac{\sqrt{x}}{2} < 1.1 \Rightarrow 1.8 < \sqrt{x} < 2.2 \Rightarrow 3.24 < x < 4.84.$

5. $|10+(t-70)\times 10^{-4}-10| < 0.0005 \Rightarrow |(t-70)\times 10^{-4}| < 0.0005 \Rightarrow -0.0005 < (t-70)\times 10^{-4} < 0.0005$
$\Rightarrow -5 < t-70 < 5 \Rightarrow 65° < t < 75° \Rightarrow$ Within 5° F.

6. We want to know in what interval to hold values of h to make V satisfy the inequality $|V-1000| = |36\pi h - 1000| \le 10$. To find out, we solve the inequality:
$|36\pi h - 1000| \le 10 \Rightarrow -10 \le 36\pi h - 1000 \le 10 \Rightarrow 990 \le 36\pi h \le 1010 \Rightarrow \frac{990}{36\pi} \le h \le \frac{1010}{36\pi} \Rightarrow 8.8 \le h \le 8.9$
where 8.8 was rounded up, to be safe, and 8.9 was rounded down, to be safe.
The interval in which we should hold h is about $8.9-8.8 = 0.1$ cm wide (1 mm). With stripes 1 mm wide, we can expect to measure a liter of water with an accuracy of 1%, which is more than enough accuracy for cooking.

7. Show $\lim_{x\to 1} f(x) = \lim_{x\to 1} (x^2-7) = -6 = f(1)$.
Step 1: $|(x^2-7)+6| < \epsilon \Rightarrow -\epsilon < x^2-1 < \epsilon \Rightarrow 1-\epsilon < x^2 < 1+\epsilon \Rightarrow \sqrt{1-\epsilon} < x < \sqrt{1+\epsilon}$.
Step 2: $|x-1| < \delta \Rightarrow -\delta < x-1 < \delta \Rightarrow -\delta+1 < x < \delta+1$.
Then $-\delta+1 = \sqrt{1-\epsilon}$ or $\delta+1 = \sqrt{1+\epsilon}$. Choose $\delta = \min\left\{1-\sqrt{1-\epsilon}, \sqrt{1+\epsilon}-1\right\}$, then $0 < |x-1| < \delta \Rightarrow$ $|(x^2-7)-6| < \epsilon$ and $\lim_{x\to -1} f(x) = -6$. By the continuity text, $f(x)$ is continuous at $x = 1$.

8. Show $\lim_{x\to \frac{1}{4}} g(x) = \lim_{x\to \frac{1}{4}} \frac{1}{2x} = 2 = g\left(\frac{1}{4}\right)$.
Step 1: $\left|\frac{1}{2x}-2\right| < \epsilon \Rightarrow -\epsilon < \frac{1}{2x}-2 < \epsilon \Rightarrow 2-\epsilon < \frac{1}{2x} < 2+\epsilon \Rightarrow \frac{1}{4-2\epsilon} > x > \frac{1}{4+2\epsilon}$.
Step 2: $\left|x-\frac{1}{4}\right| < \delta \Rightarrow -\delta < x-\frac{1}{4} < \delta \Rightarrow -\delta+\frac{1}{4} < x < \delta+\frac{1}{4}$.
Then $-\delta+\frac{1}{4} = \frac{1}{4+2\epsilon} \Rightarrow \delta = \frac{1}{4}-\frac{1}{4+2\epsilon} = \frac{\epsilon}{4(2-\epsilon)}$, or $\delta+\frac{1}{4} = \frac{1}{4-2\epsilon} \Rightarrow \delta = \frac{1}{4-2\epsilon}-\frac{1}{4} = \frac{\epsilon}{4(2-\epsilon)}$.
Choose $\delta = \frac{\epsilon}{4(2+\epsilon)}$, the smaller of the two values. Then $0 < \left|x-\frac{1}{4}\right| < \delta \Rightarrow \left|\frac{1}{2x}-2\right| < \epsilon$ and $\lim_{x\to \frac{1}{4}} \frac{1}{2x} = 2$.
By the continuity test, $g(x)$ is continuous at $x = \frac{1}{4}$.

9. Show $\lim_{x\to 2} h(x) = \lim_{x\to 2} \sqrt{2x-3} = 1 = h(2)$.
Step 1: $\left|\sqrt{2x-3}-1\right| < \epsilon \Rightarrow -\epsilon < \sqrt{2x-3}-1 < \epsilon \Rightarrow 1-\epsilon < \sqrt{2x-3} < 1+\epsilon \Rightarrow \frac{(1-\epsilon)^2+3}{2} < x < \frac{(1+\epsilon)^2+3}{2}$.
Step 2: $|x-2| < \delta \Rightarrow -\delta < x-2 < \delta$ or $-\delta+2 < x < \delta+2$.
Then $-\delta+2 = \frac{(1-\epsilon)^2+3}{2} \Rightarrow \delta = 2-\frac{(1-\epsilon)^2+3}{2} = \frac{1-(1-\epsilon)^2}{2} = \epsilon-\frac{\epsilon^2}{2}$, or $\delta+2 = \frac{(1+\epsilon)^2+3}{2} \Rightarrow \delta = \frac{(1+\epsilon)^2+3}{2}-2 = \frac{(1+\epsilon)^2-1}{2} =$ $\epsilon+\frac{\epsilon^2}{2}$. Choose $\delta = \epsilon-\frac{\epsilon^2}{2}$, the smaller of the two values. Then, $0 < |x-2| < \delta \Rightarrow \left|\sqrt{2x-3}-1\right| < \epsilon$,
so $\lim_{x\to 2} \sqrt{2x-3} = 1$. By the continuity test, $h(x)$ is continuous at $x = 2$.

10. Show $\lim_{x\to 5} F(x) = \lim_{x\to 5} \sqrt{9-x} = 2 = F(5)$.
Step 1: $\left|\sqrt{9-x}-2\right| < \epsilon \Rightarrow -\epsilon < \sqrt{9-x}-2 < \epsilon \Rightarrow 9-(2-\epsilon)^2 > x > 9-(2+\epsilon)^2$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

Step 2: $0<|x-5|<\delta \Rightarrow -\delta<x-5<\delta \Rightarrow -\delta+5<x<\delta+5$.
Then $-\delta+5=9-(2+\epsilon)^2 \Rightarrow \delta=(2+\epsilon)^2-4=\epsilon^2+2\epsilon$, or $\delta+5=9-(2-\epsilon)^2 \Rightarrow \delta=4-(2-\epsilon)^2=\epsilon^2-2\epsilon$.
Choose $\delta=\epsilon^2-2\epsilon$, the smaller of the two values. Then, $0<|x-5|<\delta \Rightarrow \left|\sqrt{9-x}-2\right|<\epsilon$, so $\lim\limits_{x\to 5}\sqrt{9-x}=2$.
By the continuity test, $F(x)$ is continuous at $x=5$.

11. Suppose L_1 and L_2 are two different limits. Without loss of generality assume $L_2>L_1$. Let $\epsilon=\frac{1}{3}(L_2-L_1)$. Since $\lim\limits_{x\to x_0} f(x)=L_1$ there is a $\delta_1>0$ such that $0<|x-x_0|<\delta_1 \Rightarrow |f(x)-L_1|<\epsilon \Rightarrow -\epsilon<f(x)-L_1<\epsilon$
$\Rightarrow -\frac{1}{3}(L_2-L_1)+L_1<f(x)<\frac{1}{3}(L_2-L_1)+L_1 \Rightarrow 4L_1-L_2<3f(x)<2L_1+L_2$. Likewise, $\lim\limits_{x\to x_0} f(x)=L_2$ so there is a δ_2
such that $0<|x-x_0|<\delta_2 \Rightarrow |f(x)-L_2|<\epsilon \Rightarrow -\epsilon<f(x)-L_2<\epsilon \Rightarrow -\frac{1}{3}(L_2-L_1)+L_2<f(x)<\frac{1}{3}(L_2-L_1)+L_2$
$\Rightarrow 2L_2+L_1<3f(x)<4L_2-L_1 \Rightarrow L_1-4L_2<-3f(x)<-2L_2-L_1$. If $\delta=\min\{\delta_1,\delta_2\}$ both inequalities must hold for $0<|x-x_0|<\delta$: $\left.\begin{array}{l} 4L_1-L_2<3f(x)<2L_1+L_2 \\ L_1-4L_2<-3f(x)<-2L_2-L_1 \end{array}\right\} \Rightarrow 5(L_1-L_2)<0<L_1-L_2$. That is, $L_1-L_2<0$ and $L_1-L_2>0$, a contradiction.

12. Suppose $\lim\limits_{x\to c} f(x)=L$. If $k=0$, then $\lim\limits_{x\to c} kf(x)=\lim\limits_{x\to c} 0=0=0\cdot\lim\limits_{x\to c} f(x)$ and we are done. If $k\neq 0$, then given any $\epsilon>0$, there is a $\delta>0$ so that $0<|x-c|<\delta<\Rightarrow |f(x)-L|<\frac{\epsilon}{|k|} \Rightarrow |k||f(x)-L|<\epsilon \Rightarrow |k(f(x)-L)|<\epsilon$
$\Rightarrow |(kf(x))-(kL)|<\epsilon$. Thus $\lim\limits_{x\to c} kf(x)=kL=k\left(\lim\limits_{x\to c} f(x)\right)$.

13. (a) Since $x\to 0^+, 0<x^3<x<1 \Rightarrow (x^3-x)\to 0^- \Rightarrow \lim\limits_{x\to 0^+} f(x^3-x)=\lim\limits_{y\to 0^-} f(y)=B$ where $y=x^3-x$.

(b) Since $x\to 0^-, -1<x<x^3<0 \Rightarrow (x^3-x)\to 0^+ \Rightarrow \lim\limits_{x\to 0^-} f(x^3-x)=\lim\limits_{y\to 0^+} f(y)=A$ where $y=x^3-x$.

(c) Since $x\to 0^+, 0<x^4<x^2<1 \Rightarrow (x^2-x^4)\to 0^+ \Rightarrow \lim\limits_{x\to 0^+} f(x^2-x^4)=\lim\limits_{y\to 0^+} f(y)=A$ where $y=x^2-x^4$.

(d) Since $x\to 0^-, -1<x<0 \Rightarrow 0<x^4<x^2<1 \Rightarrow (x^2-x^4)\to 0^+ \Rightarrow \lim\limits_{x\to 0^+} f(x^2-x^4)=A$ as in part (c).

14. (a) True, because if $\lim\limits_{x\to a}(f(x)+g(x))$ exists then $\lim\limits_{x\to a}(f(x)+g(x))-\lim\limits_{x\to a} f(x)=\lim\limits_{x\to a}[(f(x)+g(x))-f(x)]=\lim\limits_{x\to a} g(x)$ exists, contrary to assumption.

(b) False; for example take $f(x)=\frac{1}{x}$ and $g(x)=-\frac{1}{x}$. Then neither $\lim\limits_{x\to 0} f(x)$ nor $\lim\limits_{x\to 0} g(x)$ exists, but $\lim\limits_{x\to 0}(f(x)+g(x))=\lim\limits_{x\to 0}\left(\frac{1}{x}-\frac{1}{x}\right)=\lim\limits_{x\to 0} 0=0$ exists.

(c) True, because $g(x)=|x|$ is continuous $\Rightarrow g(f(x))=|f(x)|$ is continuous (it is the composite of continuous functions).

(d) False; for example let $f(x)=\begin{cases} -1, & x\le 0 \\ 1, & x>0 \end{cases} \Rightarrow f(x)$ is discontinuous at $x=0$. However $|f(x)|=1$ is continuous at $x=0$.

15. Show $\lim\limits_{x\to -1} f(x)=\lim\limits_{x\to -1}\frac{x^2-1}{x+1}=\lim\limits_{x\to -1}\frac{(x+1)(x-1)}{(x+1)}=-2, x\neq -1$.

Define the continuous extension of $f(x)$ as $F(x)=\begin{cases} \frac{x^2-1}{x+1}, & x\neq -1 \\ -2, & x=-1 \end{cases}$. We now prove the limit of $f(x)$ as $x\to -1$ exists and has the correct value.
Step 1: $\left|\frac{x^2-1}{x+1}-(-2)\right|<\epsilon \Rightarrow -\epsilon<\frac{(x+1)(x-1)}{(x+1)}+2<\epsilon \Rightarrow -\epsilon<(x-1)+2<\epsilon, x\neq -1 \Rightarrow -\epsilon-1<x<\epsilon-1$.
Step 2: $|x-(-1)|<\delta \Rightarrow -\delta<x+1<\delta \Rightarrow -\delta-1<x<\delta-1$.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

Then $-\delta-1=-\epsilon-1\Rightarrow\delta=\epsilon$, or $\delta-1=\epsilon-1\Rightarrow\delta=\epsilon$. Choose $\delta=\epsilon$. Then $0<|x-(-1)|<\delta$ $\Rightarrow\left|\frac{x^2-1}{x+1}-(-2)\right|<\epsilon\Rightarrow\lim_{x\to-1}F(x)=-2$. Since the conditions of the continuity test are met by $F(x)$, then $f(x)$ has a continuous extension to $F(x)$ at $x=-1$.

16. Show $\lim_{x\to3}g(x)=\lim_{x\to3}\frac{x^2-2x-3}{2x-6}=\lim_{x\to3}\frac{(x-3)(x+1)}{2(x-3)}=2,\ x\neq3$.

Define the continuous extension of $g(x)$ as $G(x)=\begin{cases}\frac{x^2-2x-3}{2x-6}, & x\neq3\\ 2, & x=3\end{cases}$. We now prove the limit of $g(x)$ as $x\to3$ exists and has the correct value.

Step 1: $\left|\frac{x^2-2x-3}{2x-6}-2\right|<\epsilon\Rightarrow-\epsilon<\frac{(x-3)(x+1)}{2(x-3)}-2<\epsilon\Rightarrow-\epsilon<\frac{x+1}{2}-2<\epsilon,\ x\neq3\Rightarrow3-2\epsilon<x<3+2\epsilon$.

Step 2: $|x-3|<\delta\Rightarrow-\delta<x-3<\delta\Rightarrow3-\delta<x<\delta+3$.

Then, $3-\delta=3-2\epsilon\Rightarrow\delta=2\epsilon$, or $\delta+3=3+2\epsilon\Rightarrow\delta=2\epsilon$. Choose $\delta=2\epsilon$. Then $0<|x-3|<\delta$ $\Rightarrow\left|\frac{x^2-2x-3}{2x-6}-2\right|<\epsilon\Rightarrow\lim_{x\to3}\frac{(x-3)(x+1)}{2(x-3)}=2$. Since the conditions of the continuity test hold for $G(x)$, $g(x)$ can be continuously extended to $G(x)$ at $x=3$.

17. (a) Let $\epsilon>0$ be given. If x is rational, then $f(x)=x\Rightarrow|f(x)-0|=|x-0|<\epsilon\Leftrightarrow|x-0|<\epsilon$; i.e., choose $\delta=\epsilon$. Then $|x-0|<\delta\Rightarrow|f(x)-0|<\epsilon$ for x rational. If x is irrational, then $f(x)=0\Rightarrow|f(x)-0|<\epsilon\Leftrightarrow0<\epsilon$ which is true no matter how close irrational x is to 0, so again we can choose $\delta=\epsilon$. In either case, given $\epsilon>0$ there is a $\delta=\epsilon>0$ such that $0<|x-0|<\delta\Rightarrow|f(x)-0|<\epsilon$. Therefore, f is continuous at $x=0$.

(b) Choose $x=c>0$. Then within any interval $(c-\delta,c+\delta)$ there are both rational and irrational numbers. If c is rational, pick $\epsilon=\frac{c}{2}$. No matter how small we choose $\delta>0$ there is an irrational number x in $(c-\delta,c+\delta)\Rightarrow|f(x)-f(c)|=|0-c|=c>\frac{c}{2}=\epsilon$. That is, f is not continuous at any rational $c>0$. On the other hand, suppose c is irrational $\Rightarrow f(c)=0$. Again pick $\epsilon=\frac{c}{2}$. No matter how small we choose $\delta>0$ there is a rational number x in $(c-\delta,c+\delta)$ with $|x-c|<\frac{c}{2}=\varepsilon\Leftrightarrow\frac{c}{2}<x<\frac{3c}{2}$. Then $|f(x)-f(c)|=|x-0|$ $=|x|>\frac{c}{2}=\epsilon\Rightarrow f$ is not continuous at any irrational $c>0$.

If $x=c<0$, repeat the argument picking $\epsilon=\frac{|c|}{2}=\frac{-c}{2}$. Therefore f fails to be continuous at any nonzero value $x=c$.

18. (a) Let $c=\frac{m}{n}$ be a rational number in $[0,1]$ reduced to lowest terms $\Rightarrow f(c)=\frac{1}{n}$. Pick $\epsilon=\frac{1}{2n}$. No matter how small $\delta>0$ is taken, there is an irrational number x in the interval $(c-\delta,c+\delta)\Rightarrow|f(x)-f(c)|$ $=\left|0-\frac{1}{n}\right|=\frac{1}{n}>\frac{1}{2n}=\epsilon$. Therefore f is discontinuous at $x=c$, a rational number.

(b) Now suppose c is an irrational number $\Rightarrow f(c)=0$. Let $\epsilon>0$ be given. Notice that $\frac{1}{2}$ is the only rational number reduced to lowest terms with denominator 2 and belonging to $[0,1]$; $\frac{1}{3}$ and $\frac{2}{3}$ the only rationals with denominator 3 belonging to $[0,1]$; $\frac{1}{4}$ and $\frac{3}{4}$ with denominator 4 in $[0,1]$; $\frac{1}{5}$, $\frac{2}{5}$, $\frac{3}{5}$ and $\frac{4}{5}$ with denominator 5 in $[0,1]$; etc. In general, choose N so that $\frac{1}{N}<\epsilon\Rightarrow$ there exist only finitely many rationals in $[0,1]$ having denominator $\leq N$, say $r_1,r_2,\ldots,r_p$. Let $\delta=\min\{|c-r_i|:i=1,\ldots,p\}$. Then the interval $(c-\delta,c+\delta)$ contains no rational numbers with denominator $\leq N$. Thus, $0<|x-c|<\delta\Rightarrow|f(x)-f(c)|=|f(x)-0|=$ $|f(x)|\leq\frac{1}{N}<\epsilon\Rightarrow f$ is continuous at $x=c$ irrational.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(c) The graph looks like the markings on a typical ruler when the points $(x, f(x))$ on the graph of $f(x)$ are connected to the x-axis with vertical lines.



$$f(x) = \begin{cases} 1/n & \text{if } x = m/n \text{ is a rational number in lowest terms} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

19. Yes. Let R be the radius of the equator (earth) and suppose at a fixed instant of time we label noon as the zero point, 0, on the equator $\Rightarrow 0 + \pi R$ represents the midnight point (at the same exact time). Suppose x_1 is a point on the equator "just after" noon $\Rightarrow x_1 + \pi R$ is simultaneously "just after" midnight. It seems reasonable that the temperature T at a point just after noon is hotter than it would be at the diametrically opposite point just after midnight: That is, $T(x_1) - T(x_1 + \pi R) > 0$. At exactly the same moment in time pick x_2 to be a point just before midnight $\Rightarrow x_2 + \pi R$ is just before noon. Then $T(x_2) - T(x_2 + \pi R) < 0$. Assuming the temperature function T is continuous along the equator (which is reasonable), the Intermediate Value Theorem says there is a point c between 0 (noon) and πR (simultaneously midnight) such that $T(c) - T(c + \pi R) = 0$; i.e., there is always a pair of antipodal points on the earth's equator where the temperatures are the same.

20. $$\lim_{x\to c} f(x)g(x) = \lim_{x\to c} \tfrac{1}{4}\left[(f(x)+g(x))^2 - (f(x)-g(x))^2\right] = \tfrac{1}{4}\left[\left(\lim_{x\to c}(f(x)+g(x))\right)^2 - \left(\lim_{x\to c}(f(x)-g(x))\right)^2\right]$$
$$= \tfrac{1}{4}(3^2 - (-1)^2) = 2.$$

21. (a) At $x = 0$: $\lim\limits_{a\to 0} r_+(a) = \lim\limits_{a\to 0} \dfrac{-1+\sqrt{1+a}}{a} = \lim\limits_{a\to 0}\left(\dfrac{-1+\sqrt{1+a}}{a}\right)\left(\dfrac{-1-\sqrt{1+a}}{-1-\sqrt{1+a}}\right) = \lim\limits_{a\to 0} \dfrac{1-(1+a)}{a(-1-\sqrt{1+a})} = \dfrac{-1}{-1-\sqrt{1+0}} = \dfrac{1}{2}$

At $x = -1$: $\lim\limits_{a\to -1^+} r_+(a) = \lim\limits_{a\to -1^+} \dfrac{1-(1+a)}{a(-1-\sqrt{1+a})} = \lim\limits_{a\to -1} \dfrac{-a}{a(-1-\sqrt{1+a})} = \dfrac{-1}{-1-\sqrt{0}} = 1$

(b) At $x = 0$: $\lim\limits_{a\to 0^-} r_-(a) = \lim\limits_{a\to 0^-} \dfrac{-1-\sqrt{1+a}}{a} = \lim\limits_{a\to 0^-}\left(\dfrac{-1-\sqrt{1+a}}{a}\right)\left(\dfrac{-1+\sqrt{1+a}}{-1+\sqrt{1+a}}\right) = \lim\limits_{a\to 0^-} \dfrac{1-(1+a)}{a(-1+\sqrt{1+a})} = \lim\limits_{a\to 0^-} \dfrac{-a}{a(-1+\sqrt{1+a})}$

$= \lim\limits_{a\to 0^-} \dfrac{-1}{-1+\sqrt{1+a}} = \infty$ (because the denominator is always negative); $\lim\limits_{a\to 0^+} r_-(a)$

$= \lim\limits_{a\to 0^+} \dfrac{-1}{-1+\sqrt{1+a}} = -\infty$ (because the denominator is always positive).

Therefore, $\lim\limits_{a\to 0} r_-(a)$ does not exist.

At $x = -1$: $\lim\limits_{a\to -1^+} r_-(a) = \lim\limits_{a\to -1^+} \dfrac{-1-\sqrt{1+a}}{a} = \lim\limits_{a\to -1^+} \dfrac{-1}{-1+\sqrt{1+a}} = 1$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

(c)





(d)





22. $f(x) = x + 2\ \cos\ x \Rightarrow f(0) = 0 + 2\ \cos\ 0 = 2 > 0$ and $f(-\pi) = -\pi + 2\ \cos(-\pi) = -\pi - 2 < 0$. Since $f(x)$ is continuous on $[-\pi, 0]$, by the Intermediate Value Theorem, $f(x)$ must take on every value between $[-\pi - 2, 2]$. Thus there is some number c in $[-\pi, 0]$ such that $f(c) = 0$; i.e., c is a solution to $x + 2\ \cos\ x = 0$.

23. (a) The function f is bounded on D if $f(x) \geq M$ and $f(x) \leq N$ for all x in D. This means $M \leq f(x) \leq N$ for all x in D. Choose B to be max $\{|M|,\ |N|\}$. Then $|f(x)| \leq B$. On the other hand, if $|f(x)| \leq B$, then $-B \leq f(x) \leq B \Rightarrow f(x) \geq -B$ and $f(x) \leq B \Rightarrow f(x)$ is bounded on D with $N = B$ an upper bound and $M = -B$ a lower bound.

(b) Assume $f(x) \leq N$ for all x and that $L > N$. Let $\epsilon = \frac{L-N}{2}$. Since $\lim_{x \to x_0} f(x) = L$ there is a $\delta > 0$ such that $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \epsilon \Leftrightarrow L - \epsilon < f(x) < L + \epsilon \Leftrightarrow L - \frac{L-N}{2} < f(x) < L + \frac{L-N}{2} \Leftrightarrow \frac{L+N}{2} < f(x) < \frac{3L-N}{2}$. But $L > N \Rightarrow \frac{L+N}{2} > N \Rightarrow N < f(x)$ contrary to the boundedness assumption $f(x) \leq N$. This contradiction proves $L \leq N$.

(c) Assume $M \leq f(x)$ for all x and that $L < M$. Let $\epsilon = \frac{M-L}{2}$. As in part (b), $0 < |x - x_0| < \delta \Rightarrow L - \frac{M-L}{2} < f(x) < L + \frac{M-L}{2} \Leftrightarrow \frac{3L-M}{2} < f(x) < \frac{M+L}{2} < M$, a contradiction.

24. (a) If $a \geq b$, then $a - b \geq 0 \Rightarrow |a - b| = a - b \Rightarrow \max\ \{a, b\} = \frac{a+b}{2} + \frac{|a-b|}{2} = \frac{a+b}{2} + \frac{a-b}{2} = \frac{2a}{2} = a$.

If $a \leq b$, then $a - b \leq 0 \Rightarrow |a - b| = -(a - b) = b - a \Rightarrow \max\ \{a, b\} = \frac{a+b}{2} + \frac{|a-b|}{2} = \frac{a+b}{2} + \frac{b-a}{2} = \frac{2b}{2} = b$.

(b) Let $\min\ \{a, b\} = \frac{a+b}{2} - \frac{|a-b|}{2}$.

25. $\lim_{x \to 0} = \frac{\sin(1-\cos\ x)}{x} = \lim_{x \to 0} \frac{\sin(1-\cos\ x)}{1-\cos x} \cdot \frac{1-\cos\ x}{x} \cdot \frac{1+\cos\ x}{1+\cos\ x} = \lim_{x \to 0} \frac{\sin(1-\cos x)}{1-\cos x} \cdot \lim_{x \to 0} \frac{1-\cos^2\ x}{x(1+\cos\ x)}$

$= 1 \cdot \lim_{x \to 0} \frac{\sin^2 x}{x(1+\cos\ x)} = \lim_{x \to 0} \frac{\sin\ x}{x} \cdot \frac{\sin\ x}{1+\cos\ x} = 1 \cdot \left(\frac{0}{2}\right) = 0.$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

26. $\lim_{x\to 0^+} \frac{\sin x}{\sin\sqrt{x}} = \lim_{x\to 0^+} \frac{\sin x}{x}\cdot\frac{\sqrt{x}}{\sin\sqrt{x}}\cdot\frac{x}{\sqrt{x}} = 1\cdot \lim_{x\to 0^+} \frac{1}{\left(\frac{\sin\sqrt{x}}{\sqrt{x}}\right)}\cdot \lim_{x\to 0^+}\sqrt{x} = 1\cdot 1\cdot 0 = 0.$

27. $\lim_{x\to 0} \frac{\sin(\sin x)}{x} = \lim_{x\to 0} \frac{\sin(\sin x)}{\sin x}\cdot\frac{\sin x}{x} = \lim_{x\to 0} \frac{\sin(\sin x)}{\sin x}\cdot \lim_{x\to 0}\frac{\sin x}{x} = 1\cdot 1 = 1.$

28. $\lim_{x\to 0} \frac{\sin(x^2+x)}{x} = \lim_{x\to 0} \frac{\sin(x^2+x)}{x^2+x}\cdot(x+1) = \lim_{x\to 0} \frac{\sin(x^2+x)}{x^2+x}\cdot \lim_{x\to 0}(x+1) = 1\cdot 1 = 1.$

29. $\lim_{x\to 2} \frac{\sin(x^2-4)}{x-2} = \lim_{x\to 2} \frac{\sin(x^2-4)}{x^2-4}\cdot(x+2) = \lim_{x\to 2} \frac{\sin(x^2-4)}{x^2-4}\cdot \lim_{x\to 2}(x+2) = 1\cdot 4 = 4.$

30. $\lim_{x\to 9} \frac{\sin(\sqrt{x}-3)}{x-9} = \lim_{x\to 9} \frac{\sin(\sqrt{x}-3)}{\sqrt{x}-3}\cdot\frac{1}{\sqrt{x}+3} = \lim_{x\to 9} \frac{\sin(\sqrt{x}-3)}{\sqrt{x}-3}\cdot \lim_{x\to 9}\frac{1}{\sqrt{x}+3} = 1\cdot\frac{1}{6} = \frac{1}{6}.$

31. Since the highest power of x in the numerator is 1 more than the highest power of x in the denominator, there is an oblique asymptote. $y = \frac{2x^{3/2}+2x-3}{\sqrt{x}+1} = 2x - \frac{3}{\sqrt{x}+1}$, thus the oblique asymptote is $y = 2x$.

32. As $x \to \pm\infty$, $\frac{1}{x} \to 0 \Rightarrow \sin\left(\frac{1}{x}\right) \to 0 \Rightarrow 1+\sin\left(\frac{1}{x}\right) \to 1$, thus as $x \to \pm\infty$, $y = x + x\sin\left(\frac{1}{x}\right) = x\left(1+\sin\left(\frac{1}{x}\right)\right) \to x$; thus the oblique asymptote is $y = x$.

33. As $x \to \pm\infty$, $x^2+1 \to x^2 \Rightarrow \sqrt{x^2+1} \to \sqrt{x^2}$; as $x \to -\infty$, $\sqrt{x^2} = -x$, and as $x \to +\infty$, $\sqrt{x^2} = x$; thus the oblique asymptotes are $y = x$ and $y = -x$.

34. As $x \to \pm\infty$, $x+2 \to x \Rightarrow \sqrt{x^2+2x} = \sqrt{x(x+2)} \to \sqrt{x^2}$; as $x \to -\infty$, $\sqrt{x^2} = -x$, and as $x \to +\infty$, $\sqrt{x^2} = x$; asymptotes are $y = x$ and $y = -x$.

35. Assume $1 < a < b$ and $\frac{a}{x} + x = \frac{1}{x-b} \Rightarrow a(x-b) + x^2(x-b) = x \Rightarrow$ $f(x) = a(x-b) + x^2(x-b) - x = 0$; f is continuous for all x-values and $f(0) = -ab < 0$, $f(a+b) = a^2 + (a+b)^2 a - (a+b)$
$$= \underbrace{a^2 - a}_{(+)} + \underbrace{(a+b)^2 a - b}_{(+)} > 0.$$
Thus, by the Intermediate Value Theorem there is at least one number c, $0 < c < a+b$, so that $f(c) = 0 \Rightarrow$ $a(c-b) + c^2(c-b) - c = 0 \Rightarrow \frac{a}{c} + c = \frac{1}{c-b}$.

36. (a) $\lim_{x\to 0} \frac{\sqrt{a+bx}-1}{x} = \frac{\text{"}\sqrt{a}-1\text{"}}{0}$, so $\sqrt{a}-1 = 0 \Rightarrow \quad a = 1$, then $\lim_{x\to 0} \frac{\sqrt{1+bx}-1}{x}\cdot\frac{\sqrt{1+bx}+1}{\sqrt{1+bx}+1} = \lim_{x\to 0} \frac{(1+bx)-1}{x(\sqrt{1+bx}+1)}$
$= \lim_{x\to 0} \frac{b}{\sqrt{1+bx}+1} = \frac{b}{2} = 2 \Rightarrow b = 4$

(b) $\lim_{x\to 1} \frac{\tan(ax-a)+b-2}{x-1} = \frac{\text{"}\tan 0+b-2\text{"}}{0} = \frac{\text{"}b-2\text{"}}{0}$, so $b-2 = 0 \Rightarrow b = 2$, then $\lim_{x\to 1} \frac{\tan(ax-a)}{x-1} = \lim_{x\to 1} a\cdot\frac{\tan a(x-1)}{a(x-1)}$
$= \lim_{x\to 1} \frac{a}{\cos a(x-1)}\cdot\frac{\sin a(x-1)}{a(x-1)} = \frac{a}{\cos 0}\cdot 1 = a = 3$

37. $\lim_{x\to 1} \frac{x^{2/3}-1}{1-x^{1/2}} = \lim_{x\to 1} \frac{(x^{1/6})^4-1^4}{1^3-(x^{1/6})^3} = \lim_{x\to 1} \frac{(x^{1/6}-1)(x^{1/6}+1)((x^{1/6})^2+1)}{(1-x^{1/6})(1+(x^{1/6})+(x^{1/6})^2)} = \lim_{x\to 1} \frac{-(x^{1/6}+1)(x^{1/3}+1)}{1+x^{1/6}+x^{1/3}} = \frac{-(2)(2)}{3} = \frac{-4}{3}$

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual

38. $\lim\limits_{x\to 0^+} \dfrac{|3x+4|-|x|-4}{x} = \lim\limits_{x\to 0^+} \dfrac{(3x+4)-x-4}{x} = \lim\limits_{x\to 0^+} \dfrac{2x}{2} = 2$; assume $x > \dfrac{-3}{4} \Rightarrow \lim\limits_{x\to 0^-} \dfrac{|3x+4|-|x|-4}{x} = \lim\limits_{x\to 0^-} \dfrac{(3x+4)-(-x)-4}{x}$

$\lim\limits_{x\to 0^-} \dfrac{4x}{x} = 4 \Rightarrow \lim\limits_{x\to 0} \dfrac{|3x+4|-|x|-4}{x}$ does not exist.

Copyright © 2018 Pearson Education, Inc.

Thomas Calculus Early Transcendentals 14th edition Hass Solutions Manual