

***Graphical Approach to Precalculus with Limits 7th
edition Hornsby Solutions Manual***

SKU: 9780134696492-SOLUTIONS

Chapter 2: Analysis of Graphs of Functions

2.1: Graphs of Basic Functions and Relations; Symmetry

1. $(-\infty, \infty)$.
2. $(-\infty, \infty); [0, \infty)$
3. $(0, 0)$
4. $[0, \infty); [0, \infty)$
5. increases
6. $(-\infty, 0]; [0, \infty)$
7. x -axis
8. even
9. odd
10. y -axis; origin
11. The domain can be all real numbers; therefore, the function is continuous for the interval $(-\infty, \infty)$.
12. The domain can be all real numbers; therefore, the function is continuous for the interval $(-\infty, \infty)$.
13. The domain can only be values where $x \geq 0$; therefore, the function is continuous for the interval $[0, \infty)$.
14. The domain can only be values where $x \leq 0$; therefore, the function is continuous for the interval $(-\infty, 0]$.
15. The domain can be all real numbers except -3 ; therefore, the function is continuous for the interval $(-\infty, -3) \cup (-3, \infty)$.
16. The domain can be all real numbers except 1 ; therefore, the function is continuous for the interval $(-\infty, 1) \cup (1, \infty)$.
17.
 - (a) The function is increasing for the interval $(3, \infty)$
 - (b) The function is decreasing for the interval $(-\infty, 3)$
 - (c) The function is never constant; therefore, none.
 - (d) The domain can be all real numbers; therefore, the interval $(-\infty, \infty)$.
 - (e) The range can only be values where $y \geq 0$; therefore, the interval $[0, \infty)$.
18.
 - (a) The function is increasing for the interval $(4, \infty)$
 - (b) The function is decreasing for the interval $(-\infty, -1)$
 - (c) The function is constant for the interval $(-1, 4)$
 - (d) The domain can be all real numbers; therefore, the interval $(-\infty, \infty)$.
 - (e) The range can only be values where $y \geq 3$; therefore, the interval $[3, \infty)$.
19.
 - (a) The function is increasing for the interval $(-\infty, 1)$
 - (b) The function is decreasing for the interval $(4, \infty)$

- (c) The function is constant for the interval $(1, 4)$
- (d) The domain can be all real numbers; therefore, the interval $(-\infty, \infty)$.
- (e) The range can only be values where $y \leq 3$; therefore, the interval $(-\infty, 3]$.
20. (a) The function is never increasing; therefore, none.
- (b) The function is always decreasing; therefore, the interval $(-\infty, \infty)$.
- (c) The function is never constant; therefore, none.
- (d) The domain can be all real numbers; therefore, the interval $(-\infty, \infty)$.
- (e) The range can be all real numbers; therefore, the interval $(-\infty, \infty)$.
21. (a) The function is never increasing; therefore, none
- (b) The function is decreasing for the intervals $(-\infty, -2)$ and $(3, \infty)$
- (c) The function is constant for the interval $(-2, 3)$.
- (d) The domain can be all real numbers; therefore, the interval $(-\infty, \infty)$.
- (e) The range can only be values where $y \leq 1.5$ or $y \geq 2$; therefore, the interval $(-\infty, 1.5] \cup [2, \infty)$.
22. (a) The function is increasing for the interval $(3, \infty)$.
- (b) The function is decreasing for the interval $(-\infty, -3)$.
- (c) The function is constant for the interval $(-3, 3)$
- (d) The domain can be all real numbers except -3 ; therefore, the interval $(-\infty, -3) \cup (-3, \infty)$.
- (e) The range can only be values where $y > 1$; therefore, the interval $(1, \infty)$.
23. Graph $f(x) = x^5$. See Figure 23. As x increases for the interval $(-\infty, \infty)$, y increases; therefore, the function is increasing.
24. Graph $f(x) = -x^3$. See Figure 24. As x increases for the interval $(-\infty, \infty)$, y decreases; therefore, the function is decreasing.
25. Graph $f(x) = x^4$. See Figure 25. As x increases for the interval $(-\infty, 0)$ y decreases; therefore, the function is decreasing on $(-\infty, 0)$
26. Graph $f(x) = x^4$. See Figure 26. As x increases for the interval $(0, \infty)$, y increases; therefore, the function is increasing on $(0, \infty)$

$[-10, 10]$ by $[-10, 10]$
Xscl = 1 Yscl = 1




Figure 23

$[-10, 10]$ by $[-10, 10]$
Xscl = 1 Yscl = 1




Figure 24

$[-10, 10]$ by $[-10, 10]$
Xscl = 1 Yscl = 1




Figure 25

$[-10, 10]$ by $[-10, 10]$
Xscl = 1 Yscl = 1






Figure 26

27. Graph $f(x) = -|x|$. See Figure 27. As x increases for the interval $(-\infty, 0)$, y increases; therefore, the function is increasing on $(-\infty, 0)$.
28. Graph $f(x) = -|x|$. See Figure 28. As x increases for the interval $(0, \infty)$, y decreases; therefore, the function is decreasing on $(0, \infty)$.
29. Graph $f(x) = -\sqrt[3]{x}$. See Figure 29. As x increases for the interval $(-\infty, \infty)$, y decreases; therefore, the function is decreasing.
30. Graph $f(x) = -\sqrt{x}$. See Figure 30. As x increases for the interval $(0, \infty)$, y decreases; therefore, the function is decreasing.



31. Graph $f(x) = 1 - x^3$. See Figure 31. As x increases for the interval $(-\infty, \infty)$, y decreases; therefore, the function is decreasing.
32. Graph $f(x) = x^2 - 2x$. See Figure 32. As x increases for the interval $(1, \infty)$, y increases; therefore, the function is increasing on $(1, \infty)$.
33. Graph $f(x) = 2 - x^2$. See Figure 33. As x increases for the interval $(-\infty, 0)$, y increases; therefore, the function is increasing on $(-\infty, 0)$.
34. Graph $f(x) = |x + 1|$. See Figure 34. As x increases for the interval $(-\infty, -1)$, y decreases; therefore, the function is decreasing on $(-\infty, -1)$.



35. (a) No (b) Yes (c) No
 36. (a) Yes (b) No (c) No
 37. (a) Yes (b) No (c) No
 38. (a) No (b) No (c) Yes
 39. (a) Yes (b) Yes (c) Yes
 40. (a) Yes (b) Yes (c) Yes
 41. (a) No (b) No (c) Yes
 42. (a) No (b) Yes (c) No
 43. Since the graph is symmetric with respect to the y -axis the function is even.
 44. Since the graph is symmetric with respect to the origin the function is odd.
 45. Since the graph is symmetric with respect to the origin the function is odd.
 46. Since the graph is symmetric with respect to the y -axis the function is even.
 47. Since the graph is not symmetric with respect to the y -axis or origin, the function is neither even nor odd.
 48. Since the graph is not symmetric with respect to the y -axis or origin, the function is neither even nor odd.
 49. (a) Since $f(-x) = f(x)$, this is an even function and is symmetric with respect to the y -axis.
 See Figure 49a.
 (b) Since $f(-x) = -f(x)$, this is an odd function and is symmetric with respect to the origin.
 See Figure 49b.




Figure 49a




Figure 49b

50. (a) Since this is an odd function and is symmetric with respect to the origin. See Figure 50a.
 (b) Since this is an even function and is symmetric with respect to the y -axis. See Figure 50b




Figure 50a




Figure 50b

51. If f is an even function then $f(-x) = f(x)$ or opposite domains have the same range. See Figure 51
 52. If g is an odd function then $g(-x) = -g(x)$ or opposite domains have the opposite range. See Figure 52

x	$f(x)$
-3	21
-2	-12
-1	-25
1	-25
2	-12
3	21

Figure 51

x	$g(x)$
-5	13
-3	1
-2	-5
0	0
2	5
3	-1
5	-13

Figure 52

53. This is an even function since opposite domains have the same range.
54. This is an even function since opposite domains have the same range.
55. This is an odd function since opposite domains have the opposite range.
56. This is an odd function since opposite domains have the opposite range.
57. This is neither even nor odd since the opposite domains are neither the opposite or same range.
58. This is neither even nor odd since the opposite domains are neither the opposite or same range.
59. If $f(x) = x^4 - 7x^2 + 6$, then $f(-x) = (-x)^4 - 7(-x)^2 + 6 \Rightarrow f(-x) = x^4 - 7x^2 + 6$. Since $f(-x) = f(x)$, the function is even.
60. If $f(x) = -2x^6 - 8x^2$, then $f(-x) = -2(-x)^6 - 8(-x)^2 \Rightarrow f(-x) = -2x^6 - 8x^2$. Since $f(-x) = f(x)$, the function is even.
61. If $f(x) = 3x^3 - x$, then $f(-x) = 3(-x)^3 - (-x) \Rightarrow f(-x) = -3x^3 + x$ and $-f(x) = -(3x^3 - x) \Rightarrow -f(x) = -3x^3 + x$. Since $f(-x) = -f(x)$, the function is odd.
62. If $f(x) = -x^5 + 2x^3 - 3x$, then $f(-x) = -(-x)^5 + 2(-x)^3 - 3(-x) \Rightarrow f(-x) = x^5 - 2x^3 + 3x$ and $-f(x) = -(-x^5 + 2x^3 - 3x) \Rightarrow -f(x) = x^5 - 2x^3 + 3x$. Since $f(-x) = -f(x)$, the function is odd.
63. If $f(x) = x^6 - 4x^4 + 5$ then $f(-x) = (-x)^6 - 4(-x)^4 + 5 \Rightarrow f(-x) = x^6 - 4x^4 + 5$. Since $f(-x) = f(x)$, the function is even.
64. If $f(x) = 8$, then $f(-x) = 8$. Since $f(-x) = f(x)$, the function is even.
65. If $f(x) = 3x^5 - x^3 + 7x$, then $f(-x) = 3(-x)^5 - (-x)^3 + 7(-x) \Rightarrow f(-x) = -3x^5 + x^3 - 7x$ and $-f(x) = -(3x^5 - x^3 + 7x) \Rightarrow -f(x) = -3x^5 + x^3 - 7x$. Since $f(-x) = -f(x)$, the function is odd.
66. If $f(x) = x^3 - 4x$, then $f(-x) = (-x)^3 - 4(-x) \Rightarrow f(-x) = -x^3 + 4x$ and $-f(x) = -(x^3 - 4x) \Rightarrow -f(x) = -x^3 + 4x$. Since $f(-x) = -f(x)$, the function is odd.
67. If $f(x) = |5x|$, then $f(-x) = |5(-x)| \Rightarrow f(-x) = |5x|$. Since $f(-x) = f(x)$, the function is even.
68. If $f(x) = \sqrt{x^2 + 1}$, then $f(-x) = \sqrt{(-x)^2 + 1} \Rightarrow f(-x) = \sqrt{x^2 + 1}$. Since $f(-x) = f(x)$, the function is even.

69. If $(-3, 11)$ and $(2, 9)$ then $f(-x) = \frac{1}{2(-x)} \Rightarrow f(-x) = -\frac{1}{2x}$ and $-f(x) = -\left(\frac{1}{2x}\right) \Rightarrow -f(x) = -\frac{1}{2x}$. Since $f(-x) = -f(x)$, the function is odd.
70. If $f(x) = 4x - \frac{1}{x}$, then $f(-x) = 4(-x) - \frac{1}{(-x)} \Rightarrow f(-x) = -4x + \frac{1}{x}$ and $-f(x) = -\left(4x - \frac{1}{x}\right) \Rightarrow -f(x) = -4x + \frac{1}{x}$. Since $f(-x) = -f(x)$, the function is odd.
71. If $f(x) = \frac{x^3 + 3x}{x}$, then $f(-x) = \frac{(-x)^3 + 3(-x)}{-x} \Rightarrow f(-x) = \frac{-x^3 - 3x}{-x} = \frac{x^3 + 3x}{x}$. Since $f(-x) = f(x)$, the function is even.
72. If $f(x) = \frac{x^2 + 5}{x}$, then $f(-x) = \frac{(-x)^2 + 5}{(-x)} \Rightarrow f(-x) = \frac{x^2 + 5}{-x} = -\left(\frac{x^2 + 5}{x}\right)$. Since $f(-x) = -f(x)$, the function is odd.
73. If $f(x) = \sqrt[3]{x^3 - 5x}$, then $f(-x) = \sqrt[3]{(-x)^3 - 5(-x)} \Rightarrow f(-x) = \sqrt[3]{-x^3 + 5x} = -\sqrt[3]{x^3 - 5x}$. Since $f(-x) = -f(x)$, the function is odd.
74. If $f(x) = x$ then $f(-x) = -x$. Since $f(-x) = -f(x)$, the function is odd.
75. If $f(x) = \frac{x^3 + 3}{|x|}$, then $f(-x) = \frac{(-x)^3 + 3}{|-x|} \Rightarrow f(-x) = \frac{-x^3 + 3}{|x|}$. Since $f(-x) = f(x)$, the function is even.
76. If $f(x) = 9$, then $f(-x) = 9$. Since $f(-x) = f(x)$, the function is even.
77. If $f(x) = -x^3 + 2x$, then $f(-x) = -(-x)^3 + 2(-x) \Rightarrow f(-x) = x^3 - 2x$ and $-f(x) = -(-x^3 + 2x) \Rightarrow -f(x) = x^3 - 2x$. Since $f(-x) = -f(x)$, the function is symmetric with respect to the origin. Graph $f(x) = -x^3 + 2x$; the graph supports symmetry with respect to the origin.
78. If $f(x) = x^5 - 2x^3$, then $f(-x) = (-x)^5 - 2(-x)^3 \Rightarrow f(-x) = -x^5 + 2x^3$ and $-f(x) = -(x^5 - 2x^3) \Rightarrow -f(x) = -x^5 + 2x^3$. Since $f(-x) = -f(x)$, the function is symmetric with respect to the origin. Graph $f(x) = x^5 - 2x^3$; the graph supports symmetry with respect to the origin.
79. If $f(x) = 0.5x^4 - 2x^2 + 1$, then $f(-x) = 0.5(-x)^4 - 2(-x)^2 + 1 \Rightarrow f(-x) = 0.5x^4 - 2x^2 + 1$. Since $f(-x) = f(x)$, the function is symmetric with respect to the y-axis. Graph $f(x) = 0.5x^4 - 2x^2 + 1$; the graph supports symmetry with respect to the y-axis.
80. If $f(x) = 0.75x^2 + |x| + 1$, then $f(-x) = 0.75(-x)^2 + |(-x)| + 1 \Rightarrow f(-x) = 0.75x^2 + |x| + 1$. Since $f(-x) = f(x)$, the function is symmetric with respect to the y-axis. Graph $f(x) = 0.75x^2 + |x| + 1$; the graph supports symmetry with respect to the y-axis.

81. If $f(x) = x^3 - x + 3$, then $f(-x) = (-x)^3 - (-x) + 3 \Rightarrow f(-x) = -x^3 + x + 3$ and $-f(x) = -(x^3 - x + 3) \Rightarrow -f(x) = -x^3 + x - 3$. Since $f(x) \neq f(-x) \neq -f(x)$, the function is not symmetric with respect to the y -axis or the origin.
82. If $f(x) = x^4 - 5x + 2$, then $f(-x) = (-x)^4 - 5(-x) + 2 \Rightarrow f(-x) = x^4 + 5x + 2$ and $-f(x) = -(x^4 - 5x + 2) \Rightarrow -f(x) = -x^4 + 5x - 2$. Since $f(x) \neq f(-x) \neq -f(x)$, the function is not symmetric with respect to the y -axis or the origin. Graph $f(x) = x^4 - 5x + 2$; the graph supports no symmetry with respect to the y -axis or the origin.
83. If $f(x) = x^6 - 4x^3$, then $f(-x) = (-x)^6 - 4(-x)^3 \Rightarrow f(-x) = x^6 + 4x^3$ and $-f(x) = -(x^6 - 4x^3) \Rightarrow -f(x) = -x^6 + 4x^3$. Since $f(x) \neq f(-x) \neq -f(x)$, the function is not symmetric with respect to the y -axis or the origin. Graph $f(x) = x^6 - 4x^3$; the graph supports no symmetry with respect to the y -axis or the origin.
84. If $f(x) = x^3 - 3x$, then $f(-x) = (-x)^3 - 3(-x) \Rightarrow f(-x) = -x^3 + 3x$ and $-f(x) = -(x^3 - 3x) \Rightarrow -f(x) = -x^3 + 3x$. Since $f(-x) = -f(x)$, the function is symmetric with respect to the origin. Graph $f(x) = x^3 - 3x$; the graph supports symmetry with respect to the origin.
85. If $f(x) = -6$, then $f(-x) = -6$. Since $f(-x) = f(x)$, the function is symmetric with respect to the y -axis. Graph $f(x) = -6$; the graph supports symmetry with respect to the y -axis.
86. If $f(x) = |x|$, then $f(-x) = |-x| \Rightarrow f(-x) = |x|$. Since $f(-x) = f(x)$, the function is symmetric with respect to the y -axis. Graph $f(x) = |x|$; the graph supports symmetry with respect to the y -axis.
87. If $f(x) = \frac{1}{4x^3}$, then $f(-x) = \frac{1}{4(-x)^3} \Rightarrow f(-x) = -\frac{1}{4x^3}$ and $-f(x) = -\left(\frac{1}{4x^3}\right) \Rightarrow -f(x) = -\frac{1}{4x^3}$. Since $f(-x) = -f(x)$, the function is symmetric with respect to the origin. Graph $f(x) = \frac{1}{4x^3}$; the graph supports symmetry with respect to the origin.
88. If $f(x) = \sqrt{x^2} \Rightarrow f(x) = |x|$, then $f(-x) = \sqrt{(-x)^2} \Rightarrow f(-x) = \sqrt{x^2} \Rightarrow f(-x) = |x|$. Since $f(-x) = f(x)$, the function is symmetric with respect to the y -axis. Graph $f(x) = \sqrt{x^2}$; the graph

2.2: Vertical and Horizontal Shifts of Graphs

1. The equation $y = x^2$ shifted 3 units upward is $y = x^2 + 3$.
2. The equation $y = x^3$ shifted 2 units downward is $y = x^3 - 2$.
3. The equation $y = \sqrt{x}$ shifted 4 units downward is $y = \sqrt{x} - 4$.
4. The equation $y = \sqrt[3]{x}$ shifted 6 units upward is $y = \sqrt[3]{x} + 6$.

5. The equation $y = |x|$ shifted 4 units to the right is $y = |x - 4|$.
6. The equation $y = |x|$ shifted 3 units to the left is $y = |x + 3|$.
7. The equation $y = x^3$ shifted 7 units to the left is $y = (x + 7)^3$.
8. The equation $y = \sqrt{x}$ shifted 9 units to the right is $y = \sqrt{x - 9}$.
9. The equation $y = x^2$ shifted 2 units downward and 3 units right is $y = (x - 3)^2 - 2$.
10. The equation $y = x^2$ shifted 4 units upward and 1 unit left is $y = (x + 1)^2 + 4$.
11. The equation $y = \sqrt{x}$ shifted 3 units upward and 6 units to the left is $y = \sqrt{x + 6} + 3$.
12. The equation $y = |x|$ shifted 1 unit downward and 5 units to the right is $y = |x - 5| - 1$.
13. The equation $y = x^2$ shifted 500 units upward and 2000 units right is $y = (x - 2000)^2 + 500$.
14. The equation $y = x^2$ shifted 255 units downward and 1000 units left is $y = (x + 1000)^2 - 255$.
15. Shift the graph of f 4 units upward to obtain the graph of g .
16. Shift the graph of f 4 units to the left to obtain the graph of g .
17. The equation $y = x^2 - 3$ is $y = x^2$ shifted 3 units downward; therefore, graph B.
18. The equation $y = (x - 3)^2$ is $y = x^2$ shifted 3 units to the right; therefore, graph C.
19. The equation $y = (x + 3)^2$ is $y = x^2$ shifted 3 units to the left; therefore, graph A.
20. The equation $y = |x| + 4$ is $y = |x|$ shifted 4 units upward; therefore, graph A.
21. The equation $y = |x + 4| - 3$ is $y = |x|$ shifted 4 units to the left and 3 units downward; therefore, graph B.
22. The equation $y = |x - 4| - 3$ is $y = f(x)$ shifted 4 units to the right and 3 units downward; therefore, graph C.
23. The equation $y = (x - 3)^3$ is $y = x^3$ shifted 3 units to the right; therefore, graph C.
24. The equation $y = (x - 2)^3 - 4$ is $y = x^3$ shifted 2 units to the right and 4 units downward; therefore, graph A.
25. The equation $y = (x + 2)^3 - 4$ is $[-a, -b]$ shifted 2 units to the left and 4 units downward; therefore, graph B.
26. Using $Y_2 = Y_1 + k$ and $x = 0$, we get $19 = 15 + k \Rightarrow k = 4$.
27. Using $Y_2 = Y_1 + k$ and $x = 0$, we get $-5 = -3 + k \Rightarrow k = -2$.
28. Using $Y_2 = Y_1 + k$ and $x = 0$, we get $5.5 = 4 + 1.5 \Rightarrow k = 1.5$.
29. From the graphs, $(6, 2)$ is a point on Y_1 and $(6, -1)$ a point on Y_2 . Using $Y_2 = Y_1 + k$ and $x = 6$, we get $-1 = 2 + k \Rightarrow k = -3$.
30. From the graphs, $(-4, 3)$ is a point on Y_1 and $(-4, 8)$ a point on Y_2 . Using $Y_2 = Y_1 + k$ and $x = -4$, we get $8 = 3 + k \Rightarrow k = 5$.

31. For the equation $y = x^2$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 3 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[-3, \infty)$.
32. For the equation $y = x^2$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 3 units to the right gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[0, \infty)$.
33. For the equation $y = |x|$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 4 units to the left and 3 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[-3, \infty)$.
34. For the equation $y = |x|$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 4 units to the right and 3 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[-3, \infty)$.
35. For the equation $y = x^3$, the Domain is $(-\infty, \infty)$ and the Range is $(-\infty, \infty)$. Shifting this 3 units to the right gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $(-\infty, \infty)$.
36. For the equation $y = x^3$, the Domain is $(-\infty, \infty)$ and the Range is $(-\infty, \infty)$. Shifting this 2 units to the right and 4 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $(-\infty, \infty)$.
37. For the equation $y = x^2$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 1 unit to the right and 5 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[-5, \infty)$.
38. For the equation $y = x^2$, the Domain is $(-\infty, \infty)$ and the Range is $[0, \infty)$. Shifting this 8 units to the left and 3 units upward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $[3, \infty)$.
39. For the equation $y = \sqrt{x}$, the Domain is $[0, \infty)$ and the Range is $[0, \infty)$. Shifting this 4 units to the right gives us: (a) Domain: $[4, \infty)$ (b) Range: $[0, \infty)$.
40. For the equation $y = \sqrt{x}$, the Domain is $[0, \infty)$ and the Range is $[0, \infty)$. Shifting this 1 units to the left and 10 units downward gives us: (a) Domain: $[-1, \infty)$ (b) Range: $[-10, \infty)$.
41. For the equation $y = x^3$, the Domain is $(-\infty, \infty)$ and the Range is $(-\infty, \infty)$. Shifting this 1 unit to the right and 4 units upward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $(-\infty, \infty)$.
42. For the equation $y = \sqrt[3]{x}$, the Domain is $(-\infty, \infty)$ and the Range is $(-\infty, \infty)$. Shifting this 7 units to the left and 10 units downward gives us: (a) Domain: $(-\infty, \infty)$ (b) Range: $(-\infty, \infty)$.
43. The graph of $y = f(x)$ is the graph of the equation $y = x^2$ shifted 1 unit to the right. See Figure 43.
44. The graph of $y = \sqrt{x+2}$ is the graph of the equation $y = \sqrt{x}$ shifted 2 units to the left. See Figure 44.
45. The graph of $y = x^3 + 1$ is the graph of the equation $y = x^3$ shifted 1 unit upward. See Figure 45.




Figure 43




Figure 44




Figure 45

46. The graph of $y = |x + 2|$ is the graph of the equation $y = |x|$ shifted 2 units to the left. See Figure 46.
47. The graph of $y = (x - 1)^3$ is the graph of the equation $y = x^3$ shifted 1 unit to the right. See Figure 47.
48. The graph of $y = |x| - 3$ is the graph of the equation $y = |x|$ shifted 3 units downward. See Figure 48.




Figure 46




Figure 47




Figure 48

49. The graph of $y = \sqrt{x - 2} - 1$ is the graph of the equation $y = \sqrt{x}$ shifted 2 units to the right and 1 unit downward. See Figure 49.
50. The graph of $y = \sqrt{x + 3} - 4$ is the graph of the equation $y = \sqrt{x}$ shifted 3 units to the left and 4 units downward. See Figure 50.
51. The graph of $f(x)$ is the graph of the equation $y = x^2$ shifted 2 units to the left and 3 units upward. See Figure 51.




Figure 49




Figure 50




Figure 51

52. The graph of $y = (x - 4)^2 - 4$ is the graph of the equation $y = x^2$ shifted 4 units to the right and 4 units downward. See Figure 52.
53. The graph of $y = |x + 4| - 2$ is the graph of the equation $y = |x|$ shifted 4 units to the left and 2 units downward. See Figure 53.
54. The graph of $y = (x + 3)^3 - 1$ is the graph of the equation $y = x^3$ shifted 3 units to the left and 1 unit downward. See Figure 54.




Figure 52




Figure 53




Figure 54

55. Since h and k are positive, the equation is $y = x^2$ shifted to the right and down; therefore, B.
56. Since h and k are positive, the equation is $y = x^2$ shifted to the left and down; therefore, D.
57. Since h and k are positive, the equation is $y = x^2$ shifted to the left and up; therefore, A.
58. Since h and k are positive, the equation is $y = x^2$ shifted to the right and up; therefore, C.
59. The equation $y = f(x) + 2$ is $y = f(x)$ shifted up 2 units or add 2 to the y -coordinate of each point as follows: $(-3, 2) \Rightarrow (-3, 0)$; $(-1, 4) \Rightarrow (-1, 6)$; $(5, 0) \Rightarrow (5, 2)$. See Figure 59.
60. The equation $y = f(x) - 2$ is $y = f(x)$ shifted down 2 units or subtract 2 from the y -coordinate of each point as follows: $(-3, 2) \Rightarrow (-3, -4)$; $(-1, 4) \Rightarrow (-1, 2)$; $(5, 0) \Rightarrow (5, -2)$. See Figure 60.




Figure 59




Figure 60

61. The equation $y = f(x + 2)$ is $y = f(x)$ shifted left 2 units or subtract 2 from the x -coordinate of each point as follows: $(-3, -2) \Rightarrow (-5, -2)$; $(-1, 4) \Rightarrow (-3, 4)$; $(5, 0) \Rightarrow (3, 0)$. See Figure 61.
62. The equation $y = f(x - 2)$ is $y = f(x)$ shifted right 2 units or add 2 to the x -coordinate of each point as follows: $(-3, -2) \Rightarrow (-1, -2)$; $(-1, 4) \Rightarrow (1, 4)$; $(5, 0) \Rightarrow (7, 0)$. See Figure 62.




Figure 61




Figure 62

63. The graph is the basic function $y = x^2$ translated 4 units to the left and 3 units up; therefore, the new equation is $y = (x + 4)^2 + 3$. The equation is now increasing for the interval: (a) $(-4, \infty)$ and decreasing for the interval: (b) $(-\infty, -4)$.
64. The graph is the basic function $y = \sqrt{x}$ translated 5 units to the left; therefore, the new equation is $y = \sqrt{x + 5}$. The equation is now increasing for the interval: (a) $(-5, \infty)$ and does not decrease; therefore: (b) none.
65. The graph is the basic function $y = x^3$ translated 5 units down; therefore, the new equation is $y = x^3 - 5$. The equation is now increasing for the interval: (a) $(-\infty, \infty)$ and does not decrease; therefore: (b) none.
66. The graph is the basic function $y = |x|$ translated 10 units to the left; therefore, the new equation is $y = |x + 10|$. The equation is now increasing for the interval: (a) $(-10, \infty)$ and decreasing for the interval: (b) $(-\infty, -10)$.
67. The graph is the basic function $y = \sqrt{x}$ translated 2 units to the right and 1 unit up; therefore, the new equation is $y = \sqrt{x - 2} + 1$. The equation is now increasing for the interval: (a) $(2, \infty)$ and does not decrease; therefore: (b) none.
68. The graph is the basic function $y = x^2$ translated 2 units to the right and 3 units down; therefore, the new equation is $y = (x - 2)^2 - 3$. The equation is now increasing for the interval: (a) $(2, \infty)$ and decreasing for the interval: (b) $(-\infty, 2)$.
69. The translation is 3 units to the left and 1 unit up; therefore, the new equation is $y = |x + 3| + 1$. The form $y = |x - h| + k$ will equal $y = |x + 3| + 1$ when: $h = -3$ and $k = 1$.
70. The equation $y = x^2$ has a Domain: $(-\infty, \infty)$ and a Range: $[0, \infty)$. After the translation the Domain is still: $(-\infty, \infty)$ but now the Range is $(38, \infty)$, a positive or upward shift of 38 units. Therefore, the horizontal shift can be any number of units, but the vertical shift is up 38. This makes h any real number and $k = 38$.
71. (a) $O(9) = 2649.4(9) + 13,260 = 37,104.6$; In 2014, organic food sales totaled \$37,104.6 million.
- (b) We will use the point (2005, 13260) and the slope of 2649.4 in the point slope form for the equation of a line. $y - y_1 = m(x - x_1) \Rightarrow y - 13260 = 2649.4(x - 2005) \Rightarrow y = 2649.4(x - 2005) + 13260$
- (c) $y = 2649.4(2014 - 2005) + 13260 = 37,104.6$
- (d) $26,507 = 2649.4(x - 2005) + 13260 \Rightarrow 12,887 = 2649.4(x - 2005) \Rightarrow \frac{12,887}{2649.4} = x - 2005 \Rightarrow x \approx 2009.86$
- The organic food sales reached \$26,507 around 2010.
72. (a) $S(3) = -23.8(3) + 248 = 176.6$; In 2013, 176.6 million units were sold.

- (b) We will use the point (2010, 248) and the slope of -23.8 in the point slope form for the equation of a line. $y - y_1 = m(x - x_1) \Rightarrow y - 248 = -23.8(x - 2010) \Rightarrow y = -23.8(x - 2010) + 248$
- (c) $y = -23.8(2013 - 2010) + 248 = -23.8(3) + 248 = 176.6$; In 2013, 176.6 million units were sold.
- (d) $129 = -23.8(x - 2010) + 248 \Rightarrow -119 = -23.8(x - 2010) \Rightarrow 5 = x - 2010 \Rightarrow x = 2015$
73. $U(2011) = 13(2011 - 2006)^2 + 115 = 13(25) + 115 = 440$; The average U.S. household spent \$440 on Apple products in 2011.
74. The formula for $W(x)$ can be found by shifting $U(x) = 13(x - 2006)^2 + 115$ to the right 4 units.
 $W(x) = 13(x - 2010)^2 + 115$; $W(x) = 13(2015 - 2010)^2 + 115 = 13(25) + 115 = 440$
 In 2015, the average worldwide household spending on Apple products was \$440, which equaled U.S. spending 4 years earlier.
75. (a) Enter the year in L_1 and enter tuition and fees in L_2 . The year 2000 corresponds to $x = 0$ and so on. The regression equation is $y \approx 397.2x + 3526$.
- (b) Since $x = 0$ corresponds to 2000, the equation when the exact year is entered is
 $y = 397.2(x - 2000) + 3526$
- (c) $y \approx 397.2(2014 - 2000) + 3526 \approx 9087$
76. (a) Enter the year in L_1 and enter the percent of women in the workforce in L_2 . The year 2015 corresponds to $x = 0$ and so on. The regression equation is $y \approx 404.12x + 9136.25$.
- (b) Since $x = 0$ corresponds to 2015, the equation when the exact year is entered is
 $y \approx 404.12(x - 2015) + 9136.25$.
- (c) $y \approx 404.12(2025 - 2015) + 9136.25 \approx 13,177$.

77. See Figure 77.




Figure 77

78. $m = \frac{2 - (-2)}{3 - 1} \Rightarrow m = \frac{4}{2} = 2$

79. Using slope-intercept form yields: $y_1 - 2 = 2(x - 3) \Rightarrow y_1 - 2 = 2x - 6 \Rightarrow y_1 = 2x - 4$

80. $(1, -2 + 6)$ and $(3, 2 + 6) \Rightarrow (1, 4)$ and $(3, 8)$

81. $m = \frac{8-4}{3-1} \Rightarrow m = \frac{4}{2} = 2$

82. Using slope-intercept form yields: $y_2 - 4 = 2(x-1) \Rightarrow y_2 - 4 = 2x - 2 \Rightarrow y_2 = 2x + 2$.

83. Graph $y_1 = 2x - 4$ and $y_2 = 2x + 2$. See Figure 87. The graph y_2 can be obtained by shifting the graph of y_1 upward 6 units. The constant 6, comes from the 6 we added to each y -value in Exercise 84.




Figure 83

84. c ; c ; the same as; c ; upward (or positive vertical)

2.3: Stretching, Shrinking, and Reflecting Graphs

1. The function $y = x^2$ vertically stretched by a factor of 2 is $y = 2x^2$.
2. The function $y = x^3$ vertically shrunk by a factor of one half is $y = \frac{1}{2}x^3$.
3. The function $y = \sqrt{x}$ reflected across the y -axis is $y = \sqrt{-x}$.
4. The function $y = \sqrt[3]{x}$ reflected across the x -axis is $y = -\sqrt[3]{x}$.
5. The function $y = |x|$ vertically stretched by a factor of 3 and reflected across the x -axis is $y = -3|x|$.
6. The function $y = |x|$ vertically shrunk by a factor of one-half and reflected across the y -axis is $y = \frac{1}{3}|-x|$.
7. The function $y = x^3$ vertically shrunk by a factor of 0.25 and reflected across the y -axis is $y = 0.25(-x^3)$ or $y = -0.25x^3$.
8. The function $y = \sqrt{x}$ vertically shrunk by a factor of 0.2 and reflected across the x -axis is $y = -0.2\sqrt{x}$.
9. Graph $y_1 = x$, $y_2 = x + 3$ (y_1 shifted up 3 units), and $y_3 = x - 3$ (y_1 shifted down 3 units). See Figure 9.
10. Graph $y_1 = x^3$, $y_2 = x^3 + 4$ (y_1 shifted up 4 units), and $y_3 = x^3 - 4$ (y_1 shifted down 4 units). See Figure 10.
11. Graph $y_1 = |x|$, $y_2 = |x - 3|$ (y_1 shifted right 3 units), and $y_3 = |x + 3|$ (y_1 shifted left 3 units). See Figure 11.




Figure 9




Figure 10




Figure 11

12. Graph $y_1 = |x|$, $y_2 = |x| - 3$ (y_1 shifted down 3 units), and $y_3 = |x| + 3$ (y_1 shifted up 3 units).

See Figure 12.

13. Graph $y_1 = \sqrt{x}$, $y_2 = \sqrt{x+6}$ (y_1 shifted left 6 units), and $y_3 = \sqrt{x-6}$ (y_1 shifted right 6 units). See Figure 13.

14. Graph $y_1 = |x|$, $y_2 = 2|x|$ (y_1 stretched vertically by a factor of 2), and $y_3 = 2.5|x|$ (y_1 stretched vertically by a factor of 2.5). See Figure 14




Figure 12




Figure 13




Figure 14

15. Graph $y_1 = \sqrt[3]{x}$, $y_2 = -\sqrt[3]{x}$ (y_1 reflected across the x -axis), and $y_3 = -2\sqrt[3]{x}$ (y_1 reflected across the x -axis and stretched vertically by a factor of 2). See Figure 15.

16. Graph $y_1 = x^2$, $y_2 = (x-2)^2 + 1$ (y_1 shifted right 2 units and up 1 unit), and $y_3 = -(x+2)^2$ (y_1 shifted left 2 units and reflected across the x -axis). See Figure 16

17. Graph $y_1 = |x|$, $y_2 = -2|x-1|+1$ (y_1 reflected across the x -axis, stretched vertically by a factor of 2, shifted right 1 unit, and shifted up 1 unit), and $y_3 = -\frac{1}{2}|x|-4$ (y_1 reflected across the x -axis, shrunk by factor of $\frac{1}{2}$, and shifted down 4 units). See Figure 17




Figure 15




Figure 16




Figure 17

18. Graph $y_1 = \sqrt{x}$, $y_2 = -\sqrt{x}$ (y_1 reflected across the x -axis), and $y_3 = \sqrt{-x}$ (y_1 reflected across the y -axis). See Figure 18.
19. Graph $y_1 = x^2 - 1$ (which is $y = x^2$ shifted down 1 unit), $y_2 = \left(\frac{1}{2}x\right)^2 - 1$ (y_1 shrunk vertically by a factor of $\frac{1}{2}$), and $y_3 = (2x)^2 - 1$ (y_1 stretched vertically by a factor of 2^2 or 4). See Figure 19.
20. Graph $y_1 = 3 - |x|$ (which is $y = |x|$ reflected across the x -axis and shifted up 3 units), $y_2 = 3 - |3x|$ stretched vertically by a factor of 3), and $y_3 = 3 - \left|\frac{1}{3}x\right|$ (y_1 shrunk vertically by a factor of $\frac{1}{3}$). See Figure 20




Figure 18




Figure 19




Figure 20

21. Graph $y_1 = \sqrt[3]{x}$, $y_2 = \sqrt[3]{-x}$ (y_1 reflected across the y -axis), and $y_3 = \sqrt[3]{-(x-1)}$ (y_1 reflected across the y -axis and shifted right 1 unit). See Figure 21.
22. Graph $y_1 = \sqrt[3]{x}$, $y_2 = 2 - \sqrt[3]{x}$ (y_1 reflected across the x -axis and shifted up 2 units), and $y_3 = 1 + \sqrt[3]{x}$ (y_1 shifted up 1 unit). See Figure 22.




Figure 21




Figure 22

23. The graph $y = f(x) = x^2$ has been reflected across the x -axis, shifted 5 units to the right, and shifted 2 units downward; therefore, the equation of $g(x)$ is $g(x) = -(x-5)^2 - 2$.
24. The graph $y = f(x) = x^3$ has been shifted 4 units to the right and shifted 3 units upward; therefore, the equation of $g(x)$ is $g(x) = (x-4)^3 + 3$.
25. The graph $y = f(x) = \sqrt{x}$ has been reflected across the y -axis and shifted 1 unit upward; therefore, the equation of $g(x)$ is $g(x) = \sqrt{-x} + 1$.
26. The graph $y = f(x) = \sqrt[3]{x}$ has been reflected across the x -axis and shifted 2 units to the right; therefore, the equation of $g(x)$ is $g(x) = -\sqrt[3]{x-2}$.
27. 4; x
28. 6; x
29. 2; left; $\frac{1}{4}$; x ; 3; downward (or negative)
30. y ; $\frac{2}{5}$; x ; 6 upward (or positive)
31. 3; right; 6
32. 2; left; 0.5
33. The function $y = x^2$ is vertically shrunk by a factor of $\frac{1}{2}$ and shifted 7 units down; therefore, $y = \frac{1}{2}x^2 - 7$.
34. The function $y = x^3$ is vertically stretched by a factor of 3, reflected across the x -axis, and shifted 8 units upward; therefore, $y = 3x + 8$.
35. The function $y = \sqrt{x}$ is shifted 3 units right, vertically stretched by a factor of 4.5, and shifted 6 units down; therefore, $y = 4.5\sqrt{x-3} - 6$.
36. The function $y = \sqrt[3]{x}$ is shifted 2 units left, vertically stretched by a factor of 1.5, and shifted 8 units upward; therefore, $y = 1.5\sqrt[3]{x+2} + 8$.
37. The function $f(x) = \sqrt{x-3} + 2$ is $f(x) = \sqrt{x}$ shifted 3 units right and 2 units upward. See Figure 37.

38. The function $f(x) = |x+2| - 3$ is $f(x) = |x|$ shifted 2 units left and 3 units downward. See Figure 38.

39. The function $f(x) = \sqrt{2x} = \sqrt{2}\sqrt{x}$ is $f(x) = \sqrt{x}$ stretched vertically by a factor of $\sqrt{2}$. See Figure 39.




Figure 37




Figure 38




Figure 39

40. The function $f(x) = \frac{1}{2}(x+2)^2$ is $f(x) = x^2$ shifted 2 units left and shrunk vertically by a factor of $\frac{1}{2}$.

See Figure 40.

41. The function $f(x) = |2x| = 2|x|$ is $f(x) = |x|$ stretched vertically by a factor of 2. See Figure 41.

42. The function $f(x) = \frac{1}{2}|x|$ is $f(x) = |x|$ shrunk vertically by a factor of $\frac{1}{2}$. See Figure 42




Figure 40




Figure 41




Figure 42

43. The function $f(x) = 1 - \sqrt{x}$ is $f(x) = \sqrt{x}$ reflected across the x -axis and shifted 1 unit upward. See Figure 43.

44. The function $f(x) = 2\sqrt{x-2} - 1$ is $f(x) = \sqrt{x}$ shifted 2 units right, stretched vertically by a factor of 2, and shifted 1 unit downward. See Figure 44.

45. The function $f(x) = -\sqrt{1-x} = -\sqrt{-(x-1)}$ is $f(x) = \sqrt{x}$ reflected across both the x -axis, the y -axis and shifted 1 unit right. See Figure 45.




Figure 43




Figure 44




Figure 45

46. The function $f(x) = \sqrt{-x} - 1$ is $f(x) = \sqrt{x}$ reflected across the y -axis and shifted 1 unit downward.

See Figure 46.

47. The function $f(x) = \sqrt{-(x+1)}$ is $f(x) = \sqrt{x}$ reflected across the y -axis and shifted 1 unit left. See Figure 47.

48. The function $f(x) = 2 + \sqrt{-(x-3)}$ is $f(x) = \sqrt{x}$ reflected across the y -axis, shifted 3 units right, and shifted 2 units upward. See Figure 48.




Figure 46




Figure 47




Figure 48

49. The function $f(x) = (x-1)^3$ is $f(x) = x^3$ shifted 1 unit right. See Figure 49.

50. The function $f(x) = (x+2)^3$ is $f(x) = x^3$ shifted 2 unit left. See Figure 50.

51. The function $f(x) = -x^3$ is $f(x) = x^3$ reflected across the x -axis. See Figure 51.




Figure 49




Figure 50




Figure 51

52. The function $f(x) = (-x)^3 + 1$ is $f(x) = x^3$ reflected across the y -axis and shifted 1 unit upward. See Figure 52.




Figure 52




Figure 53




Figure 54

53. The function $f(x) = |-x-2|$ is $f(x) = |x|$ shifted two units to the right and reflected across the y -axis. See Figure 53.

54. The function $f(x) = (-x+1)^2$ is $f(x) = x^2$ shifted one unit to the left and reflected across the y -axis. See Figure 54.

55. The function $f(x) = |-x + 3|$ is $f(x) = |x|$ shifted three units to the left and reflected across the y -axis. See Figure 55.




Figure 55

56. The function $f(x) = |-2x + 1|$ is $f(x) = |x|$ stretched vertically by a factor of 2, shifted one unit to the left and reflected across the y -axis. See Figure 56.




Figure 56

57. (a) The equation $y = -f(x)$ is $y = f(x)$ reflected across the x -axis. See Figure 57a.
 (b) The equation $y = f(-x)$ is $y = f(x)$ reflected across the y -axis. See Figure 57b.
 (c) The equation $y = 2f(x)$ is $y = f(x)$ stretched vertically by a factor of 2. See Figure 57c.
 (d) From the graph $f(0) = 1$.




Figure 57a




Figure 57b




Figure 57c

58. (a) The equation $y = -f(x)$ is $y = f(x)$ reflected across the x -axis. See Figure 58a.
 (b) The equation $y = f(-x)$ is $y = f(x)$ reflected across the y -axis. See Figure 58b.
 (c) The equation $y = 3f(x)$ is $y = f(x)$ stretched vertically by a factor of 3. See Figure 58c.
 (d) From the graph $f(4) = 1$




Figure 58a




Figure 58b




Figure 58c

59. (a) The equation $f(x)$ is $y = f(x)$ reflected across the x -axis. See Figure 59a.
 (b) The equation $y = f(-x)$ is $y = f(x)$ reflected across the y -axis. See Figure 59b.
 (c) The equation $y = f(x+1)$ is $y = f(x)$ shifted 1 unit to the left. See Figure 59c.
 (d) From the graph, there are two x -intercepts, $(-1, 0)$ and $(4, 0)$.




Figure 59a




Figure 59b




Figure 59c

60. (a) The equation $y = -f(x)$ is $y = f(x)$ reflected across the x -axis. See Figure 60a.
 (b) The equation $y = f(-x)$ is $y = f(x)$ reflected across the y -axis. See Figure 60b.
 (c) The equation $y = \frac{1}{2}f(x)$ is $(20, 5)$ shrunk vertically by a factor of $\frac{1}{2}$. See Figure 60c.
 (d) From the graph $f(x) < 0$ for the interval: $(-\infty, 0)$.




Figure 60a




Figure 60b




Figure 60c

61. (a) The equation $y = -f(x)$ is $y = f(x)$ reflected across the x -axis. See Figure 61a.
 (b) The equation $y = f\left(\frac{1}{3}x\right)$ is $y = f(x)$ stretched horizontally by a factor of 3. See Figure 61b.
 (c) The equation $y = 0.5f(x)$ is $y = f(x)$ shrunk vertically by a factor of 0.5. See Figure 61c.
 (d) From the graph, symmetry with respect to the origin.




Figure 61a




Figure 61b




Figure 61c

62. (a) The equation $y = f(2x)$ is $y = f(x)$ stretched horizontally by a factor of $\frac{1}{2}$. See Figure 62a.
- (b) The equation $y = f(-x)$ is $y = f(x)$ reflected across the y-axis. See Figure 62b.
- (c) The equation $y = 3f(x)$ is $y = f(x)$ stretched vertically by a factor of 3. See Figure 62c.
- (d) From the graph, symmetry with respect to the y-axis.




Figure 62a




Figure 62b




Figure 62c

63. (a) The equation $y = f(x) + 1$ is $y = f(x)$ shifted 1 unit upward. See Figure 63a.
- (b) The equation -30° is $y = f(x)$ reflected across the x-axis and shifted 1 unit down. See Figure 63b.
- (c) The equation $y = 2f\left(\frac{1}{2}x\right)$ is $y = f(x)$ stretched vertically by a factor of 2 and horizontally by a factor of 2. See Figure 63c.




Figure 63a




Figure 63b




Figure 63c

64. (a) The equation $y = f(x) - 2$ is $y = f(x)$ shifted 2 units downward. See Figure 64a.
- (b) The equation $y = f(x - 1) + 2$ is $y = f(x)$ shifted 1 unit right and 2 units upward. See Figure 64b.
- (c) The equation $y = 2f(x)$ is $y = f(x)$ stretched vertically by a factor of 2. See Figure 64c.




Figure 64a




Figure 64b




Figure 64c

65. (a) The equation $y = f(2x) + 1$ is $y = f(x)$ shrunk horizontally by a factor of $\{2\}$ and shifted 1 unit upward. See Figure 65a.
- (b) The equation $y = 2f\left(\frac{1}{2}x\right) + 1$ is $y = f(x)$ stretched vertically by a factor of 2, stretched horizontally by a factor of 2, and shifted 1 unit upward. See Figure 65b.
- (c) The equation $y = \frac{1}{2}f(x - 2)$ is $y = f(x)$ shrunk vertically by a factor of $\frac{1}{2}$ and shifted 2 units to the right. See Figure 65c.




Figure 65a




Figure 65b




Figure 65c

66. (a) The equation $y = f(2x)$ is $y = f(x)$ shrunk horizontally by a factor of $\frac{1}{2}$. See Figure 66a.
- (b) The equation $y = f\left(\frac{1}{2}x\right) - 1$ is $y = f(x)$ stretched horizontally by a factor of 2, and shifted 1 unit downward. See Figure 66b.
- (c) The equation $y = 2f(x) - 1$ is $y = f(x)$ stretched vertically by a factor of 2 and shifted 1 unit downward. See Figure 66c.




Figure 66a




Figure 66b




Figure 66c

67. (a) If $(r, 0)$ is the x -intercept of $y = f(x)$ and $y = -f(x)$ is $y = f(x)$ reflected across the x -axis, then $(r, 0)$ is also the x -intercept of $y = -f(x)$.
- (b) If $(r, 0)$ is the x -intercept of $y = f(x)$ and $y = f(-x)$ is $f(x)$ reflected across the y -axis, then $(-r, 0)$ is the x -intercept of $y = -f(x)$.
- (c) If $(r, 0)$ is the x -intercept of $y = f(x)$ and $y = -f(-x)$ is $y = f(x)$ reflected across both the x -axis and y -axis, then $(-r, 0)$ is the x -intercept of $y = -f(-x)$.
68. (a) If $(0, b)$ is the y -intercept of $y = f(x)$ and $y = -f(x)$ is $y = f(x)$ reflected across the x -axis, then $(0, -b)$ is the y -intercept of $y = -f(x)$.
- (b) If $(0, b)$ is the y -intercept of $y = f(x)$ and $y = f(-x)$ is $y = f(x)$ reflected across the y -axis, then $(0, b)$ is also the y -intercept of $y = f(-x)$.
- (c) If $(0, b)$ is the y -intercept of $y = f(x)$ and $y = 5f(x)$ is $y = f(x)$ stretched vertically by a factor of 5, then $(0, 5b)$ is the y -intercept of $y = 5f(x)$.
- (d) If $(0, b)$ is the y -intercept of $y = f(x)$ and $y = -3f(x)$ is $y = f(x)$ reflected across the x -axis and stretched vertically by a factor of 3, then $(0, -3b)$ is the y -intercept of $y = -3f(x)$.
69. Since $y = f(x-2)$ is $y = f(x)$ shifted 2 units to the right, the domain of $f(x-2)$ is $[-1+2, 2+2]$ or $[1, 4]$, and the range is the same: $[0, 3]$.
70. Since $5f(x+1)$ is $f(x)$ shifted 1 unit to the left, the domain of $5f(x+1)$ is $[-1-1, 2-1]$ or $[-2, 1]$ and, stretched vertically by a factor of 5, the range is $[5(0), 5(3)]$ or $[0, 15]$.
71. Since $-f(x)$ is $f(x)$ reflected across the x -axis, the domain of $-f(x)$ is the same: $[-1, 2]$, and the range is $[-3, 0]$.
72. Since $f(x-3)+1$ is $f(x)$ shifted 3 unit to the right, $f(x-3)+1$ is $[-1+3, 2+3]$ or $[2, 5]$, an shifted 1 unit upward, the range is $[0+1, 3+1]$ or $[1, 4]$.

73. Since $f(2x)$ is $f(x)$ shrunk horizontally by a factor of $\frac{1}{2}$, the domain of $f(2x)$ is $\left[\frac{1}{2}(-1), \frac{1}{2}(2)\right]$ or $\left[-\frac{1}{2}, 1\right]$, and the range is the same: $[0, 3]$.
74. Since $2f(x-1)$ is $f(x)$ shifted 1 unit to the right, the domain of $2f(x-1)$ is $[-1+1, 2+1]$ or $[0, 3]$, and stretched vertically by a factor of 3, the range is $[2(-2), 2(3)]$ or $[0, 6]$.
75. Since $3f\left(\frac{1}{4}x\right)$ is $f(x)$ stretched horizontally by a factor of 4, the domain of $3f\left(\frac{1}{4}x\right)$ is $[4(-1), 4(2)]$ or $[-4, 8]$, and stretched vertically by a factor of 3, the range is $[3(0), 3(3)]$ or $[0, 9]$.
76. Since $-2f(4x)$ is $f(x)$ shrunk horizontally by a factor of $\frac{1}{4}$, the domain of $2f(4x)$ is $\left[\frac{1}{4}(-1), \frac{1}{4}(2)\right]$ or $\left[-\frac{1}{4}, \frac{1}{2}\right]$; and reflected across the x -axis while being stretched vertically by a factor of 2, the range is $[-2(0), -2(3)] = [0, -6]$ or $[-6, 0]$.
77. Since $f(-x)$ is $f(x)$ reflected across the y -axis, the domain of $f(-x)$ is $[-(-1), -(2)] = [1, -2]$ or $[-2, 1]$; and the range is the same: $[0, 3]$.
78. Since $-2f(-x)$ is $f(x)$ reflected across the y -axis, the domain of $-2f(-x)$ is $[-(-1), -(2)] = [1, -2]$ or $[-2, 1]$, and reflected across the x -axis while being stretched vertically by a factor of 2, the range is $[-2(0), -2(3)] = [0, -6]$ or $[-6, 0]$.
79. Since $f(-3x)$ is $f(x)$ reflected across the y -axis and shrunk horizontally by a factor of $\frac{1}{3}$, the domain of $f(-3x)$ is $\left[-\frac{1}{3}(-1), -\frac{1}{3}(2)\right] = \left[\frac{1}{3}, -\frac{2}{3}\right]$ or $\left[-\frac{2}{3}, \frac{1}{3}\right]$, and the range is the same: $[0, 3]$.
80. Since $\frac{1}{3}f(x-3)$ is $f(x)$ shifted 3 units to the right, the domain of $\frac{1}{3}f(x-3)$ is $[-1+3, 2+3]$ or $[2, 5]$, and shrunk vertically by a factor of $\frac{1}{3}$, the range is $\left[\frac{1}{3}(0), \frac{1}{3}(3)\right]$ or $[0, 1]$.
81. Since $y = \sqrt{x}$ has an endpoint of $(0, 0)$, and the graph of $y = 10\sqrt{x-20} + 5$ is the graph of $y = \sqrt{x}$ shifted 20 units right, stretched vertically by a factor of 10, and shifted 5 units upward, the endpoint of $y = 10\sqrt{x-20} + 5$ is $(0+20, 10(0)+5)$ or $(20, 5)$. Therefore, the domain is $[20, \infty)$, and the range is $[5, \infty)$.

82. Since $y = \sqrt{x}$ has an endpoint of $(0, 0)$, and the graph of $y = -2\sqrt{x+15} - 18$ is the graph of $y = \sqrt{x}$ shifted 15 units left, reflected across the x -axis, stretched vertically by a factor of 2, and shifted 18 units downward, the endpoint of $y = -2\sqrt{x+15} - 18$ is $(0-15, -2(0)-18)$ or $(-15, -18)$. Therefore, the domain is $[-15, \infty)$, and the range, because of the reflection across the x -axis, is $(-\infty, -18]$.
83. Since $y = \sqrt{x}$ has an endpoint of $(0, 0)$, and the graph of $y = -.5\sqrt{x+10} + 5$ is the graph of $y = \sqrt{x}$ shifted 10 units left, reflected across the x -axis, shrunk vertically by a factor of .5, and shifted 5 units upward, the endpoint of $y = -.5\sqrt{x+10} + 5$ is $(0-10, -.5(0)+5)$ or $(-10, 5)$. Therefore, the domain is $[-10, \infty)$, and the range, because of the reflection across the x -axis, is $(-\infty, 5]$.
84. Using ex. 81, the domain is $[h, \infty)$, and the range is $[k, \infty)$.
85. The graph of $y = -f(x)$ is $y = f(x)$ reflected across the x -axis; therefore, $y = -f(x)$ is decreasing for the interval (a, b) .
86. The graph of $y = f(-x)$ is $y = f(x)$ reflected across the y -axis; therefore, $y = f(-x)$ is decreasing for the interval $(-b, -a)$.
87. The graph of $y = -f(-x)$ is $y = f(x)$ reflected across both the x -axis and y -axis; therefore, $y = -f(-x)$ is increasing for the interval $(-b, -a)$.
88. The graph of $y = -c \cdot f(x)$ is $y = f(x)$ reflected across the x -axis; therefore, $y = -c \cdot f(x)$ is decreasing for the interval (a, b) .
89. (a) the function is increasing for the interval: $(-1, 2)$.
(b) the function is decreasing for the interval: $(-\infty, -1)$.
(c) the function is constant for the interval: $(2, \infty)$.
90. (a) the function is increasing for the interval: $(-\infty, -1)$.
(b) the function is decreasing for the interval: $(-1, 2)$.
(c) the function is constant for the interval: $(2, \infty)$.
91. (a) the function is increasing for the interval: $(1, \infty)$.
(b) the function is decreasing for the interval: $(-2, 1)$.
(c) the function is constant for the interval: $(-\infty, -2)$.
92. (a) the function is increasing for the interval: $(-\infty, -3)$.
(b) the function is decreasing for the interval: $(-3, \infty)$.
(c) the function is not constant for any interval.

93. From the graph, the point on y_2 is approximately $(8,10)$.

94. From the graph, the point on y_2 is approximately $(-27,-15)$.

95. Use two points on the graph to find the slope. Two points are $(-2,-1)$ and $(-1,1)$; therefore, the slope is

$$m = \frac{1 - (-1)}{-1 - (-2)} = \frac{2}{1} \Rightarrow m = 2. \text{ The stretch factor is 2 and the graph has been shifted 2 units to the left and 1}$$

unit down; therefore, the equation is $y = 2|x + 2| - 1$.

96. Use two points on the graph to find the slope. Two points are $(1,2)$ and $(5,0)$; therefore, the slope is

$$m = \frac{0 - 2}{5 - 1} = \frac{-2}{4} \Rightarrow m = -\frac{1}{2}. \text{ The shrinking factor is } \frac{1}{2}, \text{ the graph has been reflected across the } x\text{-axis,}$$

shifted 1 unit to the right, and shifted 2 units upward; therefore, the equation is $y = -\frac{1}{2}|x - 1| + 2$.

97. Use two points on the graph to find the slope. Two points are $(0,2)$ and $(1,-1)$; therefore, the slope is

$$m = \frac{-1 - 2}{1 - 0} = \frac{-3}{1} \Rightarrow m = -3. \text{ The stretch factor is 3, the graph has been reflected across the } x\text{-axis, and}$$

shifted 2 units upward; therefore, the equation is $y = -3|x| + 2$.

98. Use two points on the graph to find the slope. Two points are $(-1,-2)$ and $(0,1)$; therefore, the slope is

$$m = \frac{1 - (-2)}{0 - (-1)} = \frac{3}{1} \Rightarrow m = 3. \text{ The stretch factor is 3 and the graph has been shifted 1 unit to the left and 2}$$

units down; therefore, the equation is $y = 3|x + 1| - 2$.

99. Use two points on the graph to find the slope. Two points are $(0,-4)$ and $(3,0)$; therefore, the slope is

$$m = \frac{-4 - 0}{0 - 3} = \frac{4}{3} \Rightarrow m = \frac{4}{3}. \text{ The stretch factor is } \frac{4}{3} \text{ and the graph has been shifted 4 units down.; therefore,}$$

the equation is $y = \frac{4}{3}|x| - 4$.

100. Use two points on the graph to find the slope. Two points are $(0,4)$ and $(-3,5)$; therefore, the slope is

$$m = \frac{5 - 4}{-3 - 0} = -\frac{1}{3} \Rightarrow m = -\frac{1}{3}. \text{ The stretch factor is } -\frac{1}{3} \text{ and the graph has been shifted 3 units to the left and}$$

5 units up.; therefore, the equation is $y = -\frac{1}{3}|x + 3| + 5$.

101. Since $y = f(x)$ is symmetric with respect to the y -axis, for every (x,y) on the graph, $(-x,y)$ is also on the graph. Reflection across the y -axis reflect onto itself and will not change the graph. It will be the same.

102. The graph is no longer symmetric to the origin since it has been moved away from the origin.

Reviewing Basic Concepts (Sections 2.1—2.3)

1. (a) The function $f(x) = |x|$ shifted up one unit yields the function $f(x) = |x| + 1$. Therefore, this function has a domain of $(-\infty, \infty)$ and a range of $[1, \infty)$. The function is increasing from $(0, \infty)$ and decreasing from $(-\infty, 0)$.
- (b) The function $f(x) = x^2$ shifted to the right 2 units yields the function $f(x) = (x - 2)^2$. Therefore, this function has a domain of $(-\infty, \infty)$ and a range of $[0, \infty)$. The function is increasing from $(2, \infty)$ and decreasing from $(-\infty, 2)$.
- (c) The function $f(x) = \sqrt{x}$ reflected over the x -axis yields the function $f(x) = -\sqrt{x}$. Therefore, this function has a domain of $[0, \infty)$ and a range of $(-\infty, 0]$. The function is never increasing and decreasing from $(0, \infty)$.
2. (a) If $y = f(x)$ is symmetric with respect to the origin, then another function value is $f(-3) = -6$.
- (b) If $y = f(x)$ is symmetric with respect to the y -axis, then another function value is $f(-3) = 6$.
- (c) If $f(-x) = -f(x)$, $y = f(x)$ is symmetric with respect to both the x -axis and y -axis, then another function value is $f(-3) = -6$.
- (d) If $y = f(-x)$, $y = f(x)$ is symmetric with respect to the y -axis, then another function value is $f(-3) = 6$.
3. (a) The equation $y = (x - 7)^2$ is $y = x^2$ shifted 7 units to the right: B.
- (b) The equation $y = x^2 - 7$ is $y = x^2$ shifted 7 units downward: D.
- (c) The equation $y = 7x^2$ is $y = x^2$ stretches vertically by a factor of 7: E.
- (d) The equation $y = (x + 7)^2$ is $y = x^2$ shifted 7 units to the left: A.
- (e) The equation $y = \left(\frac{1}{3}x\right)^2$ is $y = x^2$ stretches horizontally by a factor of 3: C.
4. (a) The equation $y = x^2 + 2$ is $y = x^2$ shifted 2 units upward: B.
- (b) The equation $y = x^2 - 2$ is $y = x^2$ shifted 2 units downward: A.
- (c) The equation $y = (x + 2)^2$ is $y = x^2$ shifted 2 units to the left: G.
- (d) The equation $y = (x - 2)^2$ is $y = x^2$ shifted 2 units to the right: C.
- (e) The equation $y = 2x^2$ is $y = x^2$ stretched vertically by a factor of 2: F.
- (f) The equation $y = -x^2$ is $y = x^2$ reflected across the x -axis D.
- (g) The equation $y = (x - 2)^2 + 1$ is $y = x^2$ shifted 2 units to the right and 1 unit upward: H.
- (h) The equation $y = (x + 2)^2 + 1$ is $y = x^2$ shifted 2 units to the left and 1 unit upward: E.
5. (a) The equation $y = |x| + 4$ is $y = |x|$ shifted 4 units upward. See Figure 5a.
- (b) The equation $y = |x + 4|$ is $y = |x|$ shifted 4 units to the left. See Figure 5b.

- (c) The equation $y = |x - 4|$ is $y = |x|$ shifted 4 units to the right. See Figure 5c.
- (d) The equation $y = |x + 2| - 4$ is $y = |x|$ shifted 2 units to the left and 4 units down. See Figure 5d.
- (e) The equation $y = -|x - 2| + 4$ is $y = |x|$ reflected across the x -axis, shifted 2 units to the right, and 4 units upward. See Figure 5e.




Figure 5a




Figure 5b




Figure 5c




Figure 5d




Figure 5e

6. (a) The graph is the function $f(x) = |x|$ reflected across the x -axis, shifted 1 unit left and 3 units upward. Therefore, the equation is $y = -|x + 1| + 3$.
- (b) The graph is the function $g(x) = \sqrt{x}$ reflected across the x -axis, shifted 4 units left and 2 units upward. Therefore, the equation is $y = -\sqrt{x + 4} + 2$.
- (c) The graph is the function $g(x) = \sqrt{x}$ stretches vertically by a factor of 2, shifted 4 units left and 4 units downward. Therefore, the equation is $y = 2\sqrt{x + 4} - 4$.
- (d) The graph is the function $f(x) = |x|$ shrunk vertically by a factor of $\frac{1}{2}$, shifted 2 units right and 1 unit downward. Therefore, the equation is $y = \frac{1}{2}|x - 2| - 1$.
7. (a) The graph of $g(x)$ is the graph $f(x)$ shifted 2 units upward. Therefore, $c = 2$.
- (b) The graph of $g(x)$ is the graph $f(x)$ shifted 4 units to the left. Therefore, $c = 4$.
8. The graph of $y = F(x + h)$ is a horizontal translation of the graph of $y = F(x)$. The graph of $y = F(x) + h$ is not the same as the graph of $y = F(x + h)$, because the graph of $y = F(x) + h$ is a vertical translation of the graph of $y = F(x)$.
9. (a) If f is even, then $f(x) = f(-x)$. See Figure 9a.
- (b) If f is odd, then $f(-x) = -f(x)$. See Figure 9b.

x	$f(x)$
-3	4
-2	-6
-1	5
1	5
2	-6
3	4

Figure 9a

x	$f(x)$
-3	4
-2	-6
-1	5
1	-5
2	6
3	-4

Figure 9b

10. (a) $R(7) = 8.25(7) + 27.5 = 85.25$, In 2017, Google's ad revenues were \$85.25 billion.
- (b) Using the point (2010, 27.5) and the slope of 8.255 with the point slope formula we will have
 $y - 27.5 = 8.25(x - 2010) \Rightarrow y = 8.25(x - 2010) + 27.5$.
- (c) $y = 8.25(2017 - 2010) + 27.5 = 8.25(7) + 27.5 = 85.25$, In 2017, Google's ad revenues were \$85.25 billion.
- (d) $69 = 8.25(x - 2010) + 27.5 \Rightarrow 41.5 = 8.25(x - 2010) \Rightarrow \frac{41.5}{8.25} + 2010 = x \Rightarrow x \approx 2015$
11. (a) $f(x) = 0$: $\{3, 4\}$
- (b) $f(x) > 0$: for the intervals $(-\infty, 3) \cup (4, \infty)$.
- (c) $f(x) < 0$: for the interval $(3, 4)$.
12. (a) $f(x) = 0$: $\{\sqrt{2}\}$
- (b) $f(x) > 0$: for the interval $(\sqrt{2}, \infty)$.
- (c) $f(x) < 0$: for the interval $(-\infty, \sqrt{2})$.
13. (a) $f(x) = 0$: $\{-4, 5\}$
- (b) $f(x) \geq 0$: for the intervals $(-\infty, -4] \cup [5, \infty)$
- (c) $f(x) \leq 0$: for the interval $[-4, 5]$.
14. (a) $f(x) = 0$: never; therefore: $\emptyset$.
- (b) $f(x) \geq 0$: for the interval $[1, \infty)$.
- (c) $f(x) \leq 0$: never; therefore: $\emptyset$.

2.4: Absolute Value Functions

- We reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 1. The domain and range of $f(x)$ is $(-\infty, \infty)$ and $[-1, \infty)$ respectively. The domain and range of $|f(x)|$ is $(-\infty, \infty)$ and $[0, \infty)$ respectively.
- We reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 2. The domain and range of $f(x)$ is $(-\infty, \infty)$ and $(-\infty, 1]$ respectively. The domain and range of $|f(x)|$ is $(-\infty, \infty)$ and $[0, \infty)$ respectively.

3. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 3. The domain and range of $f(x)$ is $(-\infty, \infty)$ and $(-\infty, \infty)$ respectively. The domain and range of $|f(x)|$ is $(-\infty, \infty)$ and $[0, \infty)$ respectively.




Figure 1




Figure 2




Figure 3

4. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 4. The domain and range of $f(x)$ is $(-\infty, \infty)$ and $(-\infty, \infty)$ respectively. The domain and range of $|f(x)|$ is $(-\infty, \infty)$ and $(0, \infty)$ respectively.
5. Since for all y , $y \geq 0$, the graph remains unchanged. That is, $y = |f(x)|$ has the same graph as $y = f(x)$. The domain and range of $f(x)$ is $[0, \infty)$ and $[0, \infty)$ respectively. The domain and range of $|f(x)|$ is $[0, \infty)$ and $[0, \infty)$ respectively.
6. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 6. The domain and range of $f(x)$ is $[0, \infty)$ and $[0, \infty)$ respectively. The domain and range of $|f(x)|$ is $[0, \infty)$ and $[0, \infty)$ respectively.




Figure 4




Figure 6

7. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 7. The domain and range of $f(x)$ is $(-\infty, 0) \cup (0, \infty)$ and $(-\infty, 0) \cup (0, \infty)$ respectively. The domain and range of $|f(x)|$ is $(-\infty, 0) \cup (0, \infty)$ and $(0, \infty)$ respectively.
8. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 8. The domain and range of $f(x)$ is $[0, \infty)$ and $[-1, \infty)$ respectively. The domain and range of $|f(x)|$ is $[0, \infty)$ and $[0, \infty)$ respectively.

9. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 9. The domain and range of $f(x)$ is $[-2, \infty)$ and $[-1, \infty)$ respectively. The domain and range of $|f(x)|$ is $[-2, \infty)$ and $[0, \infty)$ respectively.




Figure 7




Figure 8




Figure 9

10. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged.
11. If $f(a) = -5$, then $|f(a)| = |-5| = 5$.
12. Since $f(x) = x^2$ is an even function, $f(x) = x^2$ and $f(x) = |x^2|$ are the same graph.
13. If $f(x) = -x^2$, then $y = |f(x)| \Rightarrow y = |-x^2| \Rightarrow y = x^2$. Therefore, the range of $y = |f(x)|$ is $[0, \infty)$.
14. If the range of $y = f(x)$ is $[-2, \infty)$, the range of $y = |f(x)|$ is $[0, \infty)$ since all negative values of y are reflected across the x -axis.
15. If the range of $y = f(x)$ is $(-\infty, -2]$, the range of $y = |f(x)|$ is $[2, \infty)$ since all negative values of y are reflected across the x -axis.
16. $|f(x)|$ is greater than or equal to 0 for any value of x . Since -1 is less than 0, -1 cannot be in the range of f .
17. From the graph of $y = (x+1)^2 - 2$ the domain of $f(x)$ is $(-\infty, \infty)$, and the range is $[-2, \infty)$.
From the graph of $y = |(x+1)^2 - 2|$ the domain of $|f(x)|$ is $(-\infty, \infty)$, and the range is $[0, \infty)$.
18. From the graph of $y = 2 - \frac{1}{2}x$ the domain of $f(x)$ is $(-\infty, \infty)$, and the range is $(-\infty, \infty)$. From the graph of $y = \left|2 - \frac{1}{2}x\right|$ the domain of $|f(x)|$ is $(-\infty, \infty)$, and the range is $[0, \infty)$.
19. From the graph of $y = -1 - (x-2)^2$ the domain of $f(x)$ is $(-\infty, \infty)$, and the range is $(-\infty, -1]$. From the graph of $y = \left|-1 - (x-2)^2\right|$ the domain of $|f(x)|$ is $(-\infty, \infty)$, and the range is $[1, \infty)$.
20. From the graph of $y = -|x+2| - 2$ the domain of $f(x)$ is $(-\infty, \infty)$, and the range is $(-\infty, -2]$. From the graph of $y = \left|-|x+2| - 2\right|$ the domain of $|f(x)|$ is $(-\infty, \infty)$, and the range is $[2, \infty)$.

21. From the graph, the domain of $f(x)$ is $[-2, 3]$, and the range is $[-2, 3]$. For the function $y = |f(x)|$, we reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$, and, where $y \geq 0$, the graph remains unchanged. Therefore, the domain of $y = |f(x)|$ is $[-2, 3]$, and the range is $[0, 3]$.
22. From the graph, the domain of $f(x)$ is $[-3, 2]$, and the range is $[-2, 2]$. For the function $y = |f(x)|$, we reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$ and, where $y \geq 0$, the graph remains unchanged. Therefore, the domain of $y = |f(x)|$ is $[-3, 2]$, and the range is $[0, 2]$.
23. From the graph, the domain of $f(x)$ is $[-2, 3]$, and the range is $[-3, 1]$. For the function $y = |f(x)|$, we reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$, and, where $y \geq 0$, the graph remains unchanged. Therefore, the domain of $y = |f(x)|$ is $[-2, 3]$, and the range is $[0, 3]$.
24. From the graph, the domain of $y = f(x)$ is $[-3, 3]$, and the range is $[-3, -1]$. For the function $y = |f(x)|$, we reflect the graph of $y = f(x)$ across the x-axis for all points for which $y < 0$, and, where $y \geq 0$, the graph remains unchanged. Therefore, the domain of $y = |f(x)|$ is $[-3, 3]$, and the range is $[1, 3]$.
25. (a) The function $y = f(-x)$ is the function $y = f(x)$ reflected across the y-axis. See Figure 25a.
- (b) The function $y = -f(-x)$ is the function $y = f(x)$ reflected across both the x-axis and y-axis. See Figure 25b.
- (c) For the function $y = |-f(-x)|$ we reflect the graph of $y = -f(-x)$ (ex. b) across the x-axis for all points for which $y < 0$, and where $y \geq 0$, the graph remains unchanged. See Figure 25c.




Figure 25a




Figure 25b




Figure 25c

26. (a) The function $y = f(-x)$ is the function $y = f(x)$ reflected across the y-axis. See Figure 26a.
- (b) The function $y = -f(-x)$ is the function $y = f(x)$ reflected across both the x-axis and y-axis. See Figure 26b.
- (c) For the function $y = |-f(-x)|$ we reflect the graph of $y = -f(-x)$ (ex. b) across the x-axis for all points for which $y < 0$, and, where $y \geq 0$, the graph remains unchanged. See Figure 26c.




Figure 26a




Figure 26b




Figure 26c

27. The graph of $y = |f(x)|$ can not be below the x-axis; therefore, Figure A shows the graph of $y = f(x)$, while Figure B shows the graph of $y = |f(x)|$.
28. The graph of $y = |f(x)|$ can not be below the x-axis; therefore, Figure B shows the graph of $y = f(x)$, while Figure A shows the graph of $y = |f(x)|$.
29. See Figure 29a and 29b
30. See Figure 30a and 30b.
31. See Figure 31a and 31b.




Figure 29a




Figure 29b




Figure 30a




Figure 30b




Figure 31a




Figure 31b

32. See Figure 32a and 32b.
33. See Figure 33a and 33b.
34. See Figure 34a and 34b.




Figure 32a




Figure 32b




Figure 33a




Figure 33b




Figure 34a




Figure 34b

35. See Figure 35a and 35b.

36. See Figure 36a and 36b.

37. See Figure 37a and 37b.




Figure 35a




Figure 35b




Figure 36a




Figure 36b




Figure 37a




Figure 37b

38. See Figure 38a and 38b.

39. See Figure 39a and 39b.

40. See Figure 40a and 40b.




Figure 38a




Figure 38b




Figure 39a




Figure 39b




Figure 40a




Figure 40b

41. (a) From the graph, $y_1 = y_2$ at the coordinates $(-1, 5)$ and $(6, 5)$; therefore, the solution set is $\{-1, 6\}$.
 (b) From the graph, $y_1 < y_2$ for the interval $(-1, 6)$.
 (c) From the graph, $y_1 > y_2$ for the intervals $(-\infty, -1) \cup (6, \infty)$.
42. (a) From the graph, $y_1 = y_2$ at the coordinates $(0, -2)$ and $(8, -2)$; therefore, the solution set is $\{0, 8\}$.
 (b) From the graph, $y_1 < y_2$ for the intervals $(-\infty, 0) \cup (8, \infty)$.
 (c) From the graph, $y_1 > y_2$ for the interval $(0, 8)$.
43. (a) From the graph, $y_1 = y_2$ at the coordinate $(4, 1)$; therefore, the solution set is $\{4\}$.
 (b) From the graph, $y_1 < y_2$ never occurs; therefore, the solution set is the null set. .
 (c) From the graph, $y_1 > y_2$ for all values for x except 4; therefore, the solution set is the intervals $(-\infty, 4) \cup (4, \infty)$.
44. (a) From the graph, $y_1 = y_2$ never occurs; therefore, the solution set is $\emptyset$.
 (b) From the graph, $y_1 < y_2$ for all values for x ; therefore, the solution set is $(-\infty, \infty)$.
 (c) From the graph, $y_1 > y_2$ never occurs; therefore, the solution set is $\emptyset$.
45. The V-shaped graph is that of $f(x) = |.5x + 6|$, since this is typical of the graphs of absolute value functions of the form $f(x) = |ax + b|$.
46. The straight line graph is that of $g(x) = 3x - 14$ which is a linear function.
47. The graphs intersect at $(8, 10)$, so the solution set is $\{8\}$.

48. From the graph, $f(x) > g(x)$ for the interval $(-\infty, 8)$.
49. From the graph, $f(x) < g(x)$ for the interval $(8, \infty)$.
50. If $|.5x + 6| - (3x - 14) = 0$ then $|.5x + 6| = 3x - 14$. Therefore, the solution is the intersection of the graphs, or $\{8\}$.
51. (a) $|x + 4| = 9 \Rightarrow x + 4 = 9$ or $x + 4 = -9 \Rightarrow x = 5$ or $x = -13$. The solution set is $\{-13, 5\}$, which is supported by the graphs of $y_1 = |x + 4|$ and $y_2 = 9$.
- (b) $|x + 4| > 9 \Rightarrow x + 4 > 9$ or $x + 4 < -9 \Rightarrow x > 5$ or $x < -13$. The solution is $(-\infty, -13) \cup (5, \infty)$, which is supported by the graphs of $y_1 = |x + 4|$ and $y_2 = 9$.
- (c) $|x + 4| < 9 \Rightarrow -9 < x + 4 < 9 \Rightarrow -13 < x < 5$. The solution is $(-13, 5)$, which is supported by the graphs of $|x + 4|$ and $y_2 = 9$.
52. (a) $|x - 3| = 5 \Rightarrow x - 3 = 5$ or $x - 3 = -5 \Rightarrow x = 8$ or $x = -2$. The solution set is $\{-2, 8\}$, which is supported by the graphs of $y_1 = |x - 3|$ and $y_2 = 5$.
- (b) $|x - 3| > 5 \Rightarrow x - 3 > 5$ or $x - 3 < -5 \Rightarrow x > 8$ or $x < -2$. The solution is $(-\infty, -2) \cup (8, \infty)$ which is supported by the graphs of $y_1 = |x - 3|$ and $y_2 = 5$.
- (c) $|x - 3| < 5 \Rightarrow -5 < x - 3 < 5 \Rightarrow -2 < x < 8$. The solution is $(-2, 8)$, which is supported by the graphs of $y_1 = |x - 3|$ and $y_2 = 5$.
53. (a) $|7 - 2x| = 3 \Rightarrow 7 - 2x = 3$ or $7 - 2x = -3 \Rightarrow -2x = -4$ or $-2x = -10 \Rightarrow x = 2$ or $x = 5$. The solution set is $\{2, 5\}$, which is supported by the graphs of $y_1 = |7 - 2x|$ and $y_2 = 3$.
- (b) $|7 - 2x| \geq 3 \Rightarrow 7 - 2x \geq 3$ or $7 - 2x \leq -3 \Rightarrow -2x \geq -4$ or $-2x \leq -10 \Rightarrow x \leq 2$ or $x \geq 5$. The solution set is $(-\infty, 2] \cup [5, \infty)$, which is supported by the graphs of $y_1 = |7 - 2x|$ and $y_2 = 3$.
- (c) $|7 - 2x| \leq 3 \Rightarrow -3 \leq 7 - 2x \leq 3 \Rightarrow -10 \leq -2x \leq -4 \Rightarrow 5 \geq x \geq 2$ or $2 \leq x \leq 5$. The solution is $[2, 5]$, which is supported by the graphs of $y_1 = |7 - 2x|$ and $y_2 = 3$.
54. (a) $|-9 - 3x| = 6 \Rightarrow -9 - 3x = 6$ or $-9 - 3x = -6 \Rightarrow -3x = 15$ or $-3x = 3 \Rightarrow x = -5$ or $x = -1$. The solution set is $\{-5, -1\}$, which is supported by the graphs of $y_1 = |-9 - 3x|$ and $y_2 = 6$.
- (b) $|-9 - 3x| \geq 6 \Rightarrow -9 - 3x \geq 6$ or $-9 - 3x \leq -6 \Rightarrow -3x \geq 15$ or $-3x \leq 3 \Rightarrow x \leq -5$ or $x \geq -1$. The solution is $(-\infty, -5] \cup [-1, \infty)$, which is supported by the graphs of $y_1 = |-9 - 3x|$ and $y_2 = 6$.
- (c) $|-9 - 3x| \leq 6 \Rightarrow -6 \leq -9 - 3x \leq 6 \Rightarrow 3 \leq -3x \leq 15 \Rightarrow -1 \geq x \geq -5$ or $-5 \leq x \leq -1$. The solution is $[-5, -1]$, which is supported by the graphs of $y_1 = |-9 - 3x|$ and $y_2 = 6$.

55. (a) $|2x+1|+3=5 \Rightarrow 2x+1=2$ or $2x+1=-2 \Rightarrow 2x=1$ or $2x=-3 \Rightarrow x=\frac{1}{2}$ or $x=-\frac{3}{2}$. The solution set is $\left\{-\frac{3}{2}, \frac{1}{2}\right\}$, which is supported by the graphs of $y_1=|2x+1|+3$ and $y_2=5$.
- (b) $|2x+1|+3 \leq 5 \Rightarrow -2 \leq 2x+1 \leq 2 \Rightarrow -3 \leq 2x \leq 1 \Rightarrow -\frac{3}{2} \leq x \leq \frac{1}{2}$. The solution is $\left[-\frac{3}{2}, \frac{1}{2}\right]$, which is supported by the graphs of $y_1=|2x+1|+3$ and $y_2=5$.
- (c) $|2x+1|+3 \geq 5 \Rightarrow 2x+1 \geq 2$ or $2x+1 \leq -2 \Rightarrow 2x \geq 1$ or $2x \leq -3 \Rightarrow x \geq \frac{1}{2}$ or $x \leq -\frac{3}{2}$. The solution is $\left(-\infty, -\frac{3}{2}\right] \cup \left[\frac{1}{2}, \infty\right)$, which is supported by the graphs of $y_1=|2x+1|+3$ and $y_2=5$.
56. (a) $|4x+7|+4=4 \Rightarrow 4x+7=0 \Rightarrow 4x=-7 \Rightarrow x=-\frac{7}{4}$. The solution set is $\left\{-\frac{7}{4}\right\}$, which is supported by the graphs of $y_1=|4x+7|+4$ and $y_2=4$.
- (b) $|4x+7|+4 > 4 \Rightarrow 4x+7 > 0$ or $4x+7 < 0 \Rightarrow 4x > -7$ or $4x < -7 \Rightarrow x > -\frac{7}{4}$ or $x < -\frac{7}{4}$. The solution is $\left(-\infty, -\frac{7}{4}\right) \cup \left(-\frac{7}{4}, \infty\right)$, which is supported by the graphs of $y_1=|4x+7|+4$ and $y_2=4$.
- (c) $|4x+7|+4 < 4 \Rightarrow 0 < 4x+7 < 0 \Rightarrow -7 < 4x < -7 \Rightarrow -\frac{7}{4} < x < -\frac{7}{4} \Rightarrow$ the solution set is $\emptyset$, which is supported by the graphs of $y_1=|4x+7|+4$ and $y_2=4$.
57. (a) $|5-7x|=0 \Rightarrow 5-7x=0 \Rightarrow 7x=5 \Rightarrow x=\frac{5}{7}$. The solution set is $\left\{\frac{5}{7}\right\}$, which is supported by the graphs of $y_1=|5-7x|$ and $y_2=0$.
- (b) $|5-7x| \geq 0 \Rightarrow 5-7x \geq 0$ or $5-7x \leq 0 \Rightarrow 7x \geq 5$ or $7x \leq 5 \Rightarrow x \geq \frac{5}{7}$ or $x \leq \frac{5}{7}$. The solution is $(-\infty, \infty)$, which is supported by the graphs of $y_1=|5-7x|$ and $y_2=0$.
- (c) $|5-7x| \leq 0 \Rightarrow 0 \leq 5-7x \leq 0 \Rightarrow 5 \geq 7x \geq 5 \Rightarrow \frac{5}{7} \geq x \geq \frac{5}{7}$. The solution set is $\left\{\frac{5}{7}\right\}$, which is supported by the graphs of $y_1=|5-7x|$ and $y_2=0$.
58. (a) Absolute value is always positive; therefore, the solution set is $\emptyset$, which is supported by the graphs of $y_1=|\pi x+8|$ and $y_2=-4$.
- (b) Absolute value is always positive, and so cannot be less than -4 ; therefore, the solution set is $\emptyset$, which is supported by the graphs of $y_1=|\pi x+8|$ and $y_2=-4$.

- (c) Absolute value is always positive, and so is always greater than -4 ; therefore, the solution set is $(-\infty, \infty)$, which is supported by the graphs of $y_1 = |\pi x + 8|$ and $y_2 = -4$.
59. (a) Absolute value is always positive; therefore, the solution set is $\emptyset$, which is supported by the graphs of $y_1 = |\sqrt{2x} - 3.6|$ and $y_2 = -1$.
- (b) Absolute value is always positive, and so cannot be less than or equal to -1 ; therefore, the solution set is $\emptyset$, which is supported by the graphs of $y_1 = |\sqrt{2x} - 3.6|$ and $y_2 = -1$.
- (c) Absolute value is always positive, and so is always greater than -1 ; therefore, the solution is $(-\infty, \infty)$, which is supported by the graphs of $y_1 = |\sqrt{2x} - 3.6|$ and $y_2 = -1$.
60. $|2x + 4| + 2 = 10 \Rightarrow |2x + 4| = 8 \Rightarrow 2x + 4 = 8$ or $2x + 4 = -8 \Rightarrow 2x = 4$ or $2x = -12 \Rightarrow x = 2$ and $x = -6$. Therefore, the solution set is $\{-6, 2\}$.
61. $3|4 - 3x| - 4 = 8 \Rightarrow 3|4 - 3x| = 12 \Rightarrow |4 - 3x| = 4 \Rightarrow 4 - 3x = 4$ or $4 - 3x = -4 \Rightarrow -3x = 0$ or $-3x = -8 \Rightarrow x = 0$ or $x = \frac{8}{3}$. Therefore, the solution set is $\left\{0, \frac{8}{3}\right\}$.
62. $5|x + 3| - 2 = 18 \Rightarrow 5|x + 3| = 20 \Rightarrow |x + 3| = 4 \Rightarrow x + 3 = 4$ or $x + 3 = -4 \Rightarrow x = 1$ or $x = -7$. Therefore, the solution set is $\{-7, 1\}$.
63. $\frac{1}{2}|-2x + \frac{1}{2}| = \frac{3}{4} \Rightarrow |-2x + \frac{1}{2}| = \frac{3}{2} \Rightarrow -2x + \frac{1}{2} = \frac{3}{2}$ or $-2x + \frac{1}{2} = -\frac{3}{2} \Rightarrow -2x = 1$ or $-2x = -2 \Rightarrow x = -\frac{1}{2}$ or $x = 1$. Therefore, the solution set is $\left\{-\frac{1}{2}, 1\right\}$.
64. $|3(x - 5) + 2| + 3 = 9 \Rightarrow |3(x - 5) + 2| = 6 \Rightarrow 3(x - 5) + 2 = 6$ or $3(x - 5) + 2 = -6 \Rightarrow 3(x - 5) = 4$ or $3(x - 5) = -8 \Rightarrow x - 5 = \frac{4}{3}$ or $x - 5 = -\frac{8}{3} \Rightarrow x = \frac{19}{3}$ or $x = \frac{7}{3}$. Therefore, the solution set is $\left\{\frac{7}{3}, \frac{19}{3}\right\}$.
65. $4.2|.5 - x| + 1 = 3.1 \Rightarrow 4.2|.5 - x| = 2.1 \Rightarrow |.5 - x| = .5 \Rightarrow .5 - x = .5$ or $.5 - x = -.5 \Rightarrow -x = 0$ or $-x = -1 \Rightarrow x = 0$ or $x = 1$. Therefore, the solution set is $\{0, 1\}$.
66. $|3x - 1| < 8 \Rightarrow -8 < 3x - 1 < 8 \Rightarrow -7 < 3x < 9 \Rightarrow -\frac{7}{3} < x < 3$. Therefore, the solution is $\left(-\frac{7}{3}, 3\right)$.
67. $|15 - x| < 7 \Rightarrow -7 < 15 - x < 7 \Rightarrow 22 > x > 8$ or $8 < x < 22$. Therefore, the solution is $(8, 22)$.
68. $|7 - 4x| \leq 11 \Rightarrow -11 \leq 7 - 4x \leq 11 \Rightarrow -18 \leq -4x \leq 4 \Rightarrow -1 \leq x \leq \frac{9}{2}$. Therefore, the solution is $\left[-1, \frac{9}{2}\right]$.

69. $|2x-3| > 1 \Rightarrow 2x-3 > 1$ or $2x-3 < -1 \Rightarrow 2x > 4$ or $\left(\frac{f}{g}\right)(-3) = \frac{-3(-3)-4}{(-3)^2} = \frac{5}{9}$. or $x < 1$. Therefore,

the solution is $(-\infty, 1) \cup (2, \infty)$.

70. $|4-3x| > 1 \Rightarrow 4-3x > 1$ or $4-3x < -1 \Rightarrow -3x > -3$ or $-3x < -5 \Rightarrow x < 1$ or $x > \frac{5}{3}$. Therefore, the solution is

$$(-\infty, 1) \cup \left(\frac{5}{3}, \infty\right).$$

71. $|-3x+8| \geq 3 \Rightarrow -3x+8 \geq 3$ or $-3x+8 \leq -3 \Rightarrow -3x \geq -5$ or $-3x \leq -11 \Rightarrow x \leq \frac{5}{3}$ or $x \geq \frac{11}{3}$.

Therefore, the solution is $\left(-\infty, \frac{5}{3}\right] \cup \left[\frac{11}{3}, \infty\right)$.

72. Absolute value is always positive, and so is always greater than -1 ; therefore, the solution is $(-\infty, \infty)$.

73. $\left|6 - \frac{1}{3}x\right| > 0 \Rightarrow 6 - \frac{1}{3}x > 0$ or $6 - \frac{1}{3}x < 0 \Rightarrow -\frac{1}{3}x > -6$ or $-\frac{1}{3}x < -6 \Rightarrow x < 18$ or $x > 18$. Therefore,

the solution is every real number except 18: $(-\infty, 18) \cup (18, \infty)$.

74. Absolute value is always positive, and cannot be less than 0; therefore, the solution set is $\emptyset$.

75. Absolute value is always positive, and so cannot be less than or equal to -6 ; therefore, the solution set is $\emptyset$.

76. Absolute value is always positive, and so cannot be less than -4 ; therefore, the solution set is $\emptyset$.

77. Absolute value is always positive, and so is always greater than -5 ; therefore, the solution is $(-\infty, \infty)$.

78. To solve such an equation, we must solve the compound equation $ax+b=cx+d$ or $ax+b=cx+d$. The solution set consists of the union of the two individual solution sets.

79. (a) $3x+1=2x-7 \Rightarrow x+1=-7 \Rightarrow x=-8$ $3x+1=-(2x-7) \Rightarrow 3x+1=-2x+7 \Rightarrow 5x=6 \Rightarrow x=\frac{6}{5}$.

Therefore, the solution set is $\left\{-8, \frac{6}{5}\right\}$.

(b) Graph $y_1 = |3x+1|$ and $y_2 = |2x-7|$. See Figure 79. From the graph, $|f(x)| > |g(x)|$ when $y_1 > y_2$

which is for the interval $(-\infty, -8) \cup \left(\frac{6}{5}, \infty\right)$.

(c) Graph $y_1 = |3x+1|$ and $y_2 = |2x-7|$. See Figure 79. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$

which is for the interval $\left(-8, \frac{6}{5}\right)$.

80. (a) $x-4=7x+12 \Rightarrow -6x-4=12 \Rightarrow -6x=16 \Rightarrow x=-\frac{8}{3}$ or $x-4=-(7x+12) \Rightarrow$

$x-4=-7x-12 \Rightarrow 8x-4=-12 \Rightarrow 8x=-8 \Rightarrow x=-1$. Therefore, the solution set is $\left\{-\frac{8}{3}, -1\right\}$.

(b) Graph $y_1 = |x - 4|$ and $y_2 = |7x + 12|$. See Figure 80. From the graph, $|f(x)| > |g(x)|$ when $y_1 > y_2$ which is for the interval $\left(-\frac{8}{3}, -1\right)$.

(c) Graph $y_1 = |x - 4|$ and $y_2 = |7x + 12|$. See Figure 80. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$ which is for the interval $\left(-\infty, -\frac{8}{3}\right) \cup (-1, \infty)$.

$[-20, 20]$ by $[-10, 50]$
Xscl = 2 Yscl = 5




Figure 79

$[-10, 10]$ by $[-4, 16]$
Xscl = 1 Yscl = 1




Figure 80

81. (a) $-2x + 5 = x + 3 \Rightarrow -3x = -2 \Rightarrow x = \frac{2}{3}$ or $-2x + 5 = -(x + 3) \Rightarrow -2x + 5 = -x - 3 \Rightarrow$

$-x = -8 \Rightarrow x = 8$. Therefore, the solution set is $\left\{\frac{2}{3}, 8\right\}$.

(b) Graph $y_1 = |-2x + 5|$ and $y_2 = |x + 3|$. See Figure 81. From the graph, $|f(x)| > |g(x)|$ when $y_1 > y_2$, which it is for the interval $\left(-\infty, \frac{2}{3}\right) \cup (8, \infty)$.

(c) Graph $y_1 = |-2x + 5|$ and $y_2 = |x + 3|$. See Figure 81. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(\frac{2}{3}, 8\right)$.

82. (a) $-5x + 1 = 3x - 4 \Rightarrow -8x = -5 \Rightarrow x = \frac{5}{8}$ or $-5x + 1 = -(3x - 4) \Rightarrow -5x + 1 = -3x + 4 \Rightarrow -2x = 3 \Rightarrow x = -\frac{3}{2}$.

Therefore, the solution set is $\left\{-\frac{3}{2}, \frac{5}{8}\right\}$.

(b) Graph $y_1 = |-5x + 1|$ and $y_2 = |3x - 4|$. See Figure 82. From the graph, $|f(x)| > |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\infty, -\frac{3}{2}\right) \cup \left(\frac{5}{8}, \infty\right)$.

(c) Graph $y_1 = |-5x + 1|$ and $y_2 = |3x - 4|$. See Figure 82. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\frac{3}{2}, \frac{5}{8}\right)$.




Figure 81




Figure 82




Figure 83

83. (a) $x - \frac{1}{2} = \frac{1}{2}x - 2 \Rightarrow \frac{1}{2}x = -\frac{3}{2} \Rightarrow x = -3$ or $x - \frac{1}{2} = -\left(\frac{1}{2}x - 2\right) \Rightarrow x - \frac{1}{2} = -\frac{1}{2}x + 2 \Rightarrow$

$\frac{3}{2}x = \frac{5}{2} \Rightarrow x = \frac{5}{3}$. Therefore, the solution set is $\left\{-3, \frac{5}{3}\right\}$.

(b) Graph $y_1 = \left|x - \frac{1}{2}\right|$ and $y_2 = \left|\frac{1}{2}x - 2\right|$. From the graph, $|f(x)| > |g(x)|$ when $y_1 > y_2$, which it is for the interval $(-\infty, -3) \cup \left(\frac{5}{3}, \infty\right)$.

(c) Graph $y_1 = \left|x - \frac{1}{2}\right|$ and $y_2 = \left|\frac{1}{2}x - 2\right|$. See Figure 83. From the graph $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-3, \frac{5}{3}\right)$.

84. (a) $x + 3 = \frac{1}{3}x + 8 \Rightarrow \frac{2}{3}x = 5 \Rightarrow x = \frac{15}{2}$ or $x + 3 = -\left(\frac{1}{3}x + 8\right) \Rightarrow x + 3 = -\frac{1}{3}x - 8 \Rightarrow$

$\frac{4}{3}x = -11 \Rightarrow x = -\frac{33}{4}$. Therefore, the solution set is $\left\{-\frac{33}{4}, \frac{15}{2}\right\}$.

(b) Graph $y_1 = |x + 3|$ and $y_2 = \left|\frac{1}{3}x + 8\right|$. See Figure 84. From the graph, $|f(x)| > |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\infty, -\frac{33}{4}\right) \cup \left(\frac{15}{2}, \infty\right)$.

(c) Graph $y_1 = |x + 3|$ and $y_2 = \left|\frac{1}{3}x + 8\right|$. See Figure 84. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\frac{33}{4}, \frac{15}{2}\right)$.

85. (a) $4x + 1 = 4x + 6 \Rightarrow 1 = 6 \Rightarrow \emptyset$ or $4x + 1 = -(4x + 6) \Rightarrow 4x + 1 = -4x - 6 \Rightarrow 8x = -7 \Rightarrow -\frac{7}{8}$.

Therefore, the solution set is $\left\{-\frac{7}{8}\right\}$.

(b). Graph $y_1 = |4x + 1|$ or $y_2 = |4x + 6|$. See Figure 85. From the graph $|f(x)| > |g(x)|$ when $y_1 > y_2$, which it is for the interval $\left(-\infty, \frac{7}{8}\right)$.

- (c) Graph $y_1 = |4x+1|$ or $y_2 = |4x+6|$. See Figure 85. From the graph, $|f(x)| < |g(x)|$ when $y_1 > y_2$, which is for the interval $\left(-\frac{7}{8}, \infty\right)$.




Figure 84




Figure 85




Figure 86

86. (a) $6x+9=6x-3 \Rightarrow 9=-3 \Rightarrow \emptyset$ or $6x+9=-(6x-3) \Rightarrow 6x+9=-6x+3 \Rightarrow 12x=-6 \Rightarrow -6 \Rightarrow x = -\frac{6}{12} = -\frac{1}{2}$. Therefore, the solution set is $\left\{-\frac{1}{2}\right\}$.
- (b) Graph $y_1 = |6x+9|$ and $y_2 = |6x-3|$. See Figure 86. From the graph, $|f(x)| > |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\frac{1}{2}, \infty\right)$.
- (c) Graph $y_1 = |6x+9|$ and $y_2 = |6x-3|$. See Figure 86. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $\left(-\infty, -\frac{1}{2}\right)$.
87. (a) $0.25x+1=0.75x-3 \Rightarrow -0.50x=-4 \Rightarrow x=8$ or $0.25x+1=-(0.75x-3) \Rightarrow 0.25x+1=-0.75x+3 \Rightarrow x=2$. Therefore, the solution set is $\{2, 8\}$.
- (b) Graph $y_1 = |.25x+1|$ and $y_2 = |.75x-3|$. See Figure 87. From the graph, $|f(x)| > |g(x)|$ when $y_1 < y_2$, which it is for the interval $(2, 8)$.
- (c) Graph $y_1 = |.25x+1|$ and $y_2 = |.75x-3|$. See Figure 87. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $(-\infty, 2) \cup (8, \infty)$.
88. (a) $.40x+2=.60x-5 \Rightarrow -20x=-7 \Rightarrow x=35$ or $.40x+2=-(.60x-5) \Rightarrow .40x+2=-.60x+5 \Rightarrow x=3$. Therefore, the solution set is $\{3, 35\}$.
- (b) Graph $y_1 = |.40x+2|$ and $y_2 = |.60x-5|$. See Figure 88. From the graph, $|f(x)| > |g(x)|$ when $y_1 > y_2$, which it is for the interval $(3, 35)$.
- (c) Graph $y_1 = |.40x+2|$ and $y_2 = |.60x-5|$. See Figure 88. From the graph, $|f(x)| < |g(x)|$ when $y_1 < y_2$, which it is for the interval $(-\infty, 3) \cup (35, \infty)$.

[-20,20] by [-4,16]




Figure 87

[-30,50] by [-5,30]




Figure 88

[-10,10] by [-10,10]




Figure 89

[-10,10] by [-10,10]




Figure 90

89. (a) $3x+10 = -(-3x-10) \Rightarrow 3x+10 = 3x+10 \Rightarrow$ there are an infinite number of solutions.

Therefore, the solution set is $(-\infty, \infty)$.

- (b) Graph $y_1 = |3x+10|$ and $y_2 = |-3x-10|$. See Figure 89. From the graph, $|f(x)| > |g(x)|$

when $y_1 > y_2$, for which there is no solution.

- (c) Graph $y_1 = |3x+10|$ and $y_2 = |-3x-10|$. See Figure 89. From the graph, $|f(x)| < |g(x)|$

when $y_1 < y_2$, for which there is no solution.

90. (a) $5x-6 = -(-5x+6) \Rightarrow 5x-6 = 5x-6 \Rightarrow$ there are an infinite number of solutions.

Therefore, the solution set is $(-\infty, \infty)$.

- (b) Graph $y_1 = |5x-6|$ and $y_2 = |-5x+6|$. See Figure 90. From the graph, $|f(x)| > |g(x)|$

when $y_1 > y_2$, for which there is no solution.

- (c) Graph $y_1 = |5x-6|$ and $y_2 = |-5x+6|$. See Figure 90. From the graph, $|f(x)| < |g(x)|$

when $y_1 < y_2$, for which there is no solution.

91. Graph $y_1 = |x+1| + |x-6|$ and $y_2 = 11$. See Figure 91. From the graph, the lines intersect at $(-3, 11)$

and $(2, 9)$. Therefore, the solution set is $\{-3, 8\}$.

92. Graph $y_1 = |2x+2| + |x+1|$ and $y_2 = 9$. See Figure 92. From the graph, the lines intersect at $(-4, 9)$ and

$(2, 9)$. Therefore, the solution set is $\{-4, 2\}$.

93. Graph $y_1 = |x| + |x-4|$ and $y_2 = 8$. See Figure 93. From the graph, the lines intersect at $(-2, 8)$ and $(6, 8)$.

Therefore, the solution set is $\{-2, 6\}$.

94. Graph $y_1 = |.5x+2| + |.25x+4|$ and $y_2 = 9$. See Figure 94. From the graph, the lines intersect at $(-20, 9)$

and $(4, 9)$. Therefore, the solution set is $\{-20, 4\}$.

[-10,10] by [-4,16]

Xscl = 1

Yscl = 1




Figure 91

[-10,10] by [-4,16]

Xscl = 1

Yscl = 1




Figure 92

[-10,10] by [-4,16]

Xscl = 1

Yscl = 1




Figure 93

[-30,10] by [-4,16]

Xscl = 5

Yscl = 1




Figure 94

95. (a) $|T - 50| \leq 22 \Rightarrow -22 \leq T - 50 \leq 22 \Rightarrow 28 \leq T \leq 72.$

(b) The average monthly temperatures in Boston vary between a low of 28°F and a high of 72°F . The monthly averages are always within 22° of 50°F .

96. (a) $|T - 10| \leq 36 \Rightarrow -36 \leq T - 10 \leq 36 \Rightarrow -26 \leq T \leq 46..$

(b) The average monthly temperatures in Chesterfield vary between a low of -26°F and a high of 46°F . The monthly averages are always within 36° of 10°F .

97. (a) $|T - 61.5| \leq 12.5 \Rightarrow -12.5 \leq T - 61.5 \leq 12.5 \Rightarrow 49 \leq T \leq 74..$

(b) The average monthly temperatures in Buenos Aires vary between a low of 49°F (possibly in July) and a high of 74°F (possibly in January). The monthly averages are always within 12.5° of 61.5°F .

98. (a) $|T - 43.5| \leq 8.5 \Rightarrow -8.5 \leq T - 43.5 \leq 8.5 \Rightarrow 35 \leq T \leq 52.$

(b) The average monthly temperatures in Punta Arenas vary between a low of 35°F and a high of 52°F . The monthly averages are always within 8.5° of 43.5°F .

99. (a) The max circumference is 10.1 and the minimum circumference is 9.9. This can be represented as $|C - 10| \leq 0.1$

(b) $C = \pi d \Rightarrow d = \frac{C}{\pi}$ The maximum diameter is $\frac{10}{\pi} + \frac{0.1}{\pi}$ and the minimum diameter is $\frac{10}{\pi} - \frac{0.1}{\pi}.$

This can be represented as $\left|d - \frac{10}{\pi}\right| \leq \frac{0.1}{\pi}.$

100. $|H - 5| \leq 0.05$

101. (a) $|(2x - 1) - 1| \leq 0.1$

(b) $|(2x - 1) - 1| \leq 0.1 \Rightarrow |2x| \leq 0.1 \Rightarrow -0.1 \leq 2x \leq 0.1 \Rightarrow -0.05 \leq x \leq 0.05$

102. (a) $|(3x - 6) - 0| \leq 0.3$

(b) $|(3x - 6)| \leq 0.3 \Rightarrow -0.3 \leq 3x - 6 \leq 0.3 \Rightarrow 5.7 \leq 3x \leq 6.3 \Rightarrow 1.9 \leq x \leq 2.1$

103. (a) $|(4x - 8) - 4| \leq 0.04$

- (b) $|(4x-8)-4| \leq 0.04 \Rightarrow |4x-12| \leq 0.04 \Rightarrow -0.04 \leq 4x-12 \leq 0.04 \Rightarrow 11.96 \leq 4x \leq 12.04 \Rightarrow 2.99 \leq x \leq 3.01$
104. (a) $|(5x-12)-10| \leq 0.1$
- (b) $|(5x-12)-10| \leq 0.1 \Rightarrow |5x-22| \leq 0.1 \Rightarrow -0.1 \leq 5x-22 \leq 0.1 \Rightarrow 21.9 \leq 5x \leq 22.1 \Rightarrow 4.38 \leq x \leq 4.42$
105. (a) $|(x-4)-0.5| \leq 0.001$
- (b) $|(x-4)-0.5| \leq 0.001 \Rightarrow |x-4.5| \leq 0.001 \Rightarrow -0.001 \leq x-4.5 \leq 0.001 \Rightarrow 4.499 \leq x \leq 4.501$
106. (a) $|(7-x)-1.2| \leq 0.002$
- (b) $|(7-x)-1.2| \leq 0.002 \Rightarrow |5.8-x| \leq 0.002 \Rightarrow -0.002 \leq 5.8-x \leq 0.002 \Rightarrow -5.802 \leq -x \leq -5.798 \Rightarrow 5.798 \leq x \leq 5.802$
107. $|x-8.0| \leq 1.5 \Rightarrow -1.5 \leq x-8.0 \leq 1.5 \Rightarrow 6.5 \leq x \leq 9.5$; therefore, the range is the interval $[6.5, 9.5]$.
108. If $\frac{680+780}{2} = 730$ is the midpoint, then $680 \leq F \leq 780 \Rightarrow 680-730 \leq F-730 \leq 780-730 \Rightarrow -50 \leq F-730 \leq 50 \Rightarrow |F-730| \leq 50$ (or $|730-F| \leq 50$).
109. (a) $P_d = |116-125| \Rightarrow P_d = |-9| = 9$.
- (b) $17 = |P-120| \Rightarrow P-120 = 17$ or $P-120 = -17 \Rightarrow P = 137$ or $P = 103$.
110. If $\frac{98+148}{2} = 123$ is the midpoint, then $98 \leq x \leq 148 \Rightarrow 98-123 \leq x-123 \leq 148-123 \Rightarrow -25 \leq x-123 \leq 25 \Rightarrow |x-123| \leq 25$ (or $|123-x| \leq 25$); and if $\frac{16+26}{2} = 21$ is the midpoint, then $16 \leq x \leq 26 \Rightarrow 16-21 \leq x-21 \leq 26-21 \Rightarrow -5 \leq x-21 \leq 5 \Rightarrow |x-21| \leq 5$ (or $|21-x| \leq 5$).
11. If $|2x+7| = 6x-1$ then $|2x+7| - (6x-1) = 0$. Graph $y_1 = |2x+7| - (6x-1)$, See Figure 111. The x -intercept is 2; therefore, the solution set is $\{2\}$.
112. If $-|3x-12| \geq -x-1$ then $-|3x-12| - (-x-1) \geq 0$. Graph $y_1 = -|3x-12| - (-x-1)$, See Figure 112. The equation is ≥ 0 , or the graph intersects or is above the x -axis, for the interval: $[2.75, 6.5]$.
113. If $|x-4| > .5x-6$ then $|x-4| - (.5x-6) > 0$. Graph $y_1 = |x-4| - (.5x-6)$, See Figure 113. The equation is > 0 , or the graph is above the x -axis, for the interval: $(-\infty, \infty)$.
114. If $2x+8 > -|3x+4|$ then $2x+8 - (-|3x+4|) > 0$. Graph $y_1 = 2x+8 - (-|3x+4|)$, See Figure 114. The equation is > 0 , or the graph is above the x -axis, for the interval: $(-\infty, \infty)$.

$[-10,10]$ by $[-10,10]$
Xscl = 1 Yscl = 1




Figure 111

$[-10,10]$ by $[-10,10]$
Xscl = 1 Yscl = 1




Figure 112

$[-10,10]$ by $[-10,10]$
Xscl = 1 Yscl = 1




Figure 113

$[-10,10]$ by $[-4,16]$
Xscl = 1 Yscl = 1




Figure 114

115. If $|3x+4| < -3x-14$ then $|3x+4| - (-3x-14) < 0$. Graph $y_1 = |3x+4| - (-3x-14)$. See Figure 115. The equation is < 0 , or the graph is below the x-axis, never or for the solution set: $\emptyset$.

116. If $|x-\sqrt{13}| + \sqrt{6} \leq -x-\sqrt{10}$ then $|x-\sqrt{13}| + \sqrt{6} - (-x-\sqrt{10}) \leq 0$. Graph

$y_1 = |x-\sqrt{13}| + \sqrt{6} - (-x-\sqrt{10})$. See Figure 100. The equation is ≤ 0 , or the graph intersects or is below the x-axis, never or for the solution set: $\emptyset$.

$[-10,10]$ by $[-10,10]$
Xscl = 1 Yscl = 1




Figure 115

$[-10,10]$ by $[-10,10]$
Xscl = 1 Yscl = 1




Figure 116

2.5: Piecewise-Defined Functions

- From the graph, the speed limit is 40 mph.
 - From the graph, the speed limit is 30 mph for 6 miles.
 - From the graph, $f(5) = 40$ mph; $f(13) = 30$ mph; and $f(19) = 55$ mph.
 - From the graph, the graph is discontinuous at $x = 4, 6, 8, 12$, and 16 . The speed limit changes at each discontinuity.
- From the graph, the Initial amount was: \$1,000; and the final amount was: \$600.
 - From the graph, $f(10) = \$900$; $f(50) = \$600$. The function f is not continuous.
 - From the graph, the discontinuity shows 3 drops or withdrawals.
 - From the graph, the largest drop or largest withdrawal of \$300 occurred after 15 minutes.
 - From the graph, the one increase or deposit was \$200.
- From the graph, the Initial amount was: 50,000 gal.; and the final amount was: 30,000 gal.
 - From the graph, during the first and fourth days.
 - From the graph, $f(2) = 45,000$ gal; $f(4) = 40,000$ gal.

- (d) From the graph, between days 1 and 3 the water dropped: $\frac{50,000 - 40,000}{2} = \frac{10,000}{2} = 5,000$ gal./day.
4. (a) From the graph, when $x = 1$ the tank had 20 gallons of gas.
 (b) Since $x = 0$ represents 3:15 pm and 3 hours later the tank was filled to 20 gallons, the time was 6:15 pm
 (c) When the graph is horizontal, the engine is not running; when the graph is decreasing, the engine is burning gasoline; and when the graph is increasing, gasoline is being put into the tank.
 (d) When the graph is decreasing, gasoline was burned the fastest between 1 and 2.9 hours
5. (a) $f(-5) = 2(-5) = -10$ (b) $f(-1) = 2(-1) = -2$
 (c) $f(0) = 0 - 1 = -1$ (d) $f(3) = 3 - 1 = 2$
6. (a) $f(-5) = -5 - 2 = -7$ (b) $f(-1) = -1 - 2 = -3$
 (c) $f(0) = 0 - 2 = -2$ (d) $f(3) = 5 - 3 = 2$
7. (a) $f(-5) = 2 + (-5) = -3$ (b) $f(-1) = -(-1) = 1$
 (c) $f(0) = -(0) = 0$ (d) $f(3) = 3(3) = 9$
8. (a) $f(-5) = -2(-5) = 10$ (b) $f(-1) = 3(-1) - 1 = -4$
 (c) $f(0) = 3(0) - 1 = -1$ (d) $f(3) = -4(3) = -12$
9. Yes, continuous. See Figure 9.
 10. Yes, continuous. See Figure 10.
 11. Not continuous. See Figure 11




Figure 9




Figure 10




Figure 11

12. Not continuous. See Figure 12.
 13. Not continuous. See Figure 13.
 14. Not continuous. See Figure 14.




Figure 12




Figure 13




Figure 14

15. Not continuous. See Figure 15.
 16. Not continuous. See Figure 16.
 17. Yes, continuous. See Figure 17.




Figure 15




Figure 16




Figure 17

18. Not continuous. See Figure 18.
 19. Yes, continuous. See Figure 19.
 20. Yes, continuous. See Figure 20.




Figure 18




Figure 19




Figure 20

21. Look for a $y = x^2$ graph if $x \geq 0$; and a linear graph if $x < 0$. Therefore: B.
 22. Look for a $y = |x|$ graph if $x \geq -1$; and a reflected $y = x^2$ graph if $x < -1$. Therefore: A.
 23. Look for a horizontal graph above the x-axis if $x \geq 0$; and a horizontal graph below the x-axis if $x < 0$. Therefore: D.
 24. Look for a $y = \sqrt{x}$ graph if $x \geq 0$; and a reflected $y = x^2$ graph if $x < 0$. Therefore: C.
 25. Graph $y_1 = (x-1)(x \leq 3) + (2)(x > 3)$, See Figure 25.
 26. Graph $y_1 = (6-x)(x \leq 3) + (3x-6)(x > 3)$, See Figure 26.
 27. Graph $y_1 = (4-x)(x < 2) + (1+2x)(x \geq 2)$, See Figure 27.
 28. Graph $y_1 = (2x+1)(x \geq 0) + (x)(x < 0)$, See Figure 28.




Figure 25




Figure 26




Figure 27




Figure 28

29. Graph $y_1 = (2+x)(x < -4) + (-x)(-4 \leq x \text{ and } x \leq 5) + (3x)(x > 5)$, See Figure 29.

30. Graph $y_1 = (-2x)(x < -3) + (3x-1)(-3 \leq x \text{ and } x \leq 2) + (-4x)(x > 2)$, See Figure 30.

31. Graph $y_1 = \left(-\frac{1}{2}x + 2\right)(x \leq 2) + \left(\frac{1}{2}x\right)(x > 2)$, See Figure 31.

32. Graph $y_1 = (x^3 + 5)(x \leq 0) + (-x^2)(x > 0)$, See Figure 32.




Figure 29




Figure 30




Figure 31




Figure 32

33. From the graph, the function is $f(x) = \begin{cases} 2 & \text{if } x \leq 0 \\ -1 & \text{if } x > 1 \end{cases}$; domain: $(-\infty, 0] \cup (1, \infty)$; range: $\{-1, 2\}$.

34. From the graph, the function is $f(x) = \begin{cases} 1 & \text{if } x \leq -1 \\ -1 & \text{if } x > 2 \end{cases}$; domain: $(-\infty, -1] \cup (2, \infty)$; range: $\{-1, 1\}$.

35. From the graph, the function is $f(x) = \begin{cases} x & \text{if } x \leq 0 \\ 2 & \text{if } x > 0 \end{cases}$; domain: $(-\infty, \infty)$; range: $(-\infty, 0] \cup \{2\}$.


36. From the graph, the function is $f(x) = \begin{cases} -3 & \text{if } x < 0 \\ \sqrt{x} & \text{if } x \geq 0 \end{cases}$; domain: $(-\infty, \infty)$; range: $\{-3\} \cup [0, \infty)$.

37. From the graph, the function is $f(x) = \begin{cases} \sqrt[3]{x} & \text{if } x < 1 \\ x+1 & \text{if } x \geq 1 \end{cases}$; domain: $(-\infty, \infty)$; range: $(-\infty, 1) \cup [2, \infty)$.

38. From the graph, the function is $f(x) = \begin{cases} 3 & \text{if } x = 2 \\ 2x-3 & \text{if } x \neq 2 \end{cases}$; domain: $(-\infty, \infty)$; range: $(-\infty, 1) \cup (1, \infty)$.

39. There is an overlap of intervals since the number 4 satisfies both conditions. To be a function, every x -value is used only once.

40. The value $f(4)$ cannot be found since using the first formula, $f(4) = 11$, and using the second formula, $f(4) = 16$. To have two different values violates the definition of function.
41. The graph of $y = \lfloor x \rfloor$ is shifted 1.5 units downward.
42. The graph of $y = \lfloor x \rfloor$ is reflected across the y -axis.
43. The graph of $y = \lfloor x \rfloor$ is reflected across the x -axis.
44. The graph of $y = \lfloor x \rfloor$ is shifted 2 units to the left.
45. Graph $y = \lfloor x \rfloor - 1.5$, See Figure 45.
46. Graph $y = \lfloor -x \rfloor$, See Figure 46.
47. Graph $y = -\lfloor x \rfloor$, See Figure 47.
48. Graph $y = \lfloor x + 2 \rfloor$, See Figure 48.



49. (a) In 2012 there were 200 million users (2012, 200) and in 2015 there were 700 million users (2015, 700).

The slope is given as $\frac{700 - 200}{2015 - 2012} = \frac{500}{3}$. The equation is given as $W - 200 = \frac{500}{3}(x - 2012) \Rightarrow$

$W = \frac{500}{3}(x - 2012) + 200$ between 2012 and 2015 inclusively.

In 2018 there were 1000 million users (2018, 1000). The slope is given as $\frac{1000 - 700}{2018 - 2015} = \frac{300}{3} = 100$.

The equation is $W - 700 = 100(x - 2015) \Rightarrow W = 100(x - 2015) + 700$ between 2015 and 2018.

- (b) See Figure 49. The function is continuous on this time period.
- (c) From 2012 to 2015, the number of active WhatsApp users increased, on average, by about 166.7 million users per year. From 2015 to 2018, this increase slowed to 100 million per year, on average.
50. (a) In January 2014 there were 46 million users (0, 46) and in July 2015 there were 94 million users (17, 94). The slope is given as $\frac{94 - 46}{18 - 0} = \frac{48}{18} = \frac{8}{3}$. The equation is given as $S = \frac{8}{3}x + 46$ between 0 and 18 months inclusively.

In January 2017 there were 160 million users (36, 160). The slope is given as

$\frac{160 - 94}{36 - 18} = \frac{66}{18} = \frac{11}{3}$. The equation is given as $S - 160 = \frac{11}{3}(x - 36) \Rightarrow S = \frac{11}{3}x + 28$ between 18 and 36 months.

- (b) See Figure 50. The function is continuous on the time interval.
- (c) From January 2014 to July 2015, the number of daily active Snapchat users increased, on average, by about 2.7 million per month. From July 2015 to January 2017, this increased to 3.7 million per month, on average.




Figure 49




Figure 50

51. When $0 \leq x \leq 3$ the slope is 5, which means the inlet pipe is open and the outlet pipe is closed; when $3 < x \leq 5$ the slope is 2, which means both pipes are open; when $5 < x \leq 8$ the slope is 0, which means both pipes are closed; when $8 < x \leq 10$ the slope is -3 , which means the inlet pipe is closed and the outlet pipe is open.
52. (a) Since for each year given the shoe size is 1 size smaller than his age, the formula is $y = x - 1$
- (b) From the table and question, graph the function. See figure 50.
53. (a) $f(1.5) = 0.70$, $f(3) = 0.91$. It costs \$0.70 to mail 1.5 oz and \$0.91 to mail 3 oz.
- (b) Domain: $(0, 5]$; Range: $\{0.49, 0.70, 0.91, 1.12, 1.33\}$. See figure 53.
54. (a) $f(1967) = 0.0475(1967) - 93.3 = 0.1325$, In 1967, the average Super Bowl ad cost \$0.1325 million.
- (b) $f(1998) = 0.475(1998) - 93.3 = 1.605$, In 1998, the average Super Bowl ad cost \$1.605 million.
- (c) $f(2013) = 0.179(2013) - 356.037 \approx 5.01$, In 2013, the average Super Bowl ad cost \$5.01 million
- (d) The function $f(x)$ is continuous on its domain since there are no breaks in the graph of $f(x)$.
- (e) See Figure 54.




Figure 52




Figure 53




Figure 54

55. (a) From the table, graph the piecewise function. See Figure 55.
 (b) The likelihood of being a victim peaks from age 16 up to age 20, then decreases.
56. (a) From the table, graph the piecewise function. See Figure 56.
 (b) Housing starts increased, decreased, and increased, with the maximum occurring during the 1960's.
57. (a) From the graph, the highest speed is 60 mph and the lowest speed is 20 mph.
 (b) There are approximately 4 miles of highway with a speed of 55 mph.
 (c) From the graph, $f(4) = 40$; $f(12) = 20$; $f(18) = 7$.
58. From the table, graph the piecewise function. See Figure 58.




Figure 55




Figure 56




Figure 58

59. (a) A 3.5 minute call would round up to 4 minutes. A 4 minute call would cost: $.50 + 3(.25) = \$1.25$.
 (b) We use a piecewise defined function where the cost increases after each whole number as follows:

$$f(x) = \begin{cases} .50 & \text{if } 0 < x \leq 1 \\ .75 & \text{if } 1 < x \leq 2 \\ 1.00 & \text{if } 2 < x \leq 3 \\ 1.25 & \text{if } 3 < x \leq 4 \\ 1.50 & \text{if } 4 < x \leq 5 \end{cases} \quad \text{Another possibility is } f(x) = \begin{cases} .50 & \text{if } 0 < x \leq 1 \\ .50 - .25 \lfloor 1 - x \rfloor & \text{if } 1 < x \leq 5 \end{cases}$$

60. (a) Since cost is rounded down to the nearest 2 foot interval, we can use the greatest integer function. The

function is $f(x) = .80 \left\lfloor \frac{x}{2} \right\rfloor$ from $6 \leq x \leq 18$.

- (b) Graph $f(x) = .80 \left\lfloor \frac{x}{2} \right\rfloor$ from $6 \leq x \leq 18$. See Figure 60.

(c) $f(8.5) = .80 \left\lfloor \frac{8.5}{2} \right\rfloor = .80 \lfloor 4.25 \rfloor = .80(4) \Rightarrow f(8.5) = \3.20

$f(15.2) = .80 \left\lfloor \frac{15.2}{2} \right\rfloor = .80 \lfloor 7.6 \rfloor = .80(7) \Rightarrow f(15.2) = \5.60




Figure 60

61. For x in the interval $(0, 2]$, $y = 25$. For x in the interval $(2, 3]$, $y = 25 + 3 = 28$. For x in the interval $(3, 4]$, $y = 28 + 3 = 31$ and so on. The graph is a step function. In this case, the first step has a different width. See Figure 61.
62. Sketch a piecewise function that measures miles over minutes. The first piece increases at a rate of 40 mph or $\frac{40}{60} = \frac{2}{3}$ miles per minute, for 30 minutes (the time it takes to travel 20 miles at that rate). The second piece stays at a constant distance of 20 miles for the 2 hour period at the park. The third piece decreases (returns home) at a rate of 20 mph or $\frac{20}{60} = \frac{1}{3}$ miles per minute, for 60 minutes (the time it takes to travel 20 miles at that rate). See Figure 62.
63. Sketch a piecewise function that fills a tank at a rate of 5 gallons a minute for the first 20 minutes (the time it takes to fill the 100 gallon tank) and then drains the tank at a rate of 2 gallons per minute for 50 minutes (the time it takes to drain the 100 gallon tank). See Figure 63.




Figure 61




Figure 62




Figure 63

64. (a) Use the points (1990, 1.3) and (2007, 2.7) to find the slope. $m = \frac{2.7-1.3}{2007-1990} = \frac{1.4}{17} \approx 0.082$
- $y - y_1 = m(x - x_1) \Rightarrow y - 1.3 = 0.082(x - 1990) \Rightarrow y = 0.082x + 161.88$ Use the points (2007, 2.7) and (2016, 1.5) to find the slope. $m = \frac{1.5-2.7}{2016-2007} = -\frac{1.2}{9} \approx -0.13$ $y - y_1 = m(x - x_1) \Rightarrow$
- $y - 1.5 = -0.13(x - 2016) \Rightarrow y = -0.13x + 263.58$.
- (b) Use the two functions above to create $f(x) = \begin{cases} 0.082x + 161.88 & 1990 \leq x \leq 2007 \\ -0.13x + 263.58 & 2007 < x \leq 2016 \end{cases}$

2.6: Operations and Composition

1. $x^2 + (2x - 5) = x^2 + 2x - 5 \Rightarrow E.$
2. $x^2 - (2x - 5) = x^2 - 2x + 5 \Rightarrow B.$
3. $x^2(2x - 5) = 2x^3 - 5x^2 \Rightarrow F.$
4. $\frac{x^2}{2x - 5} \Rightarrow D.$
5. $(2x - 5)^2 = 4x^2 - 20x + 25 \Rightarrow A.$
6. $(2(x^2) - 5) = 2x^2 - 5 \Rightarrow C.$
7. $(f \circ g)(3) = f(g(3)) = f(2(3) - 1) = f(5) = 5^2 + 3(5) = 40$
8. $(g \circ f)(-2) = g(f(-2)) = g((-2)^2 + 3(-2)) = g(-2) = 2(-2) - 1 = -5$
9. $(f \circ g)(x) = f(g(x)) = f(2x - 1) = (2x - 1)^2 + 3(2x - 1) = 4x^2 - 4x + 1 + 6x - 3 = 4x^2 + 2x - 2$
10. $(g \circ f)(x) = g(f(x)) = g(x^2 + 3x) = 2(x^2 + 3x) - 1 = 2x^2 + 6x - 1$
11. $(f + g)(3) = f(3) + g(3) = ((3)^2 + 3(3)) + (2(3) - 1) = 23$
12. $(f + g)(-5) = f(-5) + g(-5) = ((-5)^2 + 3(-5)) + (2(-5) - 1) = -1$
13. $(f \cdot g)(4) = f(4) \cdot g(4) = ((4)^2 + 3(4)) \cdot (2(4) - 1) = 196$
14. $(f \cdot g)(-3) = f(-3) \cdot g(-3) = ((-3)^2 + 3(-3)) \cdot (2(-3) - 1) = 0$
15. $\left(\frac{f}{g}\right)(-1) = \frac{f(-1)}{g(-1)} = \frac{(-1)^2 + 3(-1)}{2(-1) - 1} = \frac{2}{3}$
16. $\left(\frac{f}{g}\right)(4) = \frac{f(4)}{g(4)} = \frac{(4)^2 + 3(4)}{2(4) - 1} = \frac{28}{7} = 4$
17. $(f - g)(2) = f(2) - g(2) = ((2)^2 + 3(2)) - (2(2) - 1) = 7$
18. $(f - g)(-2) = f(-2) - g(-2) = ((-2)^2 + 3(-2)) - (2(-2) - 1) = 3$
19. $(g - f)(-2) = g(-2) - f(-2) = (2(-2) - 1) - ((-2)^2 + 3(-2)) = -3$
20. $\left(\frac{f}{g}\right)\left(\frac{1}{2}\right) = \frac{f\left(\frac{1}{2}\right)}{g\left(\frac{1}{2}\right)} = \frac{\left(\frac{1}{2}\right)^2 + 3\left(\frac{1}{2}\right)}{2\left(\frac{1}{2}\right) - 1} = \frac{\frac{7}{4}}{0} \Rightarrow \text{undefined.}$
21. $\left(\frac{g}{f}\right)(0) = \frac{g(0)}{f(0)} = \frac{2(0) - 1}{(0)^2 + 3(0)} = \frac{-1}{0} \Rightarrow \text{undefined.}$
22. Because the two compositions are opposites of each other.

23. (a) $(f+g)(x) = (4x-1) + (6x+3) = 10x+2$, $(f-g)(x) = (4x-1) - (6x+3) = -2x-4$
 $(fg)(x) = (4x-1)(6x+3) = 24x^2 + 12x - 6x - 3 = 24x^2 + 6x - 3$
 (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
 (c) $\left(\frac{f}{g}\right)(x) = \frac{4x-1}{6x+3}$; all values can replace x , except $-\frac{1}{2}$; therefore, the domain is $\left(-\infty, -\frac{1}{2}\right) \cup \left(-\frac{1}{2}, \infty\right)$.
 (d) $(f \circ g)(x) = f[g(x)] = 4(6x+3) - 1 = 24x + 12 - 1 = 24x + 11$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$
 (e) $(g \circ f)(x) = g[f(x)] = 6(4x-1) + 3 = 24x - 6 + 3 = 24x - 3$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
24. (a) $(f+g)(x) = (9-2x) + (-5x+2) = -7x+11$, $(f-g)(x) = (9-2x) - (-5x+2) = 3x+7$
 $(fg)(x) = (9-2x)(-5x+2) = -45x + 18 + 10x^2 - 4x = 10x^2 - 49x + 18$
 (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
 (c) $\left(\frac{f}{g}\right)(x) = \frac{9-2x}{-5x+2}$; all values can replace x , except $\frac{2}{5}$; therefore, the domain is $\left(-\infty, \frac{2}{5}\right) \cup \left(\frac{2}{5}, \infty\right)$.
 (d) $(f \circ g)(x) = f[g(x)] = 9 - 2(-5x+2) = 9 + 10x - 4 = 10x + 5$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$
 (e) $(g \circ f)(x) = g[f(x)] = -5(9-2x) + 2 = -45 + 10x + 2 = 10x - 43$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
25. (a) $(f+g)(x) = |x+3| + 2x$, $(f-g)(x) = |x+3| - 2x$, $(fg)(x) = |x+3|(2x)$
 (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
 (c) $\left(\frac{f}{g}\right)(x) = \frac{|x+3|}{2x}$; all values can replace x , except 0; therefore, the domain is $(-\infty, 0) \cup (0, \infty)$
 (d) $(f \circ g)(x) = f[g(x)] = |(2x)+3| = |2x+3|$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
 (e) $(g \circ f)(x) = g[f(x)] = 2(|x+3|) = 2|x+3|$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
26. (a) $(f+g)(x) = |2x-4| + (x+1) = |2x-4| + x + 1$, $(f-g)(x) = |2x-4| - (x+1) = |2x-4| - x - 1$
 $(fg)(x) = |2x-4|(x+1) = (x+1)|2x-4|$
 (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
 (c) $\left(\frac{f}{g}\right)(x) = \frac{|2x-4|}{x+1}$ all values can replace x , except -1 ; therefore, the domain is $(-\infty, -1) \cup (-1, \infty)$.
 (d) $(f \circ g)(x) = f[g(x)] = |2(x+1) - 4| = |2x - 2|$; all values can replace x , so the domain is $(-\infty, \infty)$.
 (e) $(g \circ f)(x) = g[f(x)] = |2x-4| + 1$; all values can replace x , so the domain is $(-\infty, \infty)$.

27. (a) $(f+g)(x) = \sqrt[3]{x+4} + (x^3+5) = \sqrt[3]{x+4} + x^3 + 5$, $(f-g)(x) = \sqrt[3]{x+4} - (x^3+5) = \sqrt[3]{x+4} - x^3 - 5$
 $(fg)(x) = (\sqrt[3]{x+4})(x^3+5)$
- (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
- (c) $\left(\frac{f}{g}\right)(x) = \frac{\sqrt[3]{x+4}}{(x^3+5)}$ all values can replace x , except $\sqrt[3]{-5}$, so the domain is $(-\infty, \sqrt[3]{-5}) \cup (\sqrt[3]{-5}, \infty)$.
- (d) $(f \circ g)(x) = f[g(x)] = \sqrt[3]{(x^3+5)+4} = \sqrt[3]{x^3+9}$; all values can replace x , so the domain is $(-\infty, \infty)$.
- (e) $(g \circ f)(x) = g[f(x)] = (\sqrt[3]{x+4})^3 + 5 = x+4+5 = x+9$; all values can replace x , so the domain is $(-\infty, \infty)$.
28. (a) $(f+g)(x) = \sqrt[3]{6-3x} + (2x^3+1) = \sqrt[3]{6-3x} + 2x^3 + 1$
 $(f-g)(x) = \sqrt[3]{6-3x} - (2x^3+1) = \sqrt[3]{6-3x} - 2x^3 - 1$
 $(fg)(x) = (\sqrt[3]{6-3x})(2x^3+1)$
- (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
- (c) $\left(\frac{f}{g}\right)(x) = \frac{\sqrt[3]{6-3x}}{(2x^3+1)}$; all values can replace x , except $\sqrt[3]{-\frac{1}{2}}$, so the domain is $\left(-\infty, \sqrt[3]{-\frac{1}{2}}\right) \cup \left(\sqrt[3]{-\frac{1}{2}}, \infty\right)$
- (d) $(f \circ g)(x) = f[g(x)] = \sqrt[3]{6-3(2x^3+1)} = \sqrt[3]{-6x^3+3}$; all values can replace x , so the domain is $(-\infty, \infty)$.
- (e) $(g \circ f)(x) = g[f(x)] = 2(\sqrt[3]{6-3x})^3 + 1 = 2(6-3x) + 1 = -6x + 13$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
29. (a) $(f+g)(x) = \sqrt{x^2+3} + (x+1) = \sqrt{x^2+3} + x + 1$, $(f-g)(x) = \sqrt{x^2+3} - (x+1) = \sqrt{x^2+3} - x - 1$
 $(fg)(x) = (\sqrt{x^2+3})(x+1)$
- (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
- (c) $\left(\frac{f}{g}\right)(x) = \frac{\sqrt{x^2+3}}{(x+1)}$; all values can replace x , except -1 ; therefore, the domain is $(-\infty, -1) \cup (-1, \infty)$.
- (d) $(f \circ g)(x) = f[g(x)] = \sqrt{(x+1)^2+3} = \sqrt{x^2+2x+1+3} = \sqrt{x^2+2x+4}$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
- (e) $(g \circ f)(x) = g[f(x)] = (\sqrt{x^2+3}) + 1 = \sqrt{x^2+3} + 1$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.

30. (a) $(f + g)(x) = \sqrt{2 + 4x^2} + (x) = \sqrt{2 + 4x^2} + x$, $(f - g)(x) = \sqrt{2 + 4x^2} - (x) = \sqrt{2 + 4x^2} - x$
 $(fg)(x) = \left(\sqrt{2 + 4x^2}\right)(x) = x\sqrt{2 + 4x^2}$
- (b) All values can replace x in all three equations; therefore, the domain is $(-\infty, \infty)$ in all cases.
- (c) $\left(\frac{f}{g}\right)(x) = \frac{\sqrt{2 + 4x^2}}{x}$; all values can replace x , except 0; therefore, the domain is $(-\infty, 0) \cup (0, \infty)$.
- (d) $(f \circ g)(x) = f[g(x)] = \sqrt{2 + 4(x)^2} = \sqrt{2 + 4x^2}$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
- (e) $(g \circ f)(x) = g[f(x)] = \left(\sqrt{2 + 4x^2}\right) = \sqrt{2 + 4x^2}$; all values can replace x ; therefore, the domain is $(-\infty, \infty)$.
31. (a) From the graph, $4 + (-2) = 2$.
- (b) From the graph, $1 - (-3) = 4$.
- (c) From the graph, $(0)(-4) = 0$.
- (d) From the graph, $\frac{1}{-3} = -\frac{1}{3}$.
32. (a) From the graph, $0 + 2 = 2$.
- (b) From the graph, $-2 - 1 = -3$.
- (c) From the graph, $(2)(1) = 2$.
- (d) From the graph, $\frac{4}{-2} = -2$.
33. (a) From the graph, $0 + 3 = 3$.
- (b) From the graph, $-1 - 4 = -5$.
- (c) From the graph, $(2)(1) = 2$.
- (d) From the graph, $\frac{3}{0} \Rightarrow$ undefined.
34. (a) From the graph, $-3 + 1 = -2$.
- (b) From the graph, $-2 - 0 = -2$.
- (c) From the graph, $(-3)(-1) = 3$.
- (d) From the graph, $\frac{-3}{1} = -3$.
35. (a) From the table, $7 + (-2) = 5$.
- (b) From the table, $10 - 5 = 5$.
- (c) From the table, $(0)(6) = 0$.
- (d) From the table, $\frac{5}{0} =$ undefined.

36. (a) From the table, $5 + 4 = 9$.
 (b) From the table, $0 - 0 = 0$.
 (c) From the table, $(-4)(2) = -8$.
 (d) From the table, $\frac{8}{-1} = -8$.

37. See Table 37.

38. See Table 38.

x	$(f + g)(x)$	$(f - g)(x)$	$(fg)(x)$	$\left(\frac{f}{g}\right)(x)$
-2	6	-6	0	0
0	5	5	0	undefined
2	5	9	-14	-3.5
4	15	5	50	2

Figure 37

x	$(f + g)(x)$	$(f - g)(x)$	$(fg)(x)$	$\left(\frac{f}{g}\right)(x)$
-2	-2	-6	-8	-2
0	7	9	-8	-8
2	9	1	20	1.25
4	0	0	0	undefined

Figure 38

39. (a) $(f \circ g)(4) = f[g(4)]$, so from the graph find $g(4) = 0$. Now find $f(0) = -4$; therefore, $(f \circ g)(4) = -4$.
 (b) $(g \circ f)(3) = g[f(3)]$, so from the graph find $f(3) = 2$. Now find $g(2) = 2$; therefore, $(g \circ f)(3) = 2$.
 (c) $(f \circ f)(2) = f[f(2)]$, so from the graph find $f(2) = 0$. Now find $f(0) = -4$; therefore, $(f \circ f)(2) = -4$.
40. (a) $(f \circ g)(2) = f[g(2)]$, so from the graph find $g(2) = -2$. Now find $f(-2) = -4$; therefore,
 $(f \circ g)(2) = -4$.
 (b) $(g \circ g)(0) = g[g(0)]$, so from the graph find $g(0) = 2$. Now find $g(2) = -2$; therefore, $(g \circ g)(0) = -2$.
 (c) $(g \circ f)(4) = g[f(4)]$, so from the graph find $f(4) = 2$. Now find $g(2) = -2$; therefore,
 $(g \circ f)(4) = -2$.
41. (a) $(f \circ g)(1) = f[g(1)]$, so from the graph find $g(1) = 2$. Now find $f(2) = -3$; therefore, $(f \circ g)(1) = -3$.
 (b) $(g \circ f)(-2) = g[f(-2)]$, so from the graph find $f(-2) = -3$. Now find $g(-3) = -2$; therefore,
 $(g \circ f)(-2) = -2$.
 (c) $(g \circ g)(-2) = g[g(-2)]$, so from the graph find $g(-2) = -1$. Now find $g(-1) = 0$; therefore,
 $(g \circ g)(-1) = 0$.
42. (a) $(f \circ g)(-2) = f[g(-2)]$, so from the graph find $g(-2) = 4$. Now find $f(4) = 2$; therefore,
 $(f \circ g)(-2) = 2$.
 (b) $(g \circ f)(1) = g[f(1)]$, so from the graph find $f(1) = 1$. Now find $g(1) = 1$; therefore, $(g \circ f)(1) = 1$.
 (c) $(f \circ f)(0) = f[f(0)]$, so from the graph find $f(0) = 0$. Now find $f(0) = 0$; therefore, $(f \circ f)(0) = 0$.
43. (a) $(g \circ f)(1) = g[f(1)]$, so from the table find $f(1) = 4$. Now find $g(4) = 5$; therefore, $(g \circ f)(1) = 5$.
 (b) $(f \circ g)(4) = f[g(4)]$, so from the table find $g(4) = 5$. $f(5)$ is undefined; therefore, $(f \circ g)(4)$ is undefined.

- (c) $(f \circ f)(3) = f[f(3)]$, so from the table find $f(3) = 1$. Now find $f(1) = 4$; therefore, $(f \circ f)(3) = 4$.
44. (a) $(g \circ f)(1) = g[f(1)]$, so from the table find $f(1) = 2$. Now find $g(2) = 4$; therefore, $(g \circ f)(1) = 4$.
- (b) $(f \circ g)(4) = f[g(4)]$, so from the table we find $g(4)$ is undefined; therefore, $(f \circ g)(4)$ is undefined.
- (c) $(f \circ f)(3) = f[f(3)]$, so from the table find $f(3) = 6$. Now find $f(6) = 7$; therefore, $(f \circ f)(3) = 7$.
45. From the table, $g(3) = 4$ and $f(4) = 2$.
46. From the table, $f(6) = 7$ and $g(7) = 0$.
47. Since $Y_3 = Y_1 \circ Y_2$ and $X = -1$, we solve $Y_1[Y_2(-1)]$. First solve $Y_2 = (-1)^2 = 1$, now solve $Y_1 = 2(1) - 5 = -3$; therefore, $Y_3 = -3$.
48. Since $Y_3 = Y_1 \circ Y_2$ and $X = -2$, we solve $Y_1[Y_2(-2)]$. First solve $Y_2 = (-2)^2 = 4$, now solve $Y_1 = 2(4) - 5 = 3$; therefore, $Y_3 = 3$.
49. Since $Y_3 = Y_1 \circ Y_2$ and $X = 7$, we solve $Y_1[Y_2(7)]$. First solve $Y_2 = (7)^2 = 49$, now solve $Y_1 = 2(49) - 5 = 93$; therefore, $Y_3 = 93$.
50. Since $Y_3 = Y_1 \circ Y_2$ and $X = 8$, we solve $Y_1[Y_2(8)]$. First solve $Y_2 = (8)^2 = 64$, now solve $Y_1 = 2(64) - 5 = 123$; therefore, $Y_3 = 123$.
51. (a) $(f \circ g)(x) = f[g(x)] = (x^2 + 3x - 1)^3$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = (x^3)^2 + 3(x^3) - 1 = x^6 + 3x^3 - 1$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = (x^3)^3 = x^9$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
52. (a) $(f \circ g)(x) = f[g(x)] = 2 - \left(\frac{1}{x^2}\right) = 2 - \frac{1}{x^2}$; all values can be input for x , except 0; therefore, the domain is $(-\infty, 0) \cup (0, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = \frac{1}{(2-x)^2}$; all values can be input for x , except 2; therefore, the domain is $(-\infty, 2) \cup (2, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = 2 - (2 - x) = x$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
53. (a) $(f \circ g)(x) = f[g(x)] = (\sqrt{1-x})^2 = 1-x$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = (\sqrt{1-(x^2)}) = \sqrt{1-x^2}$; only values where $x^2 \leq 1$ can be input for x ; therefore, the domain is $[-1, 1]$.
- (c) $(f \circ f)(x) = f[f(x)] = (x^2)^2 = x^4$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.

54. (a) $(f \circ g)(x) = f[g(x)] = (x^4 + x^2 - 3x - 4) + 2 = x^4 + x^2 - 3x - 2$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = (x+2)^4 + (x+2)^2 - 3(x+2) - 4$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = (x+2) + 2 = x+4$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
55. (a) $(f \circ g)(x) = f[g(x)] = \frac{1}{(5x)+1} = \frac{1}{5x+1}$; all values can be input for x , except $-\frac{1}{5}$; therefore, the domain is $\left(-\infty, -\frac{1}{5}\right) \cup \left(-\frac{1}{5}, \infty\right)$.
- (b) $(g \circ f)(x) = g[f(x)] = 5\left(\frac{1}{x+1}\right) = \frac{5}{x+1}$; all values can be input for x , except -1 ; therefore, the domain is $(-\infty, -1) \cup (-1, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = \frac{1}{\left(\frac{1}{x+1}\right)+1} = \frac{1}{\frac{1}{x+1} + \frac{x+1}{x+1}} = \frac{1}{\frac{x+2}{x+1}} = \frac{x+1}{x+2}$; all values can be input for x , except those that make $\frac{x+1}{x+2} = 0$ or undefined. That would be -1 and -2 ; therefore, the domain is $(-\infty, -2) \cup (-2, -1) \cup (-1, \infty)$.
56. (a) $(f \circ g)(x) = f[g(x)] = (\sqrt{4-x^2}) + 4 = \sqrt{4-x^2} + 4$; only values where $x^2 \leq 4$ can be input for x ; therefore, the domain is $[-2, 2]$.
- (b) $(g \circ f)(x) = g[f(x)] = \sqrt{4-(x+4)^2}$; only values where $(x+4)^2 \leq 4$ can be input for x ; therefore, the domain is $[-6, -2]$.
- (c) $(f \circ f)(x) = f[f(x)] = (x+4) + 4 = x+8$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
57. (a) $(f \circ g)(x) = f[g(x)] = 2(4x^3 - 5x^2) + 1 = 8x^3 - 10x^2 + 1$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = 4(2x+1)^3 - 5(2x+1)^2 = 4(8x^3 + 12x^2 + 6x + 1) - 5(4x^2 + 4x + 1) = 32x^3 + 48x^2 + 24x + 4 - (20x^2 + 20x + 5) = 32x^3 + 28x^2 + 4x - 1$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = 2(2x+1) + 1 = 4x+3$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
58. (a) $(f \circ g)(x) = f[g(x)] = \frac{(2x+3)-3}{2} = \frac{2x}{3} = x$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.

- (b) $(g \circ f)(x) = g[f(x)] = 2\left(\frac{x-3}{2}\right) + 3 = (x-3) + 3 = x$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = \frac{\left(\frac{x-3}{2}\right) - 3}{2} = \frac{\left(\frac{x-3}{2} - \frac{6}{2}\right)}{2} = \frac{\left(\frac{x-9}{2}\right)}{2} = \frac{x-9}{4}$; all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
59. (a) $(f \circ g)(x) = f(g(x)) = f(5) = 5$, all values can be input for x , therefore the domain is $(-\infty, \infty)$.
- (b) $(g \circ f)(x) = g(f(x)) = g(5) = 5$, all values can be input for x , therefore the domain is $(-\infty, \infty)$.
- (c) $(f \circ f)(x) = f(f(x)) = f(5) = 5$, all values can be input for x , therefore the domain is $(-\infty, \infty)$.
60. (a) $(f \circ g)(x) = f[g(x)] = \frac{1}{1-x}$ all values can but 1 can be input for x ; therefore, the domain is $(-\infty, 1) \cup (1, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = 1 - \frac{1}{x} = \frac{x-1}{x}$ all values but 0 can be input for x ; therefore, the domain is $(-\infty, 0) \cup (0, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = \frac{1}{\frac{1}{x}} = x$ all values can be input for x ; therefore, the domain is $(-\infty, \infty)$.
61. (a) $(f \circ g)(x) = f[g(x)] = \frac{1}{\sqrt{x-2}}$ all values can but 4 can be input for x ; therefore, the domain is $(-\infty, 4) \cup (4, \infty)$.
- (b) $(g \circ f)(x) = g[f(x)] = \sqrt{\frac{1}{x-2}} = \frac{1}{\sqrt{x-2}} = \frac{\sqrt{x-2}}{x-2}$ all values bigger than 2 can be input for x ; therefore, the domain is $(2, \infty)$.
- (c) $(f \circ f)(x) = f[f(x)] = \frac{1}{\frac{1}{x-2} - 2} = \frac{x-2}{1-2(x-2)} = \frac{x-2}{5-2x}$ all values but $\frac{5}{2}$ can be input for x ; therefore, the domain is $\left(-\infty, \frac{5}{2}\right) \cup \left(\frac{5}{2}, \infty\right)$.
62. (a) $(f \circ g)(x) = f[g(x)] = \frac{1}{3-6x+1} = \frac{1}{4-6x}$ all values can but $\frac{2}{3}$ can be input for x ; therefore, the domain is $\left(-\infty, \frac{2}{3}\right) \cup \left(\frac{2}{3}, \infty\right)$.

(b) $(g \circ f)(x) = g[f(x)] = 3 - 6\left(\frac{1}{x+1}\right) = 3 - \frac{6}{x+1}$ all values but -1 can be input for x ; therefore,

the domain is $(-\infty, -1) \cup (-1, \infty)$.

(c) $(f \circ f)(x) = f[f(x)] = \frac{1}{\frac{1}{x+1} + 1} = \frac{x+1}{1+(x+1)} = \frac{x+1}{x+2}$ all values but -2 can be input for x ; therefore, the

domain is $(-\infty, -2) \cup (-2, \infty)$.

63. Let $f(x) = c$, then $(f \circ f)(x) = f(f(x)) = f(c) = c$. Therefore, there is only one value in the range of $(f \circ f)(x)$.

64. Let $f(x) = k$ and $g(x) = mx + b$ then $(f \circ g)(x) = f(g(x)) = k$. Therefore, there is only one value k in the range of $(f \circ g)(x)$.

65. $(f \circ g)(x) = f[g(x)] = 4\left(\frac{1}{4}(x-2)\right) + 2 = x - 2 + 2 = x$, $(g \circ f)(x) = g[f(x)] = \frac{1}{4}((4x+2)-2) = \frac{1}{4}(4x) = x$

66. $(f \circ g)(x) = f[g(x)] = -3\left(-\frac{1}{3x}\right) = x$, $(g \circ f)(x) = g[f(x)] = -\frac{1}{3}(-3x) = x$

67. $(f \circ g)(x) = f[g(x)] = \sqrt[3]{5\left(\frac{1}{5}x^3 - \frac{4}{5}\right) + 4} = \sqrt[3]{(x^3 - 4) + 4} = \sqrt[3]{x^3} = x$

$(g \circ f)(x) = g[f(x)] = \frac{1}{5}\left(\sqrt[3]{5x+4}\right)^3 - \frac{4}{5} = \frac{1}{5}(5x+4) - \frac{4}{5} = x + \frac{4}{5} - \frac{4}{5} = x$

68. $(f \circ g)(x) = f[g(x)] = \sqrt[3]{(x^3 - 1) + 1} = \sqrt[3]{x^3} = x$, $(g \circ f)(x) = g[f(x)] = \left(\sqrt[3]{x+1}\right)^3 - 1 = x + 1 - 1 = x$

69. Graph $y_1 = \sqrt[3]{x-6}$, $y_2 = x^3 + 6$, and $y_3 = x$ in the same viewing window. See Figures 69. The graph of y_2 can be obtained by *reflecting* the graph of y_1 across the line $y_3 = x$.

70. Graph $y_1 = 5x - 3$, $y_2 = \frac{1}{5}(x+3)$, and $y_3 = x$ in the same viewing window. See Figures 70. The graph of y_2 can be obtained by *reflecting* the graph of y_1 across the line $y_3 = x$.




Figure 69




Figure 70

71. (a) $f(x) = x^2 - 4 \Rightarrow f(x+h) = (x+h)^2 - 4 = x^2 + 2xh + h^2 - 4$;

- (b) $f(x) + f(h) = (x^2 - 4) + (h^2 - 4) = x^2 + h^2 - 8$
72. (a) $f(x) = 5x^2 + x \Rightarrow f(x+h) = 5(x+h)^2 + (x+h) = 5(x^2 + 2xh + h^2) + (x+h) = 5x^2 + 10xh + 5h^2 + x + h;$
 (b) $f(x) + f(h) = (5x^2 + x) + (5h^2 + h) = 5x^2 + x + 5h^2 + h$
73. (a) $f(x) = 3x - x^2 \Rightarrow f(x+h) = 3(x+h) - (x+h)^2 = 3x + 3h - (x^2 + 2xh + h^2) = -x^2 - 2xh + 3x - h^2 + 3h;$
 (b) $f(x) + f(h) = (3x - x^2) + (3h - h^2) = -x^2 + 3x - h^2 + 3h$
74. (a) $f(x) = x^3 \Rightarrow f(x+h) = (x+h)^3 = (x^2 + 2xh + h^2)(x+h) = x^3 + 3hx^2 + 3h^2x + h^3$
 (b) $f(x) + f(h) = x^3 + h^3$
75. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{4(x+h) + 3 - (4x + 3)}{h} = \frac{4x + 4h + 3 - 4x - 3}{h} = \frac{4h}{h} = 4.$
76. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{5(x+h) - 6 - (5x - 6)}{h} = \frac{5x + 5h - 6 - 5x + 6}{h} = \frac{5h}{h} = 5.$
77. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{-6(x+h)^2 - (x+h) + 4 - (-6x^2 - x + 4)}{h}$

$$= \frac{-6(x^2 + 2xh + h^2) - x - h + 4 + 6x^2 + x - 4}{h} = \frac{-6x^2 - 12xh - 6h^2 - x - h + 4 + 6x^2 + x - 4}{h}$$

$$= \frac{-12xh - 6h^2 - h}{h} = -12x - 6h - 1.$$
78. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{\frac{1}{2}(x+h)^2 + 4(x+h) - \left(\frac{1}{2}x^2 + 4x\right)}{h}$

$$= \frac{\frac{1}{2}(x^2 + 2xh + h^2) + 4x + 4h - \frac{1}{2}x^2 - 4x}{h} = \frac{\frac{1}{2}x^2 + xh + \frac{1}{2}h^2 + 4x + 4h - \frac{1}{2}x^2 - 4x}{h}$$

$$= \frac{xh + \frac{1}{2}h^2 + 4h}{h} = x + \frac{1}{2}h + 4.$$
79. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{(x+h)^3 - x^3}{h} = \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h} = \frac{3x^2h + 3xh^2 + h^3}{h} = 3x^2 + 3xh + h^2.$
80. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{-2(x+h)^3 - (-2x^3)}{h} = \frac{-2(x^3 + 3x^2h + 3xh^2 + h^3) + 2x^3}{h}$

$$\frac{-2x^3 - 6x^2h - 6xh^2 - 2h^3 + 2x^3}{h} = \frac{-6x^2h - 6xh^2 - 2h^3}{h} = -6x^2 - 6xh - 2h^2.$$
81. Using $\frac{f(x+h) - f(x)}{h}$ gives: $\frac{1 - (x+h)^2 - (1 - x^2)}{h} = \frac{1 - (x^2 + 2xh + h^2) - 1 + x^2}{h}$

$$\frac{1 - x^2 - 2xh - h^2 - 1 + x^2}{h} = \frac{-2xh - h^2}{h} = -2x - h$$

$$82. \text{ Using } \frac{f(x+h)-f(x)}{h} \text{ gives: } \frac{(x+h)^2+2(x+h)-(x^2+2x)}{h} = \frac{(x^2+2xh+h^2)+2x+2h-x^2-2x}{h} = \frac{2xh+h^2+2h}{h} = 2x+h+2$$

$$83. \text{ Using } \frac{f(x+h)-f(x)}{h} \text{ gives: } \frac{3(x+h)^2-(3x^2)}{h} = \frac{3(x^2+2xh+h^2)-3x^2}{h} = \frac{3x^2+6xh+3h^2-3x^2}{h} = \frac{6xh+3h^2}{h} = 6x+3h$$

$$84. \text{ Using } \frac{f(x+h)-f(x)}{h} \text{ gives: } \frac{\sqrt{x+h}-\sqrt{x}}{h} = \frac{(\sqrt{x+h}-\sqrt{x})(\sqrt{x+h}+\sqrt{x})}{h(\sqrt{x+h}+\sqrt{x})} = \frac{x+h-x}{h(\sqrt{x+h}+\sqrt{x})} = \frac{h}{h(\sqrt{x+h}+\sqrt{x})} = \frac{1}{\sqrt{x+h}+\sqrt{x}}$$

$$85. \text{ Using } \frac{f(x+h)-f(x)}{h} \text{ gives: } \frac{\frac{1}{2(x+h)}-\frac{1}{2x}}{h} = \frac{\frac{2x-(2x+2h)}{2x(2x+2h)}}{h} = \frac{\frac{-2h}{2x(2x+2h)}}{h} = \frac{-1}{2x(x+h)}$$

$$86. \text{ Using } \frac{f(x+h)-f(x)}{h} \text{ gives: } \frac{\frac{1}{(x+h)^2}-\frac{1}{x^2}}{h} = \frac{\frac{x^2-(x+h)^2}{x^2(x+h)^2}}{h} = \frac{\frac{x^2-x^2-2xh-h^2}{x^2(x+h)^2}}{h} = \frac{\frac{-2xh-h^2}{x^2(x+h)^2}}{h} = \frac{h(-2x-h)}{x^2(x+h)^2} = \frac{-2x-h}{x^2(x+h)^2}$$

$$87. \text{ One possible solution is } f(x) = x^2 \text{ and } g(x) = 6x-2. \text{ Then } (f \circ g)(x) = f[g(x)] = (6x-2)^2.$$

$$88. \text{ One possible solution is } f(x) = x^2 \text{ and } g(x) = 11x^2+12x. \text{ Then } (f \circ g)(x) = f[g(x)] = (11x^2+12x)^2.$$

$$89. \text{ One possible solution is } f(x) = \sqrt{x} \text{ and } g(x) = x^2-1. \text{ Then } (f \circ g)(x) = f[g(x)] = \sqrt{x^2-1}.$$

$$90. \text{ One possible solution is } f(x) = x^3 \text{ and } g(x) = 2x-3. \text{ Then } (f \circ g)(x) = f[g(x)] = (2x-3)^3.$$

$$91. \text{ One possible solution is } f(x) = \sqrt{x}+12 \text{ and } g(x) = 6x. \text{ Then } (f \circ g)(x) = f[g(x)] = \sqrt{6x}+12.$$

$$92. \text{ One possible solution is } f(x) = \sqrt[3]{x}-4 \text{ and } g(x) = 2x+3. \text{ Then } (f \circ g)(x) = f[g(x)] = \sqrt[3]{2x+3}-4.$$

$$93. (a) \text{ There are 5280 feet in a mile. } F(x) = 5280x$$

$$(b) \text{ There are 12 inches in a foot. } I(x) = 12x$$

$$(c) (I \circ F)(x) = I(F(x)) = I(5280x) = 12(5280x) = 63,360x$$

$$(d) \text{ The number of inches in a mile.}$$

$$94. (a) \text{ There are 2000 pounds in a ton. } T(x) = 2000x$$

$$(b) \text{ There are 16 ounces in a pound. } O(x) = 16x$$

$$(c) (O \circ T)(x) = O(T(x)) = O(2000x) = 16(2000x) = 32,000x$$

- (d) The number of ounces in a ton.
95. (a) With a cost of \$10 to produce each item and a fixed cost of \$500, the cost function is $C(x) = 10x + 500$.
 (b) With a selling price of \$35 for each item, the revenue function is $R(x) = 35x$.
 (c) The profit function is $P(x) = R(x) - C(x) \Rightarrow P(x) = 35x - (10x + 500) \Rightarrow P(x) = 25x - 500$.
 (d) A profit is shown when $P(x) > 0 \Rightarrow 25x - 500 > 0 \Rightarrow 25x > 500 \Rightarrow x > 20$. Therefore, 21 items must be produced and sold to realize a profit.
 (e) Graph $y_1 = 25x - 500$. The smallest whole number for which $P(x) > 0$ is 21. Use a window of $[0, 30]$ by $[-1000, 500]$, for example.
96. (a) With a cost of \$11 to produce each item and a fixed cost of \$180, the cost function is $C(x) = 11x + 180$.
 (b) With a selling price of \$20 for each item, the revenue function is $R(x) = 20x$.
 (c) The profit function is $P(x) = R(x) - C(x) \Rightarrow P(x) = 20x - (11x + 180) \Rightarrow P(x) = 9x - 180$.
 (d) A profit is shown when $P(x) > 0 \Rightarrow 9x - 180 > 0 \Rightarrow 9x > 180 \Rightarrow x > 20$. Therefore, 21 items must be produced and sold to realize a profit.
 (e) Graph $y_1 = 9x - 180$. The smallest whole number for which $P(x) > 0$ is 21. Use a window of $[-5, 30]$ by $[-200, 200]$, for example.
97. (a) With a cost of \$100 to produce each item and a fixed cost of \$2700, the cost function is $C(x) = 100x + 2700$.
 (b) With a selling price of \$280 for each item, the revenue function is $R(x) = 280x$.
 (c) The profit function is $P(x) = R(x) - C(x) \Rightarrow P(x) = 280x - (100x + 2700) \Rightarrow P(x) = 180x - 2700$.
 (d) A profit is shown when $P(x) > 0 \Rightarrow 180x - 2700 > 0 \Rightarrow 180x > 2700 \Rightarrow x > 15$. Therefore, 16 items must be produced and sold to realize a profit.
 (e) Graph $y_1 = 180x - 2700$, the smallest whole number for which $P(x) > 0$ is 16. Use a window of $[0, 30]$ by $[-3000, 500]$, for example.
98. (a) With a cost of \$200 to produce each item and a fixed cost of \$1000, the cost function is $C(x) = 200x + 1000$.
 (b) With a selling price of \$240 for each item, the revenue function is $R(x) = 240x$.
 (c) The profit function is $P(x) = R(x) - C(x) \Rightarrow P(x) = 240x - (200x + 1000) \Rightarrow P(x) = 40x - 1000$.
 (d) A profit is shown when $P(x) > 0 \Rightarrow 40x - 1000 > 0 \Rightarrow 40x > 1000 \Rightarrow x > 25$. Therefore, 26 items must be produced and sold to realize a profit.
 (e) Graph $y_1 = 40x - 1000$, the smallest whole number for which $P(x) > 0$ is 26. Use a window of $[-5, 40]$ by $[-1200, 600]$, for example.
99. (a) If $V(r) = \frac{4}{3}\pi r^3$, then a 3 inch increase would be $V(r) = \frac{4}{3}\pi(r+3)^3$, and the volume gained would be
- $$V(r) = \frac{4}{3}\pi(r+3)^3 - \frac{4}{3}\pi r^3.$$
- (b) Graph $y_1 = \frac{4}{3}\pi(x+3)^3 - \frac{4}{3}\pi x^3$ in the window $[0, 10]$ by $[0, 1500]$. See Figure 99. Although this appears to be a portion of a parabola, it is actually a cubic function.
- (c) From the graph in exercise 91b, an input value of $x = 4$ results in a gain of $y \approx 1168.67$.

$$(d) \quad V(4) = \frac{4}{3}\pi(4+3)^3 - \frac{4}{3}\pi(4)^3 = \frac{4}{3}\pi(343) - \frac{4}{3}\pi(64) = \frac{1372}{3}\pi - \frac{256}{3}\pi = \frac{1116}{3}\pi = 372\pi \approx 1168.67.$$

100. If $S(r) = 4\pi r^2$, then doubling the radius would give us a surface area gained function of

$$S(r) = 4\pi(2r)^2 - 4\pi r^2 = 16\pi r^2 - 4\pi r^2 = 12\pi r^2.$$

101. (a) If $x =$ width, then $2x =$ length. Since the perimeter formula is $P = 2W + 2L$ our perimeter function is $P(x) = 2(x) + 2(2x) = 2x + 4x \Rightarrow P(x) = 6x$. This is a linear function.
- (b) Graph $P(x) = 6x$ in the window $[0,10]$ by $[1,100]$. See Figure 101b. From the graph when $x = 4, y = 24$. The 4 represents the width of a rectangle and 24 represents the perimeter.
- (c) If $x = 4$ is the width of a rectangle then $2x = 8$ is the length. See Figure 101c. Using the standard perimeter formula yields $P = 2(4) + 2(8) = 24$. This compares favorably with the graph result in part b.
- (d) (Answers may vary.) If the perimeter y of a rectangle satisfying the given conditions is 36, then the width x is 6. See Figure 101d.




Figure 99




Figure 101b




Figure 101c




Figure 101d

102. (a) If $x = 4s$ then $s = \frac{x}{4}$.
- (b) If $y = s^2$ and $s = \frac{x}{4}$, then $y(x) = \left(\frac{x}{4}\right)^2 \Rightarrow y(x) = \frac{x^2}{16}$.
- (c) Since x is the perimeter and $x = 6, y(6) = \frac{(6)^2}{16} = \frac{36}{16} = \frac{9}{4} = 2.25$.
- (d) Show that the point $(6, 2.25)$ is on the graph $y = \frac{x^2}{16}$. A square with perimeter 6 will have area 2.25 square units.
103. (a) $A(2x) = \frac{\sqrt{3}}{4}(2x)^2 = \frac{\sqrt{3}}{4}(4x^2) \Rightarrow A(2x) = \sqrt{3}x^2$
- (b) $A(x) = \frac{\sqrt{3}}{4}(16)^2 = \frac{\sqrt{3}}{4}(256) \Rightarrow A(x) = 64\sqrt{3}$ square units.
- (c) On the graph of $y = \frac{\sqrt{3}}{4}x^2$, locate the point where $x = 16$ to find $y \approx 110.85$, an approximation for $64\sqrt{3}$.

104. (a) If $A(r) = \pi r^2$ and $r(t) = 2t$ then $(A \circ r)(t) = A[r(t)] = A[2t] = \pi(2t)^2 = 4\pi t^2$.
 (b) $(A \circ r)(t)$ is a composite function that expresses the area of the circular region covered by the pollutants as a function of time t (in hours).
 (c) Since $t = 0$ is 8 A.M., noon would be $t = 4$. $(A \circ r)(4) = 4\pi(4)^2 = 64\pi \text{ mi}^2$.
 (d) Graph $y_1 = 4\pi x^2$ and show that for $x = 4$, $y \approx 201$ (an approximation for 64π).
105. (a) $A(2100) = 42$, ; The average age of a person in 2100 is projected to be 42 years. $T(2100) = 430$, In 2100, the living world's population will have a combined life experience of 430 billion years.
 (b) $\frac{T(2100)}{A(2100)} = \frac{430}{42} \approx 10.2$, The world population will be about 10.2 billion in 2100.
 (c) $P(x)$ gives the world's population during year x .
106. (a) The domain is the years. See table below. Therefore, $D = \{ 2013, 2014, 2015, 2016 \}$.

x	2013	2014	2015	2016
$h(x)$	77.7	84.8	85.1	83.7

- (b) The function h computes the number of acres of soybeans harvested in the U.S. in millions of acres.
107. (a) $(f + g)(2010) = 13.0 + 74.3 = 87.3$
 (b) The function $(f + g)(x)$ computes the total SO_2 and Carbon Monoxide during year x .
 (c) Add functions f and g .

x	1970	1980	1990	2000	2010
$(f+g)(x)$	235.2	211.3	177.3	130.8	87.3

108. (a) The function h is the addition of functions f and g .

x	1990	2000	2010	2020	2030
$h(x)$	32	35.5	39	42.5	46

- (b) The function h is the addition of functions f and g . Therefore, $h(x) = f(x) + g(x)$.
109. (a) The function h is the subtraction of function f from g . Therefore, $h(x) = g(x) - f(x)$.
 (b) $h(2000) = g(2000) - f(2000) = 40 - 30 = 10$, $h(2014) = g(2014) - f(2014) = 100 - 80 = 20$
 (c) Using the points (2014, 100) and (2000, 40), the slope is $m = \frac{100 - 40}{2014 - 2000} = \frac{60}{14} = \frac{30}{7}$. The equation is $y - 100 = \frac{30}{7}(x - 2014)$ Using the points (2014, 80) and (2000, 30), the slope is

$m = \frac{80-30}{2014-2000} = \frac{50}{14} = \frac{25}{7}$. The equation is $y - 80 = \frac{25}{7}(x - 2014)$. The difference between the two

equations is given as $h = \frac{30}{7}(x - 2014) + 100 - \frac{25}{7}(x - 2014) - 80 = \frac{5}{7}(x - 2014) + 20$

110. (a) Graph $h(x) = \frac{1900(x-1982)^2 + 619}{3200(x-1982)^2 + 1586}$, in the window [1982, 1994] by [0, 1]. See Figure 11-a.

Approximately 59% of the people who contracted AIDS during this time period died.

- (b) Divide number of deaths by number of cases for each year. The results compare favorably with the graph. See Figure 110b.

Xscl = 1 Yscl = 1




Figure 110a

Year	1982	1984	1986	1988	1990	1992	1994
Ratio	.39	.51	.59	.58	.61	.60	.61

Figure 110b

Reviewing Basic Concepts (Sections 2.4–2.6)

1. (a) $\left| \frac{1}{2}x + 2 \right| = 4 \Rightarrow \frac{1}{2}x + 2 = 4 \Rightarrow \frac{1}{2}x = 2 \Rightarrow x = 4$ or $\frac{1}{2}x + 2 = -4 \Rightarrow \frac{1}{2}x = -6 \Rightarrow x = -12$.

Therefore, the solution set is $\{-12, 4\}$.

- (b) $\left| \frac{1}{2}x + 2 \right| > 4 \Rightarrow \frac{1}{2}x + 2 > 4 \Rightarrow \frac{1}{2}x > 2 \Rightarrow x > 4$ or $\frac{1}{2}x + 2 < -4 \Rightarrow \frac{1}{2}x < -6 \Rightarrow x < -12$. Therefore, the solution interval is $(-\infty, -12) \cup (4, \infty)$.

- (c) $\left| \frac{1}{2}x + 2 \right| \leq 4 \Rightarrow -4 \leq \frac{1}{2}x + 2 \leq 4 \Rightarrow -6 \leq \frac{1}{2}x \leq 2 \Rightarrow -12 \leq x \leq 4$.

Therefore, the solution interval is $[-12, 4]$.

2. For the graph of $y = |f(x)|$, we reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 2.




Figure 2




Figure 5a

$$3. \quad |2x+4| = |1-3x| \Rightarrow 2x+4 = 1-3x \Rightarrow 5x = -3 \Rightarrow x = -\frac{3}{5} \text{ or } 2x+4 = -(1-3x) \Rightarrow 2x+4 = 3x-1 \Rightarrow 5 = x$$

The solution set is $\left\{-\frac{3}{5}, 5\right\}$.

$$4. \quad (a) \quad f(-3) = 2(-3) + 3 = -3 \quad (b) \quad f(0) = (0)^2 + 4 = 4 \quad (c) \quad f(2) = (2)^2 + 4 = 8$$

5. (a) See Figure 5a.

(b) Graph $y_1 = (-x^2) * (x \leq 0) + (x-4) * (x > 0)$ in the window $[-10, 10]$ by $[-10, 10]$. See Figure 5b.




Figure 5b

$$6. \quad (a) \quad (f+g)(x) = (-3x-4) + (x^2) = x^2 - 3x - 4. \text{ Therefore, } (f+g)(1) = (1)^2 - 3(1) - 4 = -6.$$

$$(b) \quad (f-g)(x) = (-3x-4) - (x^2) = -x^2 - 3x - 4. \text{ Therefore, } (f-g)(3) = -(3)^2 - 3(3) - 4 = -22.$$

$$(c) \quad (fg)(x) = (-3x-4)(x^2) = -3x^3 - 4x^2. \text{ Therefore, } (fg)(1) = -3(-2)^3 - 4(-2)^2 = 24 - 16 = 8.$$

$$(d) \quad \left(\frac{f}{g}\right)(x) = \frac{-3x-4}{x^2}. \text{ Therefore, } \left(\frac{f}{g}\right)(-3) = \frac{-3(-3)-4}{(-3)^2} = \frac{5}{9}.$$

$$(e) \quad (f \circ g)(x) = f[g(x)] = -3(x)^2 - 4 \Rightarrow (f \circ g)(x) = -3x^2 - 4$$

$$(f) \quad (g \circ f)(x) = g[f(x)] = (-3x-4)^2 \Rightarrow (g \circ f)(x) = 9x^2 + 24x + 16$$

$$7. \quad (a) \quad (f \circ g)(x) = f[g(x)] = \sqrt{4-x} \quad D: (-\infty, 4]$$

$$(b) \quad (g \circ f)(x) = g[f(x)] = 4 - \sqrt{x} \quad D: [0, \infty)$$

8. One of many possible solutions for $(f \circ g)(x) = h(x)$ is $f(x) = x^4$ and $g(x) = x + 2$. Then

$$(f \circ g)(x) = f[g(x)] = (x+2)^4.$$

$$9. \quad \frac{-2(x+h)^2 + 3(x+h) - 5 - (-2x^2 + 3x - 5)}{h} = \frac{-2(x^2 + 2xh + h^2) + 3x + 3h - 5 + 2x^2 - 3x + 5}{h} =$$

$$\frac{-2x^2 - 4xh - 2h^2 + 3x + 3h - 5 + 2x^2 - 3x + 5}{h} = \frac{-4xh - 2h^2 + 3h}{h} = -4x - 2h + 3.$$

10. (a) At 4% simple interest the equation for interest earned is $y_1 = .04x$.

(b) If he invested x dollars in the first account, then he invested $x+500$ in the second account. The equation for the amount of interest earned on this account is $y_2 = .025(x+500) \Rightarrow y_2 = .025x + 12.5$.

(c) It represents the total interest earned in both accounts for 1 year.

(d) Graph $y_1 + y_2 = .04x + (.025x + 12.5) \Rightarrow y_1 + y_2 = .04x + .025x + 12.5$ in the window $[0,1000]$ by $[0,100]$. See Figure 10. An input value of $x = 250$, results in \$28.75 earned interest.

(e) At $x = 250$, $y_1 + y_2 = .04(250) + .025(250) + 12.5 = 10 + 6.25 + 12.5 = \28.75 .




Figure 10

11. If the radius is r , then the height is $2r$ and the equation is

$$S = \pi r \sqrt{r^2 + (2r)^2} = \pi r \sqrt{r^2 + 4r^2} = \pi r \sqrt{5r^2} \Rightarrow S = \pi r^2 \sqrt{5}.$$

Chapter 2 Review Exercises

The graphs for exercises 1–10 can be found in the “Function Capsule” boxes located in section 2.1 in the text.

1. True Both $f(x) = x^2$ and $f(x) = |x|$ have the interval: $[0, \infty)$ as the range.
2. True Both $f(x) = x^2$ and $f(x) = |x|$ increase on the interval: $[0, \infty)$.
3. False The function $f(x) = \sqrt{x}$ has the domain: $[0, \infty)$ and $f(x) = \sqrt[3]{x}$ the domain: $(-\infty, \infty)$.
4. False The function $f(x) = \sqrt[3]{x}$ increases on its entire domain.
5. True The function $f(x) = x$ has a domain and range of: $(-\infty, \infty)$.
6. False The function $f(x) = \sqrt{x}$ is not defined on $(-\infty, 0)$, so certainly cannot be continuous.
7. True All of the functions show increases on the interval: $[0, \infty)$.
8. True Both $f(x) = x$ and $f(x) = x^3$ have graphs that are symmetric with respect to the origin.
9. True Both $f(x) = x^2$ and $f(x) = |x|$ have graphs that are symmetric with respect to the y -axis.
10. True No graphs are symmetric with respect to the x -axis.
11. Only values where $x \geq 0$ can be input for x , therefore the domain of $f(x) = \sqrt{x}$ is: $[0, \infty)$.
12. Only positive solution are possible in absolute value functions, therefore the range of $f(x) = \sqrt{x}$ is: $[0, \infty)$.
13. All solution are possible in cube root functions, therefore the range of $f(x) = \sqrt[3]{x}$ is: $(-\infty, \infty)$.
14. All values can be input for x , therefore the domain of $f(x) = x^2$ is: $(-\infty, \infty)$.
15. The function $f(x) = \sqrt[3]{x}$ increases for all inputs for x , therefore the interval is: $(-\infty, \infty)$.
16. The function $f(x) = |x|$ increases for all inputs where $x \geq 0$, therefore the interval is: $[0, \infty)$.
17. The equation is the equation $y = \sqrt{x}$. Only values where $x \geq 0$ can be input for x , therefore the domain of $y = \sqrt{x}$ is: $[0, \infty)$.

18. The equation $y^2 = x$ is the equation $y = \sqrt{x}$. Square root functions have both positive and negative solutions and all solutions are possible, therefore the range of $y = \sqrt{x}$ is: $(-\infty, \infty)$.
19. The graph of $f(x) = (x+3) - 1$ is the graph $y = x$ shifted 3 units to the left and 1 unit downward. See Figure 19.
20. The graph of $f(x) = -\frac{1}{2}x + 1$ is the graph reflected $y = x$ across the x -axis, vertically shrunk by a factor of $\frac{1}{2}$, and shifted 1 unit upward. See Figure 20.
21. The graph of $f(x) = (x+1)^2 - 2$ is the graph $y = x^2$ shifted 1 unit to the left and 2 units downward. See Figure 21.




Figure 19




Figure 20




Figure 21

22. The graph of $f(x) = -2x^2 + 3$ is the graph $y = x^2$ reflected across the x -axis, vertically stretched by a factor of 2, and shifted 3 units upward. See Figure 22.
23. The graph of $f(x) = -x^3 + 2$ is the graph $y = x^3$ reflected across the x -axis and shifted 2 units upward. See Figure 23.
24. The graph of $f(x) = (x-3)^3$ is the graph $y = x^3$ shifted 3 units to the right. See Figure 24.




Figure 22




Figure 23




Figure 24

25. The graph of $f(x) = \sqrt{\frac{1}{2}x}$ is the graph $y = \sqrt{x}$ horizontally stretched by a factor of 2. See Figure 25.
26. The graph of $f(x) = \sqrt{x-2} + 1$ is the graph $y = \sqrt{x}$ shifted 2 units to the right and 1 unit upward. See Figure 26.

27. The graph of $f(x) = 2\sqrt[3]{x}$ is the graph $y = \sqrt[3]{x}$ vertically stretched by a factor of 2. See Figure 27.




Figure 25




Figure 26




Figure 27

28. The graph of $f(x) = \sqrt[3]{x} - 2$ is the graph $y = \sqrt[3]{x}$ shifted 2 units downward. See Figure 28.
29. The graph of $f(x) = |x - 2| + 1$ is the graph $y = |x|$ shifted 2 units right and 1 unit upward. See Figure 29.
30. The graph of $f(x) = |-2x + 3|$ is the graph $y = |x|$ horizontally shrunk by a factor of $\frac{1}{2}$, shifted $\left(\frac{1}{2}\right)(3)$ or $\frac{3}{2}$ units to the left, and reflected across the y -axis. See Figure 30.




Figure 28




Figure 29




Figure 30

31. (a) From the graph, the function is continuous for the intervals: $(-\infty, -2)$, $[-2, 1]$ and $(1, \infty)$
- (b) From the graph, the function is increasing for the interval: $[-2, 1]$
- (c) From the graph, the function is decreasing for the interval: $(-\infty, -2)$
- (d) From the graph, the function is constant for the interval: $(1, \infty)$
- (e) From the graph, all values can be input for x , therefore the domain is: $(-\infty, \infty)$
- (f) From the graph, the possible values of y or the range is: $\{-2\} \cup [-1, 1] \cup (2, \infty)$
32. $x = y^2 - 4 \Rightarrow y^2 = x + 4 \Rightarrow y = \sqrt{x + 4}$ and $y = -\sqrt{x + 4}$.
33. From the graph, the relation is symmetric with respect to the x -axis, y -axis, and origin. The relation is not a function since some inputs x have two outputs y .

34. If $F(x) = x^3 - 6$, then $F(-x) = (-x)^3 - 6 \Rightarrow F(-x) = -x^3 - 6$ and $-F(x) = -(x^3 - 6) \Rightarrow -F(x) = -x^3 + 6$.
Since $F(x) \neq F(-x) \neq -F(x)$ the function has no symmetry and is neither an even nor an odd function.
35. If $f(x) = |x| + 4$, then $f(-x) = |-x| + 4 \Rightarrow f(-x) = |x| + 4$ and $-f(x) = -|x| - 4$. Since $f(-x) = f(x)$ the function is symmetric with respect to the y -axis and is an even function.
36. If $f(x) = \sqrt{x-5}$ then $f(-x) = \sqrt{-x-5}$ and $-f(x) = -\sqrt{x-5}$. Since $f(x) \neq f(-x) \neq -f(x)$, the function has no symmetry and is neither an even nor an odd function.
37. If $y^2 = x - 5$ then $y = \pm\sqrt{x-5}$. Since $f(x) = -\sqrt{x-5}$ is the reflection of $f(x) = \sqrt{x-5}$ across the x -axis, the relation has symmetry with respect to the x -axis. Also, one x inputs can produce two y outputs the relation is not a function.
38. If $f(x) = 3x^4 + 2x^2 + 1$ then $f(-x) = 3(-x)^4 + 2(-x)^2 + 1 \Rightarrow f(-x) = 3x^4 + 2x^2 + 1$ and $-f(x) = -3x^4 - 2x^2 - 1$. Since $f(-x) = f(x)$ the function is symmetric with respect to the y -axis and is an even function.
39. True, a graph that is symmetrical with respect to the x -axis means that for every (x, y) there is also $(x, -y)$.
40. True, since an even function and one that is symmetric with respect to the y -axis both contain the points (x, y) and $(-x, y)$.
41. True, since an odd function and one that is symmetric with respect to the origin both contain the points (x, y) and $(-x, -y)$.
42. False, for an even function, if (a, b) is on the graph, then $(-a, b)$ is on the graph and not $(a, -b)$. For example, $f(x) = x^2$ is even, and $(2, 4)$ is on the graph, but $(2, -4)$ is not.
43. False, for an odd function, if (a, b) is on the graph, then $(-a, -b)$ is on the graph and not $(-a, b)$. For example, $f(x) = x^3$ is odd, and $(2, 8)$ is on the graph, but $(-2, 8)$ is not.
44. True, if $(x, 0)$ is on the graph of $f(x) = 0$ then $(-x, 0)$ is on the graph.
45. The graph of $y = -3(x+4)^2 - 8$ is the graph of $y = x^2$ shifted 4 units to the left, vertically stretched by a factor of 3, reflected across the x -axis, and shifted 8 units downward.
46. The equation $y = \sqrt{x}$ reflected across the y -axis is: $f(x) = \sqrt{-x}$ then reflected across the x -axis is:
 $y = -\sqrt{-x}$ now vertically shrunk by a factor of $\frac{2}{3}$ is: $y = -\frac{2}{3}\sqrt{-x}$, and finally shifted 4 units upward is:
 $y = -\frac{2}{3}\sqrt{-x} + 4$.
47. Shift the function f upward 3 units. See Figure 47.

48. Shift the function f to the right 2 units. See Figure 48.
49. Shift the function f to the left 3 units and downward 2 units. See Figure 49.




Figure 47




Figure 48




Figure 49

50. For values where $f(x) > 0$ the graph remains the same. For values where $f(x) < 0$ reflect the graph across the x -axis. See Figure 50.
51. Horizontally shrink the function f by a factor of $\frac{1}{4}$. See Figure 51.
52. Horizontally stretch the function f by a factor of 2. See Figure 52.




Figure 50




Figure 51




Figure 52

53. The function is shifted upward 4 units, therefore the domain remains the same: $[-3, 4]$ and the range is increased by 4 and is: $[2, 9]$.
54. The function is shifted left 10 units, therefore the domain is decreased by 10 and is $[-13, -6]$; and the function is stretched vertically by a factor of 5, therefore the range is multiplied by 5 and is: $[-10, 25]$.
55. The function is horizontally shrunk by a factor of $\frac{1}{2}$, therefore the domain is divided by 2 and is: $[-\frac{3}{2}, 2]$; and the function is reflected across the x -axis, therefore the range is opposites of the original and is: $[-5, 2]$.
56. The function is shifted right 1 unit, therefore the domain is increased by 1 and is: $[-2, 5]$; and the function is also shifted upward 3 units, therefore the range is increased by 3 and is: $[1, 8]$.
57. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 57.

58. We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y < 0$. Where $y \geq 0$, the graph remains unchanged. See Figure 58.
59. Since the range is $\{2\}, y \geq 0$, so the graph remains unchanged.
60. Since the range is $\{-2\}, y < 0$, so we reflect the graph across the x -axis. See Figure 60.




Figure 57




Figure 58




Figure 60

61. $|4x+3|=12 \Rightarrow 4x+3=12 \Rightarrow 4x=9 \Rightarrow x=\frac{9}{4}$ or $4x+3=-12 \Rightarrow 4x=-15 \Rightarrow x=-\frac{15}{4}$, therefore the solution set is: $\left\{-\frac{15}{4}, \frac{9}{4}\right\}$.
62. $|-2x-6|+4=1 \Rightarrow |-2x-6|=-3$. Since an absolute value equation can not have a solution less than zero, the solution set is: $\emptyset$
63. $|5x+3|=|x+11| \Rightarrow 5x+3=x+11 \Rightarrow 4x=8 \Rightarrow x=2$ or $5x+3=-(x+11) \Rightarrow 6x=-14 \Rightarrow x=-\frac{14}{6}=-\frac{7}{3}$, therefore the solution set is: $\left\{-\frac{7}{3}, 2\right\}$.
64. $|2x+5|=7 \Rightarrow 2x+5=7 \Rightarrow 2x=2 \Rightarrow x=1$ or $2x+5=-7 \Rightarrow 2x=-12 \Rightarrow x=-6$, therefore the solution set is: $\{-6, 1\}$.
65. $|2x+5| \leq 7 \Rightarrow -7 \leq 2x+5 \leq 7 \Rightarrow -12 \leq 2x \leq 2 \Rightarrow -6 \leq x \leq 1$, therefore the interval is: $[-6, 1]$.
66. $|2x+5| \geq 7 \Rightarrow 2x+5 \geq 7 \Rightarrow 2x \geq 2 \Rightarrow x \geq 1$ or $2x+5 \leq -7 \Rightarrow 2x \leq -12 \Rightarrow x \leq -6$, therefore the solution is the interval: $(-\infty, -6] \cup [1, \infty)$.
67. $|5x-12| > 0 \Rightarrow 5x-12 > 0 \Rightarrow 5x > 12 \Rightarrow x > \frac{12}{5}$ or $5x-12 < 0 \Rightarrow 5x < 12 \Rightarrow x < \frac{12}{5}$, therefore the solution is the interval: $\left(-\infty, \frac{12}{5}\right) \cup \left(\frac{12}{5}, \infty\right)$ or $\left\{x \mid x \neq \frac{12}{5}\right\}$.
68. Since an absolute value equation can not have a solution less than zero, the solution set is: $\emptyset$

69. $2|3x-1|+1=21 \Rightarrow 2|3x-1|=20 \Rightarrow |3x-1|=10 \Rightarrow 3x-1=10 \Rightarrow 3x=11 \Rightarrow x=\frac{11}{3}$ or

$$3x-1=-10 \Rightarrow 3x=-9 \Rightarrow x=-3, \text{ therefore the solution set is: } \left\{-3, \frac{11}{3}\right\}.$$

70. $|2x+1|=|-3x+1| \Rightarrow 2x+1=-3x+1 \Rightarrow 5x=0 \Rightarrow x=0$ or $2x+1=-(-3x+1) \Rightarrow -x=-2 \Rightarrow x=2$, therefore the solution set is: $\{0, 2\}$.

71. The x -coordinates of the points of intersection of the graphs are -6 and 1 . Thus, $\{-6, 1\}$ is the solution set of $y_1 = y_2$. The graph of y_1 lies on or below the graph of y_2 between -6 and 1 , so the solution set of $y_1 \leq y_2$ is $[-6, 1]$. The graph of y_1 lies above the graph of y_2 everywhere else, so the solution set of $y_1 \geq y_2$ is $(-\infty, -6] \cup [1, \infty)$.

72. Graph $y_1 = |x+1| + |x-3|$ and $\{-3, -1\}$. See Figure 72. The intersections are $x = -3$ and $x = 5$, therefore the solution set is: $\{-3, 5\}$. **Check:** $|(-3)+1| + |(-3)-3| = 8 \Rightarrow |-2| + |-6| = 8 \Rightarrow 2+6=8 \Rightarrow 8=8$ and $|(5)+1| + |(5)-3| = 8 \Rightarrow |6| + |2| = 8 \Rightarrow 6+2=8 \Rightarrow 8=8$




Figure 72

73. Initially, the car is at home. After traveling 30 mph for 1 hr, the car is 30 mi away from home. During the second hour the car travels 20 mph until it is 50 mi away. During the third hour the car travels toward home at 30 mph until it is 20 mi away. During the fourth hour the car travels away from home at 40 mph until it is 60 mi away from home. During the last hour, the car travels 60 mi at 60 mph until it arrives home.

74. See Figure 74

75. See Figure 75

76. See Figure 76




Figure 74




Figure 75




Figure 76

77. Graph $y_1 = (3x+1)*(x < 2) + (-x+4)*(x \geq 2)$ in the window $[-10, 10]$ by

$$|4x+8| > 4 \Rightarrow 4x+8 > 4 \Rightarrow 4x > -4 \Rightarrow -1 \text{ See Figure 77.}$$

78. See Figure 78.




Figure 77




Figure 78

79. From the graphs $(f+g)(1) = 2+3 = 5$

80. From the graphs $(f-g)(0) = 1-4 = -3$

81. From the graphs $(fg)(-1) = (0)(3) = 0$

82. From the graphs $\left(\frac{f}{g}\right)(2) = \frac{3}{2}$

83. From the graphs $(f \circ g)(2) = f[g(2)] = f(2) = 3$

84. From the graphs $(g \circ f)(2) = g[f(2)] = g(3) = 2$

85. From the graphs $(g \circ f)(-4) = g[f(-4)] = g(2) = 2$

86. From the graphs $(f \circ g)(-2) = f[g(-2)] = f(2) = 3$

87. From the table $(f+g)(1) = 7+1 = 8$

88. From the table $(f-g)(3) = 9-1 = 0$

89. From the table $(fg)(-1) = (3)(-2) = -6$

90. From the table $\left(\frac{f}{g}\right)(0) = \frac{5}{0}$, which is undefined.

91. From the tables $(g \circ f)(-2) = g[f(-2)] = g(1) = 2$

92. From the graphs $(f \circ g)(3) = f[g(3)] = f(-2) = 1$

93. $\frac{2(x+h)+9-(2x+9)}{h} = \frac{2x+2h+9-2x-9}{h} = \frac{2h}{h} = 2$
94. $\frac{(x+h)^2-5(x+h)+3-(x^2-5x+3)}{h} = \frac{x^2+2xh+h^2-5x-5h+3-x^2+5x-3}{h} = \frac{2xh+h^2-5h}{h} = 2x+h-5$
95. One of many possible solutions for $(f \circ g)(x) = h(x)$ is: $f(x) = x^2$ and $g(x) = x^3 - 3x$. Then
 $(f \circ g)(x) = (x^3 - 3x)^2$.
96. One of many possible solutions for $(f \circ g)(x) = h(x)$ is $f(x) = \frac{1}{x}$ and $g(x) = x - 5$. Then
 $(f \circ g)(x) = f[g(x)] = \frac{1}{x-5}$
97. If $V(r) = \frac{4}{3}\pi r^3$, then a 4 inch increase would be: $V(r) = \frac{4}{3}\pi(r+4)^3$, and the volume gained would be:
 $V(r) = \frac{4}{3}\pi(r+4)^3 - \frac{4}{3}\pi r^3$.
98. (a) Since $h = d, r = \frac{d}{2}$ and the formula for the volume of a can is: $V = \pi r^2 h$, the function is:
 $V(d) = \pi \left(\frac{d}{2}\right)^2 d \Rightarrow V(d) = \frac{\pi d^3}{4}$
- (b) Since $h = d, r = \frac{d}{2}, c = 2\pi r$ and the formula for the surface area of a can is: $A = 2\pi r h + 2\pi r^2$, the
function is: $S(d) = 2\pi \left(\frac{d}{2}\right) d + 2\pi \left(\frac{d}{2}\right)^2 \Rightarrow S(d) = \pi d^2 + \frac{\pi d^2}{2} \Rightarrow S(d) = \frac{3\pi d^2}{2}$
99. The function for changing yards to inches is: $f(x) = 36x$ and the function for changing miles to yards is:
 $g(x) = 1760x$. The composition of this which would change miles into inches is:
 $f[g(x)] = 36[1760(x)] \Rightarrow (f \circ g)(x) = 63,360x$.
100. If $x = \text{width}$, then $\text{length} = 2x$. A formula for Perimeter can now be written as: $P = x + 2x + x + 2x$ and the
function is: $P(x) = 6x$. This is a linear function.

Chapter 2 Test

1. (a) D, only values where $x \geq 0$ can be input into a square root function.
- (b) D, only values where $y \geq 0$ can be the solution to a square root function.
- (c) C, all values can be input for x in a squaring function.
- (d) B, only values where $y \geq 3$ can be the solution to $f(x) = x^2 + 3$
- (e) C, all values can be input for x in a cube root function.
- (f) C, all values can be a solution in a cube root function.
- (g) C, all values can be input for x in an absolute value function.
- (h) D, only values where $y \geq 0$ can be the solution to an absolute value function.
- (i) D, if $x = y^2$ then $y = \sqrt{x}$ and only values where $x \geq 0$ can be input into a square root function.

- (j) C, all values can be a solution in this function.
2. (a) This is $f(x)$ shifted 2 units upward. See Figure 2a.
- (b) This is $f(x)$ shifted 2 units to the left. See Figure 2b.
- (c) This is $f(x)$ reflected across the x -axis. See Figure 2c.
- (d) This is $f(x)$ reflected across the y -axis. See Figure 2d.
- (e) This is $f(x)$ vertically stretched by a factor of 2. See Figure 2e.
- (f) We reflect the graph of $y = f(x)$ across the x -axis for all points for which $y \geq 0$. Where $y < 0$ the graph remains unchanged. See Figure 2f.




Figure 2a




Figure 2b




Figure 2c




Figure 2d




Figure 2e




Figure 2f

3. (a) Since $y = f(2x)$ is $y = f(x)$ horizontally shrunk by a factor of $\frac{1}{2}$, the point $(-2, 4)$ on $y = f(x)$ becomes the point $(-1, 4)$ on the graph of $y = f(2x)$.
- (b) Since $y = f\left(\frac{1}{2}x\right)$ is $y = f(x)$ horizontally stretched by a factor of 2, the point $(-2, 4)$ on $y = f(x)$ becomes the point $(-4, 4)$ on the graph of $y = f\left(\frac{1}{2}x\right)$.
4. (a) The graph of $f(x) = -x(x-2)^2 + 4$ is the basic graph $f(x) = x^2$ reflected across the x -axis, shifted 2 units to the right, and shifted 4 units upward. See Figure 4a.
- (b) The graph of $f(x) = -2\sqrt{-x}$ is the basic graph $f(x) = \sqrt{x}$ reflected across the y -axis and vertically stretched by a factor of 2. See Figure 4b.




Figure 4a




Figure 4b

5. (a) If the graph is symmetric with respect to the y -axis, then $(x, y) \Rightarrow (-x, y)$ therefore $(3, 6) \Rightarrow (-3, 6)$.
- (b) If the graph is symmetric with respect to the x -axis, then $(x, y) \Rightarrow (x, -y)$ therefore $(3, 6) \Rightarrow (3, -6)$.
- (c) See Figure 5. We give an actual screen here. The drawing should resemble it.




Figure 5

6. (a) Shift the graph of $y = \sqrt[3]{x}$ to the left 2 units, vertically stretch by a factor of 4, and shift 5 units downward.
- (b) Graph $y = |x|$ reflected across the x -axis, vertically shrunk by a factor of $\frac{1}{2}$ shifted 3 units to the right, and shifted up 2 units. See Figure 6. From the graph the domain is: $(-\infty, \infty)$; and the range is: $(-\infty, 2]$.




Figure 6

7. (a) From the graph, the function is increasing for the interval: $(-\infty, -3)$
- (b) From the graph, the function is decreasing for the interval: $(4, \infty)$
- (c) From the graph, the function is constant for the interval: $(-3, 4)$
- (d) From the graph, the function is continuous for the intervals: $(-\infty, -3), (-3, 4), (4, \infty)$.
- (e) From the graph, the domain is: $(-\infty, \infty)$
- (f) From the graph, the range is: $(-\infty, 2)$

8. (a) $|4x+8|=4 \Rightarrow 4x+8=4 \Rightarrow 4x=-4 \Rightarrow x=-1$ or $4x+8=-4 \Rightarrow 4x=-12 \Rightarrow x=-3$, therefore the solution set is: $\{-3, -1\}$
- (b) $|4x+8|<4 \Rightarrow -4<4x+8<4 \Rightarrow -12<4x<-4 \Rightarrow -3<x<-1$, therefore the solution is: $(-3, -1)$.
- (c) $|4x+8|>4 \Rightarrow 4x+8>4 \Rightarrow 4x>-4 \Rightarrow x>-1$ or $4x+8<-4 \Rightarrow 4x<-12 \Rightarrow x<-3$ therefore the solution is: $(-\infty, -3) \cup (-1, \infty)$.
9. (a) $(f-g)(x) = 2x^2 - 3x + 2 - (-2x+1) \Rightarrow (f-g)(x) = 2x^2 - x + 1$
- (b) $\left(\frac{f}{g}\right)(x) = \frac{2x^2 - 3x + 2}{-2x+1}$
- (c) The domain can be all values for x , except any that make $g(x)=0$. Therefore
 $-2x+1 \neq 0 \Rightarrow -2x \neq -1 \Rightarrow x \neq \frac{1}{2}$ or the interval: $\left(-\infty, \frac{1}{2}\right) \cup \left(\frac{1}{2}, \infty\right)$
- (d) $(f \circ g)(x) = f[g(x)] = 2(-2x+1)^2 - 3(-2x+1) + 2 = 2(4x^2 - 4x + 1) + 6x - 3 + 2 =$
 $8x^2 - 8x + 2 + 6x - 3 + 2 = 8x^2 - 2x + 1$
- (e) $(g \circ f)(x) = g[f(x)] = -2(2x^2 - 3x + 2) + 1 = -4x^2 + 6x - 4 + 1 = -4x^2 + 6x - 3$
- (f) $\frac{2(x+h)^2 - 3(x+h) + 2 - (2x^2 - 3x + 2)}{h} = \frac{2(x^2 + 2xh + h^2) - 3x - 3h + 2 - 2x^2 + 3x - 2}{h} =$
 $\frac{2x^2 + 4xh + 2h^2 - 3x - 3h + 2 - 2x^2 + 3x - 2}{h} = \frac{4xh + 2h^2 - 3h}{h} = 4x + 2h - 3$
10. (a) $(f \circ g)(x) = f[g(x)] = 2\sqrt{x-1} + 1$ D: $[1, \infty)$
- (b) $(g \circ f)(x) = g[f(x)] = \sqrt{2x+1-1} = \sqrt{2x}$ D: $[0, \infty)$
11. (a) See Figure 11a.
- (b) Graph $y_1 = (-x^2 + 3) * (x \leq 1) + (\sqrt[3]{x} + 2) * (x > 1)$ in the window $[-4.7, 4.7]$ by $[-5.1, 5.1]$
- See Figure 11b.
- (c) The graph is not connected at $x = 1$ and thus f is not continuous when $x = 1$.




Figure 11a

$[-4.7, 4.7]$ by $[-5.1, 5.1]$

Xscl = 1 Yscl = 1




Figure 11b

 $[0, 10]$ by $[0, 6]$

Xscl = 1 Yscl = 1




Figure 12

 $[0, 1,000]$ by $[-4,000, 4,000]$

Xscl = 50 Yscl = 500





Figure 13

12. (a) See Figure 12.
- (b) Set $x = 5.5$ then \$2.75 is the cost of a 5.5 minute international call. See the display at the bottom of the screen.
13. (a) With an initial set-up cost of \$3300 and a production cost of \$4.50 the function is: $C(x) = 3300 + 4.50x$
- (b) With a selling price of \$10.50 the revenue function is: $R(x) = 10.50x$
- (c) $P(x) = R(x) - C(x) \Rightarrow P(x) = 10.50x - (3300 + 4.50x) \Rightarrow P(x) = 6x - 3300$
- (d) To make a profit $P(x) > 0$, therefore $6x - 3300 > 0 \Rightarrow 6x > 3300 \Rightarrow x > 550$
Tyler needs to sell 551 before he earns a profit.
- (e) Graph $y_1 = 6x - 3300$, See Figure 13. The first integer x -value for which $P(x) > 0$ is 551

Chapter R

Review: Basic Algebra Concepts



R.2 Review of Exponents and Polynomials



Review of Exponents and Polynomials

Product Rule

For all positive integers m and n and every real number a ,

$$a^m \cdot a^n = a^{m+n}.$$



Example: Using the Product Rule

Find each product.

a) $y^4 \cdot y^7$

b) $(6z^5)(9z^3)(2z^2)$

Solution

a) $y^4 \cdot y^7 = y^{4+7} = y^{11}$

b) $(6z^5)(9z^3)(2z^2) = (6 \cdot 9 \cdot 2)(z^5 z^3 z^2)$
 $= 108z^{5+3+2} = 108z^{10}$



Review of Exponents

Zero Exponent

For any nonzero real number a ,

$$a^0 = 1.$$

NOTE The expression 0^0 is undefined.



Example: Using the Definition of a^0

Evaluate each power.

a) 4^0

b) $(-4)^0$

c) -4^0

Solution

a) $4^0 = 1$

b) $(-4)^0 = 1$

c) $-4^0 = -1$ since $-4^0 = -(4)^0 = -1$.



Reviewing Exponents and Polynomials

Power Rules

For all positive integers m and n and all real numbers a and b ,

$$(a^m)^n = a^{mn}, \quad (ab)^m = a^m b^m,$$

$$\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m} \quad (b \neq 0).$$



Example: Using the Power Rules

Simplify each expression.

a) $(5^3)^2$

b) $(3^4 x^2)^3$

c) $\left(\frac{2^5}{b^4}\right)^3$, if $b \neq 0$

a) $(5^3)^2 = 5^{3(2)} = 5^6$

b) $(3^4 x^2)^3 = (3^4)^3 (x^2)^3 = 3^{4(3)} x^{2(3)} = 3^{12} x^6$

c) $\left(\frac{2^5}{b^4}\right)^3 = \frac{(2^5)^3}{(b^4)^3} = \frac{2^{15}}{b^{12}}$, if $b \neq 0$

Terminology for Polynomials (1 of 4)

An **algebraic expression** results from adding, subtracting, multiplying, dividing (except by 0), or finding roots or powers of any combination of variables and constants.

Examples include

$$2x^2 - 3x, \frac{5y}{2y-3}, \sqrt{m^3-8}, (3a+b)^4$$



Terminology for Polynomials (2 of 4)

A **term** is the product of a real number and one or more variables raised to powers.

The real number in a term is called the **numerical coefficient**, or just the **coefficient**.

In the term $-3m^4$, the coefficient is -3 .



Terminology for Polynomials (3 of 4)

Like terms are terms with the same variables raised to the same powers.

$-13x^3$, $4x^3$, and $-x^3$ are like terms,

$6y$ and $6y^2$ are not.



Terminology for Polynomials (4 of 4)

A **polynomial** is a term or a finite sum of terms with only nonnegative integer exponents permitted on the variables.

$$5x^3 - 8x^2 + 7x - 4 ,$$

$$9p^5 - 3 ,$$

$$8r^2 ,$$

$$6$$

are all polynomials.



Adding and Subtracting Polynomials

Polynomials are added by adding coefficients of like terms.

Polynomials are subtracted by subtracting coefficients of like terms.



Example: Adding and Subtracting Polynomials (1 of 2)

Add or subtract, as indicated.

$$\begin{aligned}\text{a)} \quad & (2y^4 - 3y^2 + y) + (4y^4 + 7y^2 + 6y) \\ &= (2 + 4)y^4 + (-3 + 7)y^2 + (1 + 6)y = 6y^4 + 4y^2 + 7y\end{aligned}$$

$$\begin{aligned}\text{b)} \quad & (-3m^3 - 8m^2 + 4) - (m^3 + 7m^2 - 3) \\ &= (-3 - 1)m^3 + (-8 - 7)m^2 + [4 - (-3)] \\ &= -4m^3 - 15m^2 + 7\end{aligned}$$

Example: Adding and Subtracting Polynomials (2 of 2)

Add or subtract, as indicated.

$$\begin{aligned}\text{c) } 8m^4 p^5 - 9m^3 p^5 + (11m^4 p^5 + 15m^3 p^5) \\ = 19m^4 p^5 + 6m^3 p^5\end{aligned}$$

$$\begin{aligned}\text{d) } 4(x^2 - 3x + 7) - 5(2x^2 - 8x - 4) \\ = 4x^2 - 4(3x) + 4(7) - 5(2x^2) - 5(-8x) - 5(-4) \\ = 4x^2 - 12x + 28 - 10x^2 + 40x + 20 \\ = -6x^2 + 28x + 48\end{aligned}$$

Multiplying Polynomials

The associative and distributive properties, together with the properties of exponents, can be used to multiply polynomials.

Polynomials can be multiplied by multiplying each term of the first polynomial by each term of the second, then combining like terms.



Example: Multiplying Polynomials

Multiply $(3p^2 - 4p + 1)(p^3 + 2p - 8)$.

Solution

$$\begin{array}{r} 3p^2 - 4p + 1 \\ \quad p^3 + 2p - 8 \\ \hline -24p^2 + 32p - 8 \\ 6p^3 - 8p^2 + 2p \\ \hline 3p^5 - 4p^4 + p^3 \\ \hline 3p^5 - 4p^4 + 7p^3 - 32p^2 + 34p - 8 \end{array}$$



Multiply Polynomials

A **binomial** is a polynomial with two terms.

The FOIL method can be used to multiply two binomials.

The FOIL method says to multiply the First terms, the Outside terms, the Inside terms, and the Last terms, then combine like terms.



Example: Using FOIL Method (1 of 2)

Find each product.

a) $(6m + 1)(4m - 3)$

F O I L

$$\begin{aligned}(6m + 1)(4m - 3) &= 6m(4m) + 6m(-3) + 1(4m) + 1(-3) \\ &= 24m^2 - 14m - 3\end{aligned}$$



Example: Using FOIL Method (2 of 2)

Find each product.

$$\text{b) } (2x + 7)(2x - 7) = 4x^2 - 14x + 14x - 49 = 4x^2 - 49$$

$$\begin{aligned}\text{c) } r^2(3r + 2)(3r - 2) &= r^2(9r^2 - 6r + 6r - 4) \\ &= r^2(9r^2 - 4) \\ &= 9r^4 - 4r^2\end{aligned}$$



Review of Exponents and Polynomials

Special Products

Product of the Sum and Difference of Two Terms

$$(x + y)(x - y) = x^2 - y^2$$

Square of a Binomial

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$



Example: Using the Special Products

Find each product.

$$\text{a) } (3p + 11)(3p - 11) = (3p)^2 - 11^2 = 9p^2 - 121$$

$$\text{b) } (5m^3 - 3)(5m^3 + 3) = (5m^3)^2 - 3^2 = 25m^6 - 9$$

$$\text{c) } (2m + 5)^2 = (2m)^2 + 2(2m)(5) + (5)^2 = 4m^2 + 20m + 25$$

$$\begin{aligned}\text{d) } (3x - 7y^4)^2 &= (3x)^2 - 2(3x)(7y^4) + (7y^4)^2 \\ &= 9x^2 - 42xy^4 + 49y^8\end{aligned}$$

