

***Probability and Statistical Inference 10th edition
Hogg Solutions Manual***

SKU: 9780135189399-SOLUTIONS

INSTRUCTOR'S SOLUTIONS MANUAL


PROBABILITY AND STATISTICAL INFERENCE TENTH EDITION

Robert V. Hogg

Elliot A. Tanis

Dale L. Zimmerman





The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2020, 2015, 2010 Pearson Education, Inc.
Publishing as Pearson, 330 Hudson Street, NY NY 10013

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-518948-1
ISBN-10: 0-13-518948-9

Contents

Preface	v
1 Probability	1
1.1 Properties of Probability	1
1.2 Methods of Enumeration	2
1.3 Conditional Probability	3
1.4 Independent Events	4
1.5 Bayes' Theorem	5
2 Discrete Distributions	7
2.1 Random Variables of the Discrete Type	7
2.2 Mathematical Expectation	9
2.3 Special Mathematical Expectations	11
2.4 The Binomial Distribution	14
2.5 The Hypergeometric Distribution	16
2.6 The Negative Binomial Distribution	17
2.7 The Poisson Distribution	18
3 Continuous Distributions	19
3.1 Random Variables of the Continuous Type	19
3.2 The Exponential, Gamma, and Chi-Square Distributions	26
3.3 The Normal Distribution	28
3.4 Additional Models	30
4 Bivariate Distributions	33
4.1 Bivariate Distributions of the Discrete Type	33
4.2 The Correlation Coefficient	34
4.3 Conditional Distributions	36
4.4 Bivariate Distributions of the Continuous Type	37
4.5 The Bivariate Normal Distribution	41
5 Distributions of Functions of Random Variables	45
5.1 Functions of One Random Variable	45
5.2 Transformations of Two Random Variables	47
5.3 Several Random Variables	52
5.4 The Moment-Generating Function Technique	53
5.5 Random Functions Associated with Normal Distributions	55
5.6 The Central Limit Theorem	58
5.7 Approximations for Discrete Distributions	59
5.8 Chebyshev's Inequality and Convergence in Probability	61
5.9 Limiting Moment-Generating Functions	62

6	Point Estimation	63
6.1	Descriptive Statistics	63
6.2	Exploratory Data Analysis	65
6.3	Order Statistics	70
6.4	Maximum Likelihood Estimation	73
6.5	A Simple Regression Problem	76
6.6	Asymptotic Distributions of Maximum Likelihood Estimators	81
6.7	Sufficient Statistics	81
6.8	Bayesian Estimation	84
7	Interval Estimation	87
7.1	Confidence Intervals for Means	87
7.2	Confidence Intervals for the Difference of Two Means	88
7.3	Confidence Intervals For Proportions	90
7.4	Sample Size	91
7.5	Distribution-Free Confidence Intervals for Percentiles	92
7.6	More Regression	93
7.7	Resampling Methods	99
8	Tests of Statistical Hypotheses	107
8.1	Tests About One Mean	107
8.2	Tests of the Equality of Two Means	109
8.3	Tests for Variances	111
8.4	Tests about Proportions	113
8.5	Some Distribution-Free Tests	114
8.6	Power of a Statistical Test	118
8.7	Best Critical Regions	121
8.8	Likelihood Ratio Tests	124
9	More Tests	127
9.1	Chi-Square Goodness-of-Fit Tests	127
9.2	Contingency Tables	130
9.3	One-Factor Analysis of Variance	131
9.4	Two-Way Analysis of Variance	134
9.5	General Factorial and 2^k Factorial Designs	135
9.6	Tests Concerning Regression and Correlation	136
9.7	Statistical Quality Control	137

Preface

This solutions manual provides answers for the even-numbered exercises in *Probability and Statistical Inference*, tenth edition, by Robert V. Hogg, Elliot A. Tanis, and Dale L. Zimmerman. Complete solutions are given for most of these exercises. You, the instructor, may decide how many of these solutions and answers you want to make available to your students. Note that the answers for the odd-numbered exercises are given in the textbook. Our hope is that this solutions manual will be helpful to each of you in your teaching.

All of the figures in this manual were generated using *Maple*, a computer algebra system. Most of the figures were generated and many of the solutions, especially those involving data, were solved using procedures that were written by Zaven Karian from Denison University. We thank him for providing these. These procedures are available free of charge for your use. They are available for download at <http://www.math.hope.edu/tanis/>. Short descriptions of these procedures are provided on the “Maple Card.” Complete descriptions of these procedures are given in *Probability and Statistics: Explorations with MAPLE*, second edition, 1999, written by Zaven Karian and Elliot Tanis, published by Prentice Hall (ISBN 0-13-021536-8). You can download a slightly revised edition of this manual at <http://www.math.hope.edu/tanis/MapleManual.pdf>.

We also want to acknowledge the many suggestions/corrections that were made by our accuracy checker, Kyle Siegrist.

If you find an error or wish to make a suggestion, please send them to dale-zimmerman@uiowa.edu. These **errata** will be posted on <http://homepage.divms.uiowa.edu/~dzimmer/>.

E.A.T.
D.L.Z.

Chapter 1

Probability

1.1 Properties of Probability

1.1-2 Sketch a figure and fill in the probabilities of each of the disjoint sets.

Let $A = \{\text{insure more than one car}\}$, $P(A) = 0.85$.

Let $B = \{\text{insure a sports car}\}$, $P(B) = 0.23$.

Let $C = \{\text{insure exactly one car}\}$, $P(C) = 0.15$.

It is also given that $P(A \cap B) = 0.17$. Since $A \cap C = \phi$, $P(A \cap C) = 0$. It follows that $P(A \cap B \cap C') = 0.17$. Thus $P(A' \cap B \cap C') = 0.06$ and $P(B' \cap C) = 0.09$.

1.1-4 (a) $S = \{\text{HHHH, HHHT, HHTH, HTHH, THHH, HHTT, HTTH, TTHH, HTHT, THTH, THHT, HTTT, THTT, TTHT, TTTH, TTTT}\}$;

(b) (i) $5/16$, (ii) 0 , (iii) $11/16$, (iv) $4/16$, (v) $4/16$, (vi) $9/16$, (vii) $4/16$.

1.1-6 (a) $P(A \cup B) = 0.5 + 0.6 - 0.4 = 0.7$;

$$\begin{aligned} \text{(b)} \quad A &= (A \cap B') \cup (A \cap B) \\ P(A) &= P(A \cap B') + P(A \cap B) \\ 0.5 &= P(A \cap B') + 0.4 \\ P(A \cap B') &= 0.1; \end{aligned}$$

$$\text{(c)} \quad P(A' \cup B') = P[(A \cap B)'] = 1 - P(A \cap B) = 1 - 0.4 = 0.6.$$

1.1-8 Let $A = \{\text{lab work done}\}$, $B = \{\text{referral to a specialist}\}$,

$$P(A) = 0.41, P(B) = 0.53, P[(A \cup B)'] = 0.21.$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$0.79 = 0.41 + 0.53 - P(A \cap B)$$

$$P(A \cap B) = 0.41 + 0.53 - 0.79 = 0.15.$$

$$\text{1.1-10} \quad A \cup B \cup C = A \cup (B \cup C)$$

$$P(A \cup B \cup C) = P(A) + P(B \cup C) - P[A \cap (B \cup C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - P[(A \cap B) \cup (A \cap C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - P(A \cap B) - P(A \cap C) + P(A \cap B \cap C).$$

1.1-12 (a) $1/3$; (b) $2/3$; (c) 0 ; (d) $1/2$.

$$\mathbf{1.1-14} \quad P(A) = \frac{2[r - r(\sqrt{3}/2)]}{2r} = 1 - \frac{\sqrt{3}}{2}.$$

1.1-16 Note that the respective probabilities are p_0 , $p_1 = p_0/4$, $p_2 = p_0/4^2, \dots$.

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{p_0}{4^k} &= 1 \\ \frac{p_0}{1 - 1/4} &= 1 \\ p_0 &= \frac{3}{4} \\ 1 - p_0 - p_1 &= 1 - \frac{15}{16} = \frac{1}{16}. \end{aligned}$$

1.2 Methods of Enumeration

1.2-2 (a) $(4)(5)(2) = 40$; (b) $(2)(2)(2) = 8$.

$$\mathbf{1.2-4} \quad (\mathbf{a}) \quad 4 \binom{6}{3} = 80;$$

$$(\mathbf{b}) \quad 4(2^6) = 256;$$

$$(\mathbf{c}) \quad \frac{(4-1+3)!}{(4-1)!3!} = 20.$$

1.2-6 $S = \{ \text{DDD, DDFD, DFDD, FDDD, DFFD, DFDF, FDDF, DFFD, FDFD, FFDD, FFF, FFDF, FDF, DFFF, FFDD, FDF, DFFD, FDDF, DFDF, DFFF} \}$ so there are 20 possibilities. Note that the winning player (2 choices) must win the last set and two of the previous sets, so the number of outcomes is

$$2 \left[\binom{2}{2} + \binom{3}{2} + \binom{4}{2} \right] = 20.$$

$$\mathbf{1.2-8} \quad 3 \cdot 3 \cdot 2^{12} = 36,864.$$

$$\begin{aligned} \mathbf{1.2-10} \quad \binom{n-1}{r} + \binom{n-1}{r-1} &= \frac{(n-1)!}{r!(n-1-r)!} + \frac{(n-1)!}{(r-1)!(n-r)!} \\ &= \frac{(n-r)(n-1)! + r(n-1)!}{r!(n-r)!} = \frac{n!}{r!(n-r)!} = \binom{n}{r}. \end{aligned}$$

$$\mathbf{1.2-12} \quad 0 = (1-1)^n = \sum_{r=0}^n \binom{n}{r} (-1)^r (1)^{n-r} = \sum_{r=0}^n (-1)^r \binom{n}{r}.$$

$$2^n = (1+1)^n = \sum_{r=0}^n \binom{n}{r} (1)^r (1)^{n-r} = \sum_{r=0}^n \binom{n}{r}.$$

$$\mathbf{1.2-14} \quad \binom{5-1+29}{29} = \frac{33!}{29!4!} = 40,920.$$

$$\mathbf{1.2-16} \quad (\mathbf{a}) \quad \frac{\binom{19}{3} \binom{52-19}{6}}{\binom{52}{9}} = \frac{102,486}{351,325} = 0.2917;$$

$$(b) \frac{\binom{19}{3}\binom{10}{2}\binom{7}{1}\binom{3}{0}\binom{5}{1}\binom{2}{0}\binom{6}{2}}{\binom{52}{9}} = \frac{7,695}{1,236,664} = 0.00622.$$

- 1.2-18** (a) $P(A) = \sum_{n=1}^5 (1/2)^n = 1 - (1/2)^5;$
 (b) $P(B) = \sum_{n=1}^{10} (1/2)^n = 1 - (1/2)^{10};$
 (c) $P(A \cup B) = P(B) = 1 - (1/2)^{10};$
 (d) $P(A \cap B) = P(A) = 1 - (1/2)^5;$
 (e) $P(C) = P(B) - P(A) = (1/2)^5 - (1/2)^{10};$
 (f) $P(B') = 1 - P(B) = (1/2)^{10}.$

1.3 Conditional Probability

1.3-2 (a) $\frac{1041}{1456};$

(b) $\frac{392}{633};$

(c) $\frac{649}{823}.$

- (d) The proportion of women who favor a gun law is greater than the proportion of men who favor a gun law.

1.3-4 (a) $P(HH) = \frac{13}{52} \cdot \frac{12}{51} = \frac{1}{17};$

(b) $P(HC) = \frac{13}{52} \cdot \frac{13}{51} = \frac{13}{204};$

(c) $P(\text{Non-Ace Heart, Ace}) + P(\text{Ace of Hearts, Non-Heart Ace})$
 $= \frac{12}{52} \cdot \frac{4}{51} + \frac{1}{52} \cdot \frac{3}{51} = \frac{51}{52 \cdot 51} = \frac{1}{52}.$

- 1.3-6** Let $H = \{\text{died from heart disease}\}; P = \{\text{at least one parent had heart disease}\}.$

$$P(H | P') = \frac{N(H \cap P')}{N(P')} = \frac{110}{648}.$$

1.3-8 (a) $\frac{3}{20} \cdot \frac{2}{19} \cdot \frac{1}{18} = \frac{1}{1140};$

(b) $\frac{\binom{3}{2}\binom{17}{1}}{\binom{20}{3}} \cdot \frac{1}{17} = \frac{1}{380};$

(c) $\sum_{k=1}^9 \frac{\binom{3}{2}\binom{17}{2k-2}}{\binom{20}{2k}} \cdot \frac{1}{20-2k} = \frac{35}{76} = 0.4605;$

(d) Draw second. The probability of winning is $1 - 0.4605 = 0.5395$.

$$\mathbf{1.3-10} \quad (\mathbf{a}) \quad P(A) = \frac{52}{52} \cdot \frac{51}{52} \cdot \frac{50}{52} \cdot \frac{49}{52} \cdot \frac{48}{52} \cdot \frac{47}{52} = \frac{8,808,975}{11,881,376} = 0.74141;$$

$$(\mathbf{b}) \quad P(A') = 1 - P(A) = 0.25859.$$

$$\mathbf{1.3-12} \quad (\mathbf{a}) \quad \text{It doesn't matter because } P(B_1) = \frac{1}{18}, \quad P(B_5) = \frac{1}{18}, \quad P(B_{18}) = \frac{1}{18};$$

$$(\mathbf{b}) \quad P(B) = \frac{2}{18} = \frac{1}{9} \quad \text{on each draw.}$$

$$\mathbf{1.3-14} \quad (\mathbf{a}) \quad 5 \cdot 4 \cdot 3 = 60;$$

$$(\mathbf{b}) \quad 5 \cdot 5 \cdot 5 = 125.$$

$$\mathbf{1.3-16} \quad \frac{3}{5} \cdot \frac{5}{8} + \frac{2}{5} \cdot \frac{4}{8} = \frac{23}{40}.$$

1.4 Independent Events

$$\begin{aligned} \mathbf{1.4-2} \quad (\mathbf{a}) \quad P(A \cap B) &= P(A)P(B) = (0.3)(0.6) = 0.18; \\ P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= 0.3 + 0.6 - 0.18 \\ &= 0.72; \end{aligned}$$

$$(\mathbf{b}) \quad P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{0.6} = 0.$$

$$\begin{aligned} \mathbf{1.4-4} \quad \text{Proof of (b):} \quad P(A' \cap B) &= P(B)P(A'|B) \\ &= P(B)[1 - P(A|B)] \\ &= P(B)[1 - P(A)] \\ &= P(B)P(A'). \end{aligned}$$

$$\begin{aligned} \text{Proof of (c):} \quad P(A' \cap B') &= P[(A \cup B)'] \\ &= 1 - P(A \cup B) \\ &= 1 - P(A) - P(B) + P(A \cap B) \\ &= 1 - P(A) - P(B) + P(A)P(B) \\ &= [1 - P(A)][1 - P(B)] \\ &= P(A')P(B'). \end{aligned}$$

$$\begin{aligned} \mathbf{1.4-6} \quad P[A \cap (B \cap C)] &= P[A \cap B \cap C] \\ &= P(A)P(B)P(C) \\ &= P(A)P(B \cap C). \end{aligned}$$

$$\begin{aligned} P[A \cap (B \cup C)] &= P[(A \cap B) \cup (A \cap C)] \\ &= P(A \cap B) + P(A \cap C) - P(A \cap B \cap C) \\ &= P(A)P(B) + P(A)P(C) - P(A)P(B)P(C) \\ &= P(A)[P(B) + P(C) - P(B \cap C)] \\ &= P(A)P(B \cup C). \end{aligned}$$

$$\begin{aligned} P[A' \cap (B \cap C')] &= P(A' \cap C' \cap B) \\ &= P(B)[P(A' \cap C') | B] \\ &= P(B)[1 - P(A \cup C | B)] \\ &= P(B)[1 - P(A \cup C)] \\ &= P(B)P[(A \cup C)'] \\ &= P(B)P(A' \cap C') \\ &= P(B)P(A')P(C') \\ &= P(A')P(B)P(C') \\ &= P(A')P(B \cap C'). \end{aligned}$$

$$\begin{aligned}
P[A' \cap B' \cap C'] &= P[(A \cup B \cup C)'] \\
&= 1 - P(A \cup B \cup C) \\
&= 1 - P(A) - P(B) - P(C) + P(A)P(B) + P(A)P(C) + \\
&\quad P(B)P(C) - P(A)P(B)P(C) \\
&= [1 - P(A)][1 - P(B)][1 - P(C)] \\
&= P(A')P(B')P(C').
\end{aligned}$$

$$1.4-8 \quad \frac{1}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} + \frac{1}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} + \frac{5}{6} \cdot \frac{2}{6} \cdot \frac{3}{6} = \frac{2}{9}.$$

$$1.4-10 \quad (a) \quad \frac{3}{4} \cdot \frac{3}{4} = \frac{9}{16};$$

$$(b) \quad \frac{1}{4} \cdot \frac{3}{4} + \frac{3}{4} \cdot \frac{2}{4} = \frac{9}{16};$$

$$(c) \quad \frac{2}{4} \cdot \frac{1}{4} + \frac{2}{4} \cdot \frac{4}{4} = \frac{10}{16}.$$

$$1.4-12 \quad (a) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(b) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(c) \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2;$$

$$(d) \quad \frac{5!}{3!2!} \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^2.$$

$$1.4-14 \quad (a) \quad 1 - (0.4)^3 = 1 - 0.064 = 0.936;$$

$$(b) \quad 1 - (0.4)^8 = 1 - 0.00065536 = 0.99934464.$$

$$1.4-16 \quad (a) \quad \sum_{k=0}^{\infty} \frac{1}{5} \left(\frac{4}{5}\right)^{2k} = \frac{5}{9};$$

$$(b) \quad \frac{1}{5} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{1}{3} + \frac{4}{5} \cdot \frac{3}{4} \cdot \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{1}{1} = \frac{3}{5}.$$

$$1.4-18 \quad (a) \quad 7; (b) \quad (1/2)^7; (c) \quad 63; (d) \quad \text{No! } (1/2)^{63} = 1/9,223,372,036,854,775,808.$$

1.4-20 No. The equations that must hold are

$$(1 - p_1)(1 - p_2) = p_1(1 - p_2) + p_2(1 - p_1) = p_1p_2.$$

There are no real solutions.

1.5 Bayes' Theorem

$$\begin{aligned}
1.5-2 \quad (a) \quad P(G) &= P(A \cap G) + P(B \cap G) \\
&= P(A)P(G|A) + P(B)P(G|B) \\
&= (0.40)(0.85) + (0.60)(0.75) = 0.79;
\end{aligned}$$

$$\begin{aligned}
(b) \quad P(A|G) &= \frac{P(A \cap G)}{P(G)} \\
&= \frac{(0.40)(0.85)}{0.79} = 0.43.
\end{aligned}$$

1.5-4 Let event B denote an accident and let A_1 be the event that age of the driver is 16–25. Then

$$\begin{aligned} P(A_1 | B) &= \frac{(0.1)(0.05)}{(0.1)(0.05) + (0.55)(0.02) + (0.20)(0.03) + (0.15)(0.04)} \\ &= \frac{50}{50 + 110 + 60 + 60} = \frac{50}{280} = 0.179. \end{aligned}$$

1.5-6 Let B be the event that the policyholder dies. Let A_1, A_2, A_3 be the events that the deceased is standard, preferred and ultra-preferred, respectively. Then

$$\begin{aligned} P(A_1 | B) &= \frac{(0.60)(0.01)}{(0.60)(0.01) + (0.30)(0.008) + (0.10)(0.007)} \\ &= \frac{60}{60 + 24 + 7} = \frac{60}{91} = 0.659; \\ P(A_2 | B) &= \frac{24}{91} = 0.264; \\ P(A_3 | B) &= \frac{7}{91} = 0.077. \end{aligned}$$

1.5-8 Let A be the event that the tablet is under warranty.

$$\begin{aligned} P(B_1 | A) &= \frac{(0.40)(0.10)}{(0.40)(0.10) + (0.30)(0.05) + (0.20)(0.03) + (0.10)(0.02)} \\ &= \frac{40}{40 + 15 + 6 + 2} = \frac{40}{63} = 0.635; \\ P(B_2 | A) &= \frac{15}{63} = 0.238; \\ P(B_3 | A) &= \frac{6}{63} = 0.095; \\ P(B_4 | A) &= \frac{2}{63} = 0.032. \end{aligned}$$

1.5-10 (a) $P(D^+) = (0.02)(0.92) + (0.98)(0.05) = 0.0184 + 0.0490 = 0.0674$;

(b) $P(A^- | D^+) = \frac{0.0490}{0.0674} = 0.727$; $P(A^+ | D^+) = \frac{0.0184}{0.0674} = 0.273$;

(c) $P(A^- | D^-) = \frac{(0.98)(0.95)}{(0.02)(0.08) + (0.98)(0.95)} = \frac{9310}{16 + 9310} = 0.998$;
 $P(A^+ | D^-) = 0.002$;

(d) Yes, particularly those in part (b).

1.5-12 Let $D = \{\text{defective roll}\}$. Then

$$\begin{aligned} P(I | D) &= \frac{P(I \cap D)}{P(D)} \\ &= \frac{P(I) \cdot P(D | I)}{P(I) \cdot P(D | I) + P(II) \cdot P(D | II)} \\ &= \frac{(0.60)(0.03)}{(0.60)(0.03) + (0.40)(0.01)} \\ &= \frac{0.018}{0.018 + 0.004} = \frac{0.018}{0.022} = 0.818. \end{aligned}$$

Chapter 2

Discrete Distributions

2.1 Random Variables of the Discrete Type

2.1-2 (a)

$$f(x) = \begin{cases} 0.6, & x = 1, \\ 0.3, & x = 5, \\ 0.1, & x = 10; \end{cases}$$

(b)




Figure 2.1-2: A probability histogram

$$\begin{aligned}
 \text{2.1-4 (a)} \quad \sum_{x=1}^9 \log_{10} \left(\frac{x+1}{x} \right) &= \sum_{x=1}^9 [\log_{10}(x+1) - \log_{10} x] \\
 &= \log_{10} 2 - \log_{10} 1 + \log_{10} 3 - \log_{10} 2 + \cdots + \log_{10} 10 - \log_{10} 9 \\
 &= \log_{10} 10 = 1;
 \end{aligned}$$

(b)

$$F(x) = \begin{cases} 0, & x < 1, \\ \log_{10} n, & n-1 \leq x < n, \quad n = 2, 3, \dots, 9, \\ 1, & 9 \leq x. \end{cases}$$

2.1-6 (a) $f(x) = \frac{1}{10}, \quad x = 0, 1, 2, \dots, 9;$

(b) $\mathcal{N}(\{0\})/150 = 11/150 = 0.073; \quad \mathcal{N}(\{5\})/150 = 13/150 = 0.087;$
 $\mathcal{N}(\{1\})/150 = 14/150 = 0.093; \quad \mathcal{N}(\{6\})/150 = 22/150 = 0.147;$
 $\mathcal{N}(\{2\})/150 = 13/150 = 0.087; \quad \mathcal{N}(\{7\})/150 = 16/150 = 0.107;$
 $\mathcal{N}(\{3\})/150 = 12/150 = 0.080; \quad \mathcal{N}(\{8\})/150 = 18/150 = 0.120;$
 $\mathcal{N}(\{4\})/150 = 16/150 = 0.107; \quad \mathcal{N}(\{9\})/150 = 15/150 = 0.100.$

(c)




Figure 2.1-6: Michigan daily lottery digits

2.1-8 (a) $f(x) = \frac{6 - |7 - x|}{36}, \quad x = 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12;$

(b)




Figure 2.1-8: Probability histogram for the sum of a pair of dice

2.1-10 (a) The space of W is $S = \{0, 1, 2, 3, 4, 5, 6, 7\}$.

$$P(W = 0) = P(X = 0, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}, \text{ assuming independence.}$$

$$P(W = 1) = P(X = 0, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 2) = P(X = 2, Y = 0) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 3) = P(X = 2, Y = 1) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 4) = P(X = 0, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 5) = P(X = 0, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 6) = P(X = 2, Y = 4) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8},$$

$$P(W = 7) = P(X = 2, Y = 5) = \frac{1}{2} \cdot \frac{1}{4} = \frac{1}{8}.$$

$$\text{That is, } f(w) = P(W = w) = \frac{1}{8}, \quad w \in S.$$

(b)




Figure 2.1-10: Probability histogram of sum of two special dice

2.1-12 Let x equal the number of orange balls and $144 - x$ the number of blue balls. Then

$$\begin{aligned} \frac{x}{144} \cdot \frac{x-1}{143} + \frac{144-x}{144} \cdot \frac{143-x}{143} &= \frac{x}{144} \cdot \frac{144-x}{143} + \frac{144-x}{144} \cdot \frac{x}{143} \\ x^2 - x + 144 \cdot 143 - 144x - 143x + x^2 &= 2 \cdot 144x - 2 \cdot x^2 \\ x^2 - 144x + 5,148 &= 0 \\ (x-78)(x-66) &= 0 \end{aligned}$$

Thus there are 78 orange balls and 66 blue balls.

2.2 Mathematical Expectation

$$\mathbf{2.2-2} \quad E(X) = (-1)\left(\frac{4}{9}\right) + (0)\left(\frac{1}{9}\right) + (1)\left(\frac{4}{9}\right) = 0;$$

$$E(X^2) = (-1)^2\left(\frac{4}{9}\right) + (0)^2\left(\frac{1}{9}\right) + (1)^2\left(\frac{4}{9}\right) = \frac{8}{9};$$

$$E(3X^2 - 2X + 4) = 3\left(\frac{8}{9}\right) - 2(0) + 4 = \frac{20}{3}.$$

$$\begin{aligned} \mathbf{2.2-4} \quad 1 &= \sum_{x=0}^6 f(x) = \frac{9}{10} + c\left(\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6}\right) \\ c &= \frac{2}{49}; \\ E(\text{Payment}) &= \frac{2}{49}\left(1 \cdot \frac{1}{2} + 2 \cdot \frac{1}{3} + 3 \cdot \frac{1}{4} + 4 \cdot \frac{1}{5} + 5 \cdot \frac{1}{6}\right) = \frac{71}{490} \text{ units.} \end{aligned}$$

2.2-6 Note that $\sum_{x=1}^{\infty} \frac{6}{\pi^2 x^2} = \frac{6}{\pi^2} \sum_{x=1}^{\infty} \frac{1}{x^2} = \frac{6}{\pi^2} \frac{\pi^2}{6} = 1$, so this is a pdf.

$$E(X) = \sum_{x=1}^{\infty} x \frac{6}{\pi^2 x^2} = \frac{6}{\pi^2} \sum_{x=1}^{\infty} \frac{1}{x} = +\infty$$

and it is well known that the sum of this harmonic series is not finite.

2.2-8 $E(|X - c|) = \frac{1}{7} \sum_{x \in S} |x - c|$, where $S = \{1, 2, 3, 5, 15, 25, 50\}$.

When $c = 5$,

$$E(|X - 5|) = \frac{1}{7} [(5 - 1) + (5 - 2) + (5 - 3) + (5 - 5) + (15 - 5) + (25 - 5) + (50 - 5)].$$

If c is either increased or decreased by 1, this expectation is increased by $1/7$. Thus $c = 5$, the median, minimizes this expectation while $b = E(X) = 101/7$ minimizes $E[(X - b)^2]$.

2.2-10 $(1) \cdot \frac{15}{36} + (-1) \cdot \frac{21}{36} = \frac{-6}{36} = \frac{-1}{6};$

$$(1) \cdot \frac{15}{36} + (-1) \cdot \frac{21}{36} = \frac{-6}{36} = \frac{-1}{6};$$

$$(4) \cdot \frac{6}{36} + (-1) \cdot \frac{30}{36} = \frac{-6}{36} = \frac{-1}{6}.$$

2.2-12 (a) The average class size is $\frac{(16)(25) + (3)(100) + (1)(300)}{20} = 50;$

(b)

$$f(x) = \begin{cases} 0.4, & x = 25, \\ 0.3, & x = 100, \\ 0.3, & x = 300; \end{cases}$$

(c) $E(X) = 25(0.4) + 100(0.3) + 300(0.3) = 130.$

2.3 Special Mathematical Expectations

2.3-2 (a)

$$\begin{aligned}
 \mu &= E(X) \\
 &= \sum_{x=1}^3 x \frac{3!}{x!(3-x)!} \left(\frac{1}{4}\right)^x \left(\frac{3}{4}\right)^{3-x} \\
 &= 3 \left(\frac{1}{4}\right) \sum_{k=0}^2 \frac{2!}{k!(2-k)!} \left(\frac{1}{4}\right)^k \left(\frac{3}{4}\right)^{2-k} \\
 &= 3 \left(\frac{1}{4}\right) \left(\frac{1}{4} + \frac{3}{4}\right)^2 = \frac{3}{4}; \\
 E[X(X-1)] &= \sum_{x=2}^3 x(x-1) \frac{3!}{x!(3-x)!} \left(\frac{1}{4}\right)^x \left(\frac{3}{4}\right)^{3-x} \\
 &= 2(3) \left(\frac{1}{4}\right)^2 \frac{3}{4} + 6 \left(\frac{1}{4}\right)^3 \\
 &= 6 \left(\frac{1}{4}\right)^2 = 2 \left(\frac{1}{4}\right) \left(\frac{3}{4}\right); \\
 \sigma^2 &= E[X(X-1)] + E(X) - \mu^2 \\
 &= (2) \left(\frac{3}{4}\right) \left(\frac{1}{4}\right) + \left(\frac{3}{4}\right) - \left(\frac{3}{4}\right)^2 \\
 &= (2) \left(\frac{3}{4}\right) \left(\frac{1}{4}\right) + \left(\frac{3}{4}\right) \left(\frac{1}{4}\right) = 3 \left(\frac{1}{4}\right) \left(\frac{3}{4}\right);
 \end{aligned}$$

(b) $\mu = E(X)$

$$\begin{aligned}
 &= \sum_{x=1}^4 x \frac{4!}{x!(4-x)!} \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{4-x} \\
 &= 4 \left(\frac{1}{2}\right) \sum_{k=0}^3 \frac{3!}{k!(3-k)!} \left(\frac{1}{2}\right)^k \left(\frac{1}{2}\right)^{3-k} \\
 &= 4 \left(\frac{1}{2}\right) \left(\frac{1}{2} + \frac{1}{2}\right)^3 = 2; \\
 E[X(X-1)] &= \sum_{x=2}^4 x(x-1) \frac{4!}{x!(4-x)!} \left(\frac{1}{2}\right)^x \left(\frac{1}{2}\right)^{4-x} \\
 &= 2(6) \left(\frac{1}{2}\right)^4 + (6)(4) \left(\frac{1}{2}\right)^4 + (12) \left(\frac{1}{2}\right)^4 \\
 &= 48 \left(\frac{1}{2}\right)^4 = 12 \left(\frac{1}{2}\right)^2; \\
 \sigma^2 &= (12) \left(\frac{1}{2}\right)^2 + \frac{4}{2} - \left(\frac{4}{2}\right)^2 = 1.
 \end{aligned}$$

2.3-4 $E[(X - \mu)/\sigma] = (1/\sigma)[E(X) - \mu] = (1/\sigma)(\mu - \mu) = 0;$

$$E\{[(X - \mu)/\sigma]^2\} = (1/\sigma^2)E[(X - \mu)^2] = (1/\sigma^2)(\sigma^2) = 1.$$

2.3-6 $f(1) = \frac{3}{8}, f(2) = \frac{2}{8}, f(3) = \frac{3}{8},$

$$\mu = 1 \cdot \frac{3}{8} + 2 \cdot \frac{2}{8} + 3 \cdot \frac{3}{8} = 2,$$

$$\sigma^2 = 1^2 \cdot \frac{3}{8} + 2^2 \cdot \frac{2}{8} + 3^2 \cdot \frac{3}{8} - 2^2 = \frac{3}{4}.$$

$$\begin{aligned} \mathbf{2.3-8} \quad E(X) &= \sum_{x=1}^4 x \cdot \frac{2x-1}{16} \\ &= \frac{50}{16} = 3.125; \end{aligned}$$

$$\begin{aligned} E(X^2) &= \sum_{x=1}^4 x^2 \cdot \frac{2x-1}{16} \\ &= \frac{85}{8}; \end{aligned}$$

$$\text{Var}(X) = \frac{85}{8} - \left(\frac{25}{8}\right)^2 = \frac{55}{64} = 0.8594;$$

$$\sigma = \frac{\sqrt{55}}{8} = 0.9270.$$

$$\mathbf{2.3-10} \quad \mu = E(X) = (-5)(1/16) + (-1)(5/8) + (3)(5/16) = 0 \text{ so}$$

$$E[(X-\mu)^3] = E(X^3) = (-5)^3(1/16) + (-1)^3(5/8) + (3^3)(5/16) = (-125 - 10 + 135)/16 = 0.$$

Thus the index of skewness likewise equals 0. The distribution is not symmetric, however.

$$\mathbf{2.3-12} \quad (\mathbf{a}) \quad f(x) = \left(\frac{364}{365}\right)^{x-1} \left(\frac{1}{365}\right), \quad x = 1, 2, 3, \dots;$$

$$(\mathbf{b}) \quad \mu = \frac{1}{\frac{1}{365}} = 365,$$

$$\sigma^2 = \frac{364}{\left(\frac{1}{365}\right)^2} = 132,860,$$

$$\sigma = 364.500;$$

$$(\mathbf{c}) \quad P(X > 400) = \left(\frac{364}{365}\right)^{400} = 0.3337,$$

$$P(X < 300) = 1 - \left(\frac{364}{365}\right)^{299} = 0.5597.$$

$$\mathbf{2.3-14} \quad P(X \geq 100) = P(X > 99) = (0.99)^{99} = 0.3697.$$

$$\mathbf{2.3-16} \quad (\mathbf{a}) \quad f(x) = (1/2)^{x-1}, \quad x = 2, 3, 4, \dots;$$

$$\begin{aligned} (\mathbf{b}) \quad M(t) &= E[e^{tx}] = \sum_{x=2}^{\infty} e^{tx} (1/2)^{x-1} \\ &= 2 \sum_{x=2}^{\infty} (e^t/2)^x \\ &= \frac{2(e^t/2)^2}{1 - e^t/2} = \frac{e^{2t}}{2 - e^t}, \quad t < \ln 2; \end{aligned}$$

$$\begin{aligned}
\text{(c)} \quad M'(t) &= \frac{4e^{2t} - e^{3t}}{(2 - e^t)^2} \\
\mu &= M'(0) = 3; \\
M''(t) &= \frac{(2 - e^t)^2(8e^{2t} - 3e^{3t}) - (4e^{2t} - e^{3t})2 \cdot (2 - e^t)(-e^t)}{(2 - e^t)^4} \\
\sigma^2 &= M''(0) - \mu^2 = 11 - 9 = 2; \\
\text{(d)} \quad \text{(i)} \quad P(X \leq 3) &= 3/4; \quad \text{(ii)} \quad P(X \geq 5) = 1/8; \quad \text{(iii)} \quad P(X = 3) = 1/4.
\end{aligned}$$

$$\begin{aligned}
\text{2.3-18} \quad P(X > k + j | X > k) &= \frac{P(X > k + j)}{P(X > k)} \\
&= \frac{q^{k+j}}{q^k} = q^j = P(X > j).
\end{aligned}$$

2.3-20 (a) Construct a square in the first quadrant with lengths of sides equal to one. Within this square construct squares with one vertex at the origin and side lengths equal to $1 - 1/2^x$;

$$\begin{aligned}
\text{(b)} \quad f(x) &= \left(1 - \frac{1}{2^x}\right)^2 - \left(1 - \frac{1}{2^{x-1}}\right)^2 = \frac{(2^x - 1)^2}{2^{2x}} - \frac{(2^{x-1} - 1)^2}{2^{2x-2}} \\
&= \frac{(2^x - 1)^2}{2^{2x}} - \frac{2^2(2^{x-1} - 1)^2}{2^{2x}} = \frac{(2^x - 1)^2}{2^{2x}} - \frac{(2^x - 2)^2}{2^{2x}} \\
&= \frac{2^{2x} - 2 \cdot 2^x + 1 - 2^{2x} + 2^2 \cdot 2^x - 4}{2^{2x}} = \frac{2^{x+1} - 3}{2^{2x}}, \quad x = 1, 2, 3, \dots;
\end{aligned}$$

$$\text{(c)} \quad \sum_{x=1}^{\infty} \left[\frac{2^{x+1}}{2^{2x}} - \frac{3}{2^{2x}} \right] = \sum_{x=1}^{\infty} \left[\frac{2 \cdot 2^x}{2^{2x}} - \frac{3}{4^x} \right] = \frac{4/4}{1 - 1/2} - \frac{3/4}{1 - 1/4} = 2 - 1 = 1;$$

$$\text{(d)} \quad M(t) = E(e^{tX}) = \sum_{x=1}^{\infty} e^{tx} \cdot \frac{2^{x+1} - 3}{2^{2x}} = \sum_{x=1}^{\infty} \left[\frac{2(2e^t)^x}{4^x} - \frac{3(e^{tx})}{4^x} \right] = \frac{2e^t}{2 - e^t} - \frac{3e^t}{4 - e^t};$$

$$\begin{aligned}
\text{(e)} \quad M'(t) &= \frac{4e^t - 2e^{2t} + 2e^{2t}}{(2 - e^t)^2} - \frac{12e^t - 3e^{2t} + 3e^{2t}}{(4 - e^t)^2} \\
&= \frac{4e^t}{(2 - e^t)^2} - \frac{12e^t}{(4 - e^t)^2} \\
\mu &= M'(0) = 4 - \frac{12}{9} = \frac{8}{3};
\end{aligned}$$

$$\begin{aligned}
\text{(f)} \quad M''(t) &= \frac{(2 - e^t)^2 \cdot 4e^t - 4e^t - 4e^t \cdot 2(2 - e^t)(-e^t)}{(2 - e^t)^4} - \\
&\quad \frac{(4 - e^t)^2 \cdot 12e^t - 12e^t - 12e^t \cdot 2(4 - e^t)(-e^t)}{(4 - e^t)^4} \\
\sigma^2 &= M''(0) - \mu^2 = \frac{4 + 8}{1} - \frac{108 + 72}{81} - \left(\frac{8}{3}\right)^2 = \frac{8}{3}.
\end{aligned}$$

2.4 The Binomial Distribution

$$\mathbf{2.4-2} \quad f(-1) = \frac{11}{18}, \quad f(1) = \frac{7}{18};$$

$$\mu = (-1)\frac{11}{18} + (1)\frac{7}{18} = -\frac{4}{18};$$

$$\sigma^2 = \left(-1 + \frac{4}{18}\right)^2 \left(\frac{11}{18}\right) + \left(1 + \frac{4}{18}\right)^2 \left(\frac{7}{18}\right) = \frac{77}{81}.$$

The variance can also be computed by noting that $X^2 = 1$ so

$$\sigma^2 = 1 - \mu^2 = 1 - \left(-\frac{4}{18}\right)^2 = \frac{77}{81}.$$

$$\mathbf{2.4-4} \quad (\mathbf{a}) \quad X \text{ is } b(7, 0.15);$$

$$(\mathbf{b}) \quad (\mathbf{i}) \quad P(X \geq 2) = 1 - P(X \leq 1) = 1 - 0.7166 = 0.2834;$$

$$(\mathbf{ii}) \quad P(X = 1) = P(X \leq 1) - P(X \leq 0) = 0.7166 - 0.3206 = 0.3960;$$

$$(\mathbf{iii}) \quad P(X \leq 3) = 0.9879.$$

$$\mathbf{2.4-6} \quad (\mathbf{a}) \quad X \text{ is } b(15, 0.75); \quad 15 - X \text{ is } b(15, 0.25);$$

$$(\mathbf{b}) \quad P(X \geq 10) = P(15 - X \leq 5) = 0.8516;$$

$$(\mathbf{c}) \quad P(X \leq 10) = P(15 - X \geq 5) = 1 - P(15 - X \leq 4) = 1 - 0.6865 = 0.3135;$$

$$(\mathbf{d}) \quad P(X = 10) = P(X \geq 10) - P(X \geq 11) \\ = P(15 - X \leq 5) - P(15 - X \leq 4) = 0.8516 - 0.6865 = 0.1651;$$

$$(\mathbf{e}) \quad \mu = (15)(0.75) = 11.25, \quad \sigma^2 = (15)(0.75)(0.25) = 2.8125; \quad \sigma = \sqrt{2.8125} = 1.67705.$$

$$\mathbf{2.4-8} \quad (\mathbf{a}) \quad 1 - 0.01^4 = 0.99999999; \quad (\mathbf{b}) \quad 0.99^4 = 0.960596.$$

$$\mathbf{2.4-10} \quad (\mathbf{a}) \quad X \text{ is } b(8, 0.90);$$

$$(\mathbf{b}) \quad (\mathbf{i}) \quad P(X = 8) = P(8 - X = 0) = 0.4305;$$

$$(\mathbf{ii}) \quad P(X \leq 6) = P(8 - X \geq 2) \\ = 1 - P(8 - X \leq 1) = 1 - 0.8131 = 0.1869;$$

$$(\mathbf{iii}) \quad P(X \geq 6) = P(8 - X \leq 2) = 0.9619.$$

$$\mathbf{2.4-12} \quad (\mathbf{a})$$

$$f(x) = \begin{cases} 125/216, & x = -1, \\ 75/216, & x = 1, \\ 15/216, & x = 2, \\ 1/216, & x = 3; \end{cases}$$

$$(\mathbf{b}) \quad \mu = (-1) \cdot \frac{125}{216} + (1) \cdot \frac{75}{216} + (2) \cdot \frac{15}{216} + (3) \cdot \frac{1}{216} = -\frac{17}{216};$$

$$\sigma^2 = E(X^2) - \mu^2 = \frac{269}{216} - \left(-\frac{17}{216}\right)^2 = 1.2392;$$

$$\sigma = 1.11;$$

(c) See Figure 2.4-12.




Figure 2.4-12: Losses in chuck-a-luck

2.4-14 Let X equal the number of winning tickets when n tickets are purchased. Then

$$\begin{aligned} P(X \geq 1) &= 1 - P(X = 0) \\ &= 1 - \left(\frac{9}{10}\right)^n. \end{aligned}$$

(a) $1 - (0.9)^n = 0.50$

$$(0.9)^n = 0.50$$

$$n \ln 0.9 = \ln 0.5$$

$$n = \frac{\ln 0.5}{\ln 0.9} = 6.58$$

so $n = 7$;

(b) $1 - (0.9)^n = 0.95$

$$(0.9)^n = 0.05$$

$$n = \frac{\ln 0.05}{\ln 0.9} = 28.43$$

so $n = 29$.

2.4-16 It is given that X is $b(10, 0.10)$. We are to find M so that

$P(1000X \leq M) \geq 0.99$ or $P(X \leq M/1000) \geq 0.99$. From Appendix Table II, $P(X \leq 4) = 0.9984 > 0.99$. Thus $M/1000 = 4$ or $M = 4000$ dollars.

2.4-18 Let X equal the number of infected individuals. X is $b(5, 0.05)$. The expected number of tests is

$$1 P(X = 0) + 6 P(X > 0) = 1 (0.7738) + 6 (1 - 0.7738) = 2.131.$$

2.4-20 (a) (i) $b(5, 0.7)$; (ii) $\mu = 3.5, \sigma^2 = 1.05$; (iii) 0.1607;

(b) (i) geometric, $p = 0.3$; (ii) $\mu = 10/3, \sigma^2 = 70/9$; (iii) 0.51;

(c) (i) Bernoulli, $p = 0.55$; (ii) $\mu = 0.55, \sigma^2 = 0.2475$; (iii) 0.55;

- (d) (ii) $\mu = 2.1, \sigma^2 = 0.89$; (iii) 0.7;
 (e) (i) discrete uniform on $1, 2, \dots, 10$; (ii) 5.5, 8.25; (iii) 0.2.

2.5 The Hypergeometric Distribution

$$\text{2.5-2 (a)} \quad \frac{\binom{4}{4}\binom{4}{0}}{\binom{8}{4}} = \frac{1}{70};$$

$$\text{(b)} \quad 2 \frac{\binom{4}{4}\binom{4}{0}}{\binom{8}{4}} = \frac{2}{70}.$$

$$\text{2.5-4} \quad P(X \geq 1) = 1 - P(X = 0) = 1 - \frac{\binom{3}{0}\binom{17}{5}}{\binom{20}{5}} = 1 - \frac{91}{228} = \frac{137}{228} = 0.60.$$

2.5-6 We have $N = N_1 + N_2$. Thus

$$\begin{aligned} E[X(X-1)] &= \sum_{x=0}^n x(x-1)f(x) \\ &= \frac{\sum_{x=2}^n x(x-1) \frac{N_1!}{x!(N_1-x)!} \cdot \frac{N_2!}{(n-x)!(N_2-n+x)!}}{\binom{N}{n}} \\ &= N_1(N_1-1) \frac{\sum_{x=2}^n \frac{(N_1-2)!}{(x-2)!(N_1-x)!} \cdot \frac{N_2!}{(n-x)!(N_2-n+x)!}}{\binom{N}{n}}. \end{aligned}$$

In the summation, let $k = x - 2$, and in the denominator, note that

$$\binom{N}{n} = \frac{N!}{n!(N-n)!} = \frac{N(N-1)}{n(n-1)} \binom{N-2}{n-2}.$$

Thus

$$\begin{aligned} E[X(X-1)] &= \frac{\frac{N_1(N_1-1)}{N(N-1)} \sum_{k=0}^{n-2} \frac{\binom{N_1-2}{k} \binom{N_2}{n-2-k}}{\binom{N-2}{n-2}}}{\frac{n(n-1)}{N(N-1)}} \\ &= \frac{N_1(N_1-1)(n)(n-1)}{N(N-1)}. \end{aligned}$$

$$\text{2.5-8 (a)} \quad \frac{\binom{5}{5}\binom{65}{0}}{\binom{70}{5}} \cdot \frac{1}{25} = \frac{1}{302,575,350};$$

$$(b) \frac{\binom{5}{5} \binom{65}{0}}{\binom{70}{5}} \cdot \frac{24}{25} = \frac{24}{302,575,350};$$

$$(c) \frac{\binom{5}{4} \binom{65}{1}}{\binom{70}{5}} \cdot \frac{1}{25} = \frac{325}{302,575,350};$$

$$(d) \frac{\binom{5}{4} \binom{65}{1}}{\binom{70}{5}} \cdot \frac{24}{25} = \frac{7,800}{302,575,350};$$

$$(e) \frac{\binom{5}{0} \binom{65}{5}}{\binom{70}{5}} \cdot \frac{1}{25} = \frac{8,259,888}{302,575,350}.$$

2.5-10 (a) $P(2, 1, 6, 10)$ means that 2 is in position 1 so 1 cannot be selected. Thus

$$P(2, 1, 6, 10) = \frac{\binom{1}{0} \binom{1}{1} \binom{8}{5}}{\binom{10}{6}} = \frac{56}{210} = \frac{4}{15};$$

$$(b) P(i, r, k, n) = \frac{\binom{i-1}{r-1} \binom{1}{1} \binom{n-i}{k-r}}{\binom{n}{k}}.$$

2.6 The Negative Binomial Distribution

$$\mathbf{2.6-2} \quad \binom{10-1}{5-1} \left(\frac{1}{2}\right)^5 \left(\frac{1}{2}\right)^5 = \frac{126}{1024} = \frac{63}{512}.$$

2.6-4 Let “being missed” be a success and let X equal the number of trials until the first success. Then $p = 0.01$.

$$P(X \leq 50) = 1 - 0.99^{50} = 1 - 0.605 = 0.395.$$

$$\mathbf{2.6-6} \quad (a) \quad R(t) = \ln(1 - p + pe^t),$$

$$R'(t) = \left[\frac{pe^t}{1 - p + pe^t} \right]_{t=0} = p,$$

$$R''(t) = \left[\frac{(1 - p + pe^t)(pe^t) - (pe^t)(pe^t)}{(1 - p + pe^t)^2} \right]_{t=0} = p(1 - p);$$

$$(b) \quad R(t) = n \ln(1 - p + pe^t),$$

$$R'(t) = \left[\frac{npe^t}{1 - p + pe^t} \right]_{t=0} = np,$$

$$R''(t) = n \left[\frac{(1 - p + pe^t)(pe^t) - (pe^t)(pe^t)}{(1 - p + pe^t)^2} \right]_{t=0} = np(1 - p);$$

$$\begin{aligned}
\text{(c)} \quad R(t) &= \ln p + t - \ln[1 - (1-p)e^t], \\
R'(t) &= \left[1 + \frac{(1-p)e^t}{1 - (1-p)e^t} \right]_{t=0} = 1 + \frac{1-p}{p} = \frac{1}{p}, \\
R''(t) &= [(-1)\{1 - (1-p)e^t\}^2\{-(1-p)e^t\}]_{t=0} = \frac{1-p}{p^2}; \\
\text{(d)} \quad R(t) &= r [\ln p + t - \ln\{1 - (1-p)e^t\}], \\
R'(t) &= r \left[\frac{1}{1 - (1-p)e^t} \right]_{t=0} = \frac{r}{p}, \\
R''(t) &= r [(-1)\{1 - (1-p)e^t\}^{-2}\{-(1-p)e^t\}]_{t=0} = \frac{r(1-p)}{p^2}.
\end{aligned}$$

$$\mathbf{2.6-8} \quad (0.7)(0.7)(0.3) = 0.147.$$

2.6-10 (a) Let X equal the number of boxes that must be purchased. Then

$$E(X) = \sum_{x=1}^{10} \frac{1}{(11-x)/10} = 29.29;$$

$$\text{(b)} \quad \frac{100 \cdot E(X)}{365} \approx 8 \text{ years.}$$

2.7 The Poisson Distribution

$$\mathbf{2.7-2} \quad \lambda = \mu = \sigma^2 = 3 \text{ so } P(X = 2) = 0.423 - 0.199 = 0.224.$$

$$\begin{aligned}
\mathbf{2.7-4} \quad 3 \frac{\lambda^1 e^{-\lambda}}{1!} &= \frac{\lambda^2 e^{-\lambda}}{2!} \\
e^{-\lambda} \lambda (\lambda - 6) &= 0 \\
\lambda &= 6
\end{aligned}$$

$$\text{Thus } P(X = 4) = 0.285 - 0.151 = 0.134.$$

$$\begin{aligned}
\mathbf{2.7-6} \quad M(t) &= e^{\lambda(e^t-1)}, \\
M'(t) &= e^{\lambda(e^t-1)} \lambda e^t, \\
M''(t) &= e^{\lambda(e^t-1)} \lambda^2 e^{2t} + e^{\lambda(e^t-1)} \lambda e^t, \\
M'''(t) &= e^{\lambda(e^t-1)} \lambda^3 e^{3t} + e^{\lambda(e^t-1)} \lambda^2 \cdot 2e^{2t} + e^{\lambda(e^t-1)} \lambda^2 e^{2t} + e^{\lambda(e^t-1)} \lambda e^t, \\
E(X^3) &= M'''(0) = \lambda^3 + 3\lambda^2 + \lambda, \\
\gamma &= [\lambda^3 + 3\lambda^2 + \lambda - \lambda^3 - 3\lambda(\lambda + \lambda^2) + 3\lambda^2 \cdot \lambda] / \lambda^{3/2} = 1/\sqrt{\lambda}.
\end{aligned}$$

Thus the index of skewness is positive. As λ increases, the index of skewness decreases to zero.

$$\mathbf{2.7-8} \quad np = 1000(0.005) = 5,$$

$$\text{(a)} \quad P(X \leq 1) \approx 0.040;$$

$$\text{(b)} \quad P(X = 4, 5, 6) = P(X \leq 6) - P(X \leq 3) \approx 0.762 - 0.265 = 0.497.$$

$$\mathbf{2.7-10} \quad \sigma = \sqrt{9} = 3,$$

$$P(3 < X < 15) = P(X \leq 14) - P(X \leq 3) = 0.959 - 0.021 = 0.938.$$

$$\mathbf{2.7-12} \quad \text{Since } E(X) = 0.2, \text{ the expected loss is } (0.2)(\$100,000) = \$2,000.$$